

WINADRYL EXPECTORANT SYRUP

Diphenhydramine HCl / Ammonium chloride (14mg/5ml / 135mg/5ml)

What is in the leaflet

1. What Winadryl Expectorant Syrup is used for
2. How Winadryl Expectorant works
3. Before you use Winadryl Expectorant Syrup
4. How to take Winadryl Expectorant Syrup
5. While you are using it
6. Side Effects
7. Storage and Disposal of Winadryl Expectorant Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Winadryl Expectorant Syrup is used for

Winadryl Expectorant Syrup is used for the relief of cough and congestion in the chest and nasal passages. It helps to loosen viscous mucous and spit out stubborn phlegm.

How Winadryl Expectorant Syrup works

Diphenhydramine works by blocking the body's natural substance (histamine) produced during an allergic reaction.

Ammonium chloride decreasing viscosity of sputum and presumably is easier to cough up.

Before you use Winadryl Expectorant Syrup

- When you must not take it

- Do not take this product if you are allergic (hypersensitivity) to diphenhydramine and ammonium or any other ingredients in it.
- Do not take if you have respiratory problem.
- Do not give this product to children under 2 years of age.

- Before you start to use it Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

Use with caution and follow the instructions of your doctor or pharmacist carefully for children aged 2 - 6 years.

Tell your doctor or pharmacist if you have liver or kidney problems.

Tell your doctor or pharmacist if you have liver or kidney problem.

- Taking others medicines

There are some medicines Winadryl Expectorant Syrup may interfere with each other. These include:

- Antimuscarinic (medicines indicated for incontinence).
- CNS depressants (including barbiturates, hypnotics, opioid anaygesics, anxiolytic sadatives and antipsychotics).
- β blockers (medicines indicated for high blood pressure, heart failure and irregular heartbeat).

How to take Winadryl Expectorant Syrup

- How much to take

➤ Adults and children above 12 years:
Give 1 to 2 teaspoons (5 to 10ml) every 4 to 6 hours.

➤ Children 6 to 12 years:
Give ½ to 1 teaspoon (2.5ml to 5ml) every 4 to 6 hours.

➤ Children 2 to 5 years:
Give ½ teaspoon (2.5ml) every 4 to 6 hours.

Do not exceed 6 doses in 24 hours unless advised by a doctor.

- When to take it

Use as directed by your doctor or pharmacist.

- How long to take it

Continue using Winadryl Syrup for the duration that the doctor recommends.

- If you forget to take it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Winadryl Expectorant Syrup

- Things you must do

Use your medicine exactly as your doctor

Tell all the doctors, dentists and pharmacists treating you that you are using Winadryl Syrup.

Tell your doctor immediately if you become pregnant while using this medication.

Take Windaryl Syrup only with a doctor's prescription.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Windaryl Syrup to anyone else, even if they have the same symptoms or condition as you. (For prescription only medicines)

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Avoid Alcohol drinks.

Side Effects

This product may cause some unwanted side effects such as:

- Drowsiness
- Dizziness
- Incoordination
- Headache
- Slowing of physical and mental activity
- Dry mouth
- Blurred vision
- Urinary difficulty or retention,
- Constipation
- Increased gastric reflux
- Nausea
- Vomiting
- Diarrhea
- Epigastric pain

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

WINADRYL EXPECTORANT SYRUP

Diphenhydramine HCl / Ammonium chloride (14mg/5ml / 135mg/5ml)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Date of revision

11/03/2021

Serial Number:

NPRA(R4/1)080321/00008

A.LWR-PIL.01

Storage and Disposal of Winadryl Expectorant Syrup
- Storage

- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.
- Keep out of reach of children.

- Disposal

Do not take this after the expiry date. An expiry date is printed on the label and box of the medicine.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description
- What it looks like

Dark brown coloured syrupy liquid with plum flavour.

- Ingredients
- Active Ingredients

Each 5ml contains Diphenhydramine HCl 14mg and Ammonium Chloride 135mg.

- Inactive Ingredients

Methyl hydroxybenzoate, propyl hydroxybenzoate, sugar, xanthan gum, liquid glucose, sodium citrate, citric acid, menthol, aniseed oil, liquid caramel, plum flavour, polyoxyl 40 hydrogenated castor oil, purified water.

- MAL Number:

MAL08010745AZ

Manufacturer and Product Registration Holder

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.