



This design is applicable for 90ml bulk pack.
A label affix on a bottle.
12 bottles packed in this plain white box.
No printed outer carton for 90ml.



This design is applicable for 120ml bulk pack.
A label affix on a bottle.
12 bottles packed in this plain white box.



Date: 25 Mar 2025

Title: Winadryl Expectorant Syrup - 100ml Box

Checked by	
Approved by	

Winadryl
Expectorant Syrup



100ml

Winadryl
Expectorant Syrup

Indications:
For the relief of cough and congestion in the chest and nasal passages. It helps to loosen viscous mucous and expectorate stubborn phlegm.

Dosage: For oral administration only.
Adults and children above 12 years:
Take 1 to 2 teaspoons (5 to 10ml) every 4 to 6 hours.
Children 6 to 12 years:
Give ½ to 1 teaspoon (2.5 to 5ml) every 4 to 6 hours.
Children 2 to 5 years:
Give ½ teaspoon (2.5ml) every 4 to 6 hours.
Do not exceed 6 doses in 24 hour.

Precautions:
If symptoms persist, please consult a doctor.
This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drinks.
Not to be used in children less than 2 years of age. To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

Storage:
Store below 30°C. Protect from light. Keep cap tightly closed.
Keep out of reach of children.
Jauhkan daripada capaian kanak-kanak.

Please refer enclosed leaflet for more information.



Free
Dosage
Cup

Winadryl
Expectorant Syrup

Name/姓名/
Nama

Date/日期/
Tarikh

Take/服食/
Ambil

ml/毫升

每日 times daily/次/
kali sehari

before food/饭后/
sebelum makan

after food/饭后/
selepas makan

Tanpa Alkohol

Winadryl
Expectorant Syrup

Indikasi:
Untuk melegakan batuk, kesesakan dada dan hidung tersumbat. Ia membantu mengurangkan kelikatan mukus dan mengeluarkan kahak.

Dos:
Orang dewasa dan kanak-kanak melebihi 12 tahun:
Ambil 1 hingga 2 camca teh (5 hingga 10ml) setiap 4 hingga 6 jam.
Kanak-kanak 6 hingga 12 tahun:
Bagi ½ hingga 1 camca teh (2.5 hingga 5ml) setiap 4 hingga 6 jam.
Kanak-kanak 2 hingga 5 tahun:
Bagi ½ camca teh (2.5ml) setiap 4 hingga 6 jam.
Jangan melebihi 6 dos dalam masa 24 jam.

Perhatian:
Jika simptom berlarutan, sila dapatkan nasihat doktor.

Sila rujuk sisip bungkusan untuk maklumat lanjut.

Each 5ml contains -
Diphenhydramine HCl 14mg
Ammonium Chloride 135mg

Preserved with Methyl Hydroxybenzoate 0.12%w/v and Propyl Hydroxybenzoate 0.012%w/v.

MAL08010745AZ BRU24063455P

Controlled
Medicine



9 557837 002321

A.LWR100-B.01

100ml

Winadryl
Expectorant Syrup

Relief of cough and
congestion in the
chest and nasal
passages

- loosen viscous mucous
- expectorate stubborn
phlegm

Antihistamine
Expectorant

Alcohol Free



Size: 54mm x 54mm x 120mm

C M Y K

Manufacturer and Product Registration Holder
WINWA MEDICAL SDN. BHD.
198101009677(75794-A)
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

BN :
MFG :
EXP :

This design is applicable for 100ml individual pack.
A label affix on a bottle.
A bottle packed in this printed outer box.

Checked by	
Approved by	



Free

■ C ■ M ■ Y ■ K

This design is applicable for 120ml individual pack.
A label affix on a bottle.
A bottle packed in this printed outer box.