

COLDTACT CAPSULE

Paracetamol / Chlorpheniramine Maleate (500mg / 2mg)

What is in the leaflet

1. What Coldtact Capsule is used for
2. How Coldtact Capsule works
3. Before you use Coldtact Capsule
4. How to take Coldtact Capsule
5. While you are using it
6. Side Effects
7. Storage and Disposal of Coldtact Capsule
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Coldtact Capsule is used for

Coldtact Capsule is used for the relief of common cold and allergy symptoms including fever, headache, body aches, sinus pain, runny nose, sneezing, itching and watery eyes.

How Coldtact Capsule works

Coldtact Capsule contains 2 active ingredients. Paracetamol belongs to a group of medicines called analgesic and antipyretic that helps to ease pain and reduce fever. Chlorpheniramine Maleate is an antihistamine used to relieve the symptoms of allergy, hay fever and common cold.

Before you use Coldtact Capsule

- When you must not take it

Do not take this product if you are allergic (hypersensitivity) to Paracetamol, Chlorpheniramine Maleate or any other ingredients in this product.

- Before you start to use it

Not to be used in children less than 2 years of age.

To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

Caution should be taken for individual with impaired kidney or liver functions or with alcohol dependence.

Caution should be taken for individual with eyes disease, gastric outlet obstruction, prostate enlargement or difficulty urinating.

- Taking other medicines

- Warfarin (prevents blood clots)
- Metoclopramide (helps control acid reflux)
- Phenytoin (Enzyme-inducer)
- Antimuscarinic drugs (reduces bladder muscle spasms)
- Propantheline (reduces gastric acid secretion)
- antidepressants with anticholinergic properties and narcotic analgesics.
- alcohol and anticonvulsant agents
- Probenecid (prevents gout)
- Chloramphenicol (antibiotic)
- Colestyramine (prevents reabsorption)
- Central nervous system depressants (e.g. alcohol, sedatives, opioid analgesics, hypnotics)
- Atropine, monoamine oxidase inhibitors (MAOIs) and tricyclic antidepressants (TCAs)

How to take Coldtact Capsule

- How much to take

Adults and children above 12 years:

Take 2 capsules every 4 to 6 hours. Do not exceed 8 capsules per day.

Children 7 to 12 years:

Take 1 capsule every 4 to 6 hours. Do not exceed 4 capsules per day.

Not recommended for children under 7 years.

- How long to take it

Take as directed by your doctor or pharmacist.

- If you forget to take it

If you forget to take Coldtact Capsule, give it as soon as you remember. Never double up on the next dose to make up for the one missed.

- If you take too much (overdose)

In case of overdose, prompt specific treatment is essential.

Call National Poison Centre (1800 888 099) or your doctor or go to the accident and emergency centre at your nearest hospital immediately.

While you are using Coldtact Capsule

- Things you must do

Consult a doctor if fever persists for more than 3 days or pain persists for more than 10 days or if there is redness.

Caution should be taken to individual with impaired kidney or liver functions or with alcohol dependence.

- Things you must not do

This preparation contains Paracetamol. Do not take any other Paracetamol containing medicines at the same time.

- Things to be careful of

This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drinks.

Side Effects

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Central nervous system depressive effects of Chlorpheniramine Maleate include sedation, lack of energy, dizziness and incoordination.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.moh.gov.my (Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Coldtact Capsule

- Storage

- Store below 30°C.
- Protect from light.
- Keep container tightly closed.
- Keep out of reach of children.

- Disposal

Do not take this after the expiry date. An expiry date is printed on the label of the medicine.

Product Description

- How it looks like

Coldtact Capsule is a red and natural transparent size # 0 hard gelatin capsule containing tri-coloured granules of white, orange and red.

COLDTACT CAPSULE

Paracetamol / Chlorpheniramine Maleate (500mg / 2mg)

Ingredients

- Active Ingredients

Each capsule contains Paracetamol 500mg,
Chlorpheniramine Maleate 2mg

- Inactive Ingredients

Povidone K30, Quinoline Yellow, Sunset
Yellow, Ponceau 4R

- MAL Number:

MAL08051763AZ

**Manufacturer and Product Registration
Holder**

Winwa Medical Sdn Bhd
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

26/07/2022

Serial Number:

NPRA(R4/1)190719/00198

X.CMD-PIL.01