

CALPO SUSPENSION (Anthelmintic)	CALPO SUSPENSION (Ubat Cacing)
Active Ingredient: Each 5ml contains Mebendazole 100mg Preservatives - Methyl Hydroxybenzoate 0.1%w/v Propyl Hydroxybenzoate 0.01%w/v Sodium Benzoate 0.1%w/v Presentations: Bottles of 10ml and 25ml. Product Descriptions: Creamy white suspension. Pharmacology: Mebendazole is an anthelmintic against Threadworm (Enterobius), Roundworm (Ascaris), Hookworm (Ancylostoma and Necator) and Whipworm (Trichuris). The mechanism of action is probably as a result of an inhibitory effects on the ability of the organism to utilise exogenous glucose. This interferes with the generation of phosphate bonds (ATP), ultimately leading to the death of the parasite. Calpo Suspension when taken by mouth is poorly absorbed. It remains in the intestine and inhibits the metabolic action of the worms. This ultimately leads to the death of the parasite. Poorly absorbed after oral administration. Metabolised to the 5-(-hydroxy) derivative and by decarboxylation to the 2-amino metabolite, both of which are detectable in plasma at concentrations higher than those of unchanged Mebendazole. Less than 10% of a dose is excreted in the urine; the major urine metabolite is the 2-amino-5-(-hydroxy) derivative. Biliary excretion and enterohepatic circulation haven been reported. Indications: For the treatment of single or mixed worm infestations, including threadworm, roundworm, whipworm and hookworm. Dosage and Administration: Adults and children above 2 years: Threadworm Infestation: 1 teaspoon (5ml) as a single dose. May be repeated after 2 to 3 weeks. Roundworm, Hookworm and Whipworm Infestation: 1 teaspoon (5ml) twice a day for 3 consecutive days. OR 1 bottle (25ml) as a single dose. Shake well before use. Symptoms and Treatments of Overdose: In case of accidental poisoning, please seek professional assistance from your doctor. Symptoms: Persistent gastrointestinal upset and diarrhoea. Treatment: Supportive therapy be initiated. Antidiarrhoeal treatment might be considered if diarrhoea is excessive. Contraindications: Not suitable for people with known hypersensitivity to the drug or other Benzimidazole derivatives, pregnant women and children under 2 years. Precautions: Use during lactation: May pass into maternal milk. Therefore use in nursing mothers requires that anticipated benefits be weighed against possible risks. Undesirable Effects: Nausea, vomiting, abdominal pain and diarrhoea may occasionally occur. Storage: Store below 30°C. Protect from light. Keep cap tightly closed. Keep out of reach of children. Shelf Life: 3 years from the date of manufacture. Manufacturer and Product Registration Holder WINWA MEDICAL SDN. BHD. 1952, Lorong 1, K.S. Bukit Minyak 3, Taman 1, K.S. Bukit Minyak, 14000 Bukit Mertajam, Pulau Pinang, Malaysia. MAL19912835X BRU16082065P Date of Revision: 1 March 2021 X.LCP-I.07	Saiz Bungkusan: Botol 10ml dan 25ml. Deskripsi Produk: Ampaian putih kekuningan. Farmakologi: Ubat Cacing Calpo bila diambil, tidak diserap ke dalam badan. Ia bertindak di usus dengan menindas keupayaan cacing mendapat sumber tenaga, jadi mengakibatkan kematian cacing dalam badan dan kemudian disingkirkan melalui najis. Indikasi: Ubat Cacing Calpo dapat menghapuskan jangkitan cacing-cacing biasa yang berada dalam usus, seperti Cacing Kerawit, Cacing Gelang, Cacing Kait dan Cacing Cambuk. Dos: Dewasa dan kanak-kanak 2 tahun ke atas: Cacing Kerawit: 1 camca teh (5ml) diambil sekali sahaja. Diulangi jika perlu selepas 2 atau 3 minggu. Cacing Gelang, Cacing Kait dan Cacing Cambuk: 1 camca teh (5ml) 2 kali sehari, diambil 3 hari berturutan. ATAU 1 botol (25ml) sebagai dos tunggal. Goncangkan botol sebelum diambil. Kontraindikasi: Tidak sesuai untuk sesiapa yang hipersensitif pada Mebendazole atau ubat terbitan Benzimidazole, wanita yang hamil dan kanak-kanak di bawah umur 2 tahun. Perhatian: Semasa Penyusuan: Berkeupayaan melintasi susu ibu. Penggunaan oleh ibu yang menyusu perlu dipertimbangkan terhadap kebaikan dan risiko. Kesan Sampingan: Mual, muntah, sakit abdomen dan cirit-birit. Cara Penyimpanan: Simpan di bawah 30°C. Jauhkan daripada cahaya. Tutupkan penutup dengan ketat. Jauhkan daripada capaian kanak-kanak. 健儿宝疳积露 包装: 10 及 25 毫升瓶装。 主治: 健儿宝疳积露能消灭体内之寄生虫包括蛲虫、蛔虫、钩虫及鞭虫。 服法: 成人及儿童 2 岁以上: 蛲虫: 每次服 1 茶匙 (5ml), 2 至 3 个星期后可重复服用。 蛔虫, 钩虫及鞭虫: 每次服 1 茶匙 (5ml), 每日服 2 次, 连续服 3 天。 或 服 1 瓶 (25ml), 1 次即可。 服用前, 先摇匀。 注意: 孕妇忌服。 贮存: 置于摄氏 30 度以下。避光。确保把盖关紧。储存在远离孩童的地方。

Size: 55mm x 120mm

Dark Blue (Pantone 2767U)