

MIFULIN OINTMENT

Salicylic acid / Zinc oxide (12.5% w/w / 6% w/w)

What is in this leaflet

1. What Mifulin Ointment is used for
2. How Mifulin Ointment works
3. Before you use Mifulin Ointment
4. How to use Mifulin Ointment
5. While you are using it
6. Side effects
7. Storage and disposal of Mifulin Ointment
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Mifulin Ointment is used for

Mifulin Ointment is used to treat skin conditions like flaky or greasy patches (seborrhoeic dermatitis), red and scaly skin (psoriasis), very dry and thickened skin (ichthyosis), as well as warts, and areas of hard or thickened skin like corns and calluses.

How Mifulin Ointment works

Mifulin Ointment contains the active ingredients salicylic acid and zinc oxide. Salicylic acid helps by softening and removing dead skin cells, making it easier to get rid of rough patches, warts, and thickened skin. Zinc oxide soothes and protects the skin, reducing irritation and helping it heal.

Before you use Mifulin Ointment

- When you must not use it

Do not use Mifulin Ointment if:

- you are allergic to salicylic acid, zinc oxide or any other ingredients of this medicine (listed in Product description)
- your child is under 2 years of age.

- Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Inform your doctor or pharmacist if you are taking pharmaceutical products, including any that you get without a prescription from your pharmacy, supermarket, or health food store.

How to use Mifulin Ointment

Clean and dry the affected areas. Apply Mifulin Ointment to the affected area and not to the surrounding healthy skin.

- How much to use

Apply a thin film of Mifulin Ointment directly to the affected area.

- When to use it

Use it two times a day.

- How long to use it

Use this medicine for the duration that your doctor recommends.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much salicylic acid can cause symptoms like dizziness, ringing in the ears, hearing loss, sweating, headache, nausea, vomiting, fast breathing, restlessness, and confusion. It can also affect your body's acid levels and brain function.

While you are using it

- Things you must do

Use your medicine exactly as your doctor or pharmacist has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Mifulin Ointment.

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Mifulin Ointment to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

As Mifulin Ointment contains camphor, avoid direct application into the nostrils, especially in children. A small amount applied in this way may cause immediate collapse. Caution must be exercised when older children are treated with this medicine.

Avoid contact with eyes or mucous membrane.

If any irritation occurs, or your skin condition worsens or does not improve, please see your doctor again immediately.

Be careful when using on large skin areas.

Side effects

Like all medicines, Mifulin Ointment can cause side effects, although not everybody gets them.

This product may cause skin irritation (dermatitis) such as redness, itchiness, or inflamed skin.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Mifulin Ointment

- Storage

- Keep out of the reach and sight of children.
- Store below 30°C.
- Protect from light.
- Keep container tightly closed.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A white smooth ointment.

- Ingredients

- Active ingredient
Salicylic acid, zinc oxide.
- Inactive ingredients
Liquid paraffin, white soft paraffin, menthol crystal, camphor.

- MAL number:

MAL19912830X

MIFULIN OINTMENT

Salicylic acid / Zinc oxide (12.5% w/w / 6% w/w)

**Manufacturer and Product
Registration Holder**

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of Revision

09/05/2025

Serial Number

NPRA(R4/1)280425/00030