

MIFULIN LOTION

Salicylic acid (12% w/v)

What is in this leaflet

1. What Mifulin Lotion is used for
2. How Mifulin Lotion works
3. Before you use Mifulin Lotion
4. How to use Mifulin Lotion
5. While you are using it
6. Side effects
7. Storage and Disposal of Mifulin Lotion
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Mifulin Lotion is used for

Mifulin Lotion is used for the treatment of corns, warts, and ringworm.

How Mifulin Lotion works

Mifulin Lotion contains Salicylic acid, which helps control bacteria, fungi and removes dead skin. It works by softening keratin, a protein in your skin, making it easier to remove dry and scaly skin.

Before you use Mifulin Lotion

- When you must not use it

Do not use Mifulin Lotion if:

- you are allergic to Salicylic acid or any other ingredients of this medicine (listed in Product Description)
- if you have wounds or damaged skin
- patients are an infant below 2 years of age as it may cause convulsions.

- Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Inform your doctor or pharmacist if you are taking pharmaceutical products, including any that you get without a prescription from your pharmacy, supermarket, or health food store.

How to use Mifulin Lotion

Clean and dry the affected areas. Apply Mifulin Lotion to the affected area and not to the surrounding healthy skin.

- How much to use

Apply a thin film of Mifulin Lotion directly to the affected area.

- When to use it

Use it once daily.

- How long to use it

Use until the corn/wart/ringworm has completely cleared.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Salicylate intoxication can occur after accidental ingestion or following excessive topical application of salicylate.

You may experience symptoms of discomfort such as dizziness, ringing in the ears, hearing loss, sweating, headache, nausea, vomiting, rapid breathing, restlessness and confusion.

Discontinue use and contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Use your medicine exactly as your doctor or pharmacist has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Mifulin Lotion.

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not stop using the medicine unless advised by your doctor or pharmacist.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Mifulin Lotion to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Mifulin Lotion contains Camphor and it can cause convulsions. Caution must be exercised when older children are treated.

Avoid direct application into the nostrils. It is dangerous to place any camphor-containing product into the nostrils of children. A small amount applied in this way may cause immediate collapse.

Avoid contact with eyes or mucous membrane. If any irritation occurs, discontinue use.

Be careful driving or operating machinery until you know how Mifulin Lotion affects you.

Side effects

Like all medicines, Mifulin Lotion can cause side effects, although not everybody gets them.

This product may cause skin irritation (dermatitis) such as redness, itchiness or inflamed skin.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Mifulin Lotion

- Storage

- Keep out of the reach and sight of children.
- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

MIFULIN LOTION

Salicylic acid (12% w/v)

Product Description

- What it looks like

A clear, colourless solution with thymol and camphor odour.

- Ingredients

- Active ingredient
Salicylic acid
- Inactive ingredients
Methylated spirit, thymol, camphor, purified water.

- MAL number:

MAL19912812XZ

Manufacturer and Product

Registration Holder

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of Revision

17/12/2024

Serial Number

NPRA(R4/1)061224/00126