

M-ZOLE CHEWABLE TABLET

Mebendazole (500mg)

What is in the leaflet

1. What M-zole Chewable Tablet is used for
2. How M-zole Chewable Tablet works
3. Before you use M-zole Chewable Tablet
4. How to take M-zole Chewable Tablet
5. While you are using it
6. Side Effects
7. Storage and Disposal of M-zole Chewable Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What M-zole Chewable Tablet is used for

M-zole Chewable Tablet is a single dose treatment against single or mixed intestinal worm infections.

How M-zole Chewable Tablet works

M-zole Chewable Tablet contains mebendazole which works by stopping the worms from using glucose sugar and leading to the death of the worms.

Before you use M-zole Chewable Tablet

- When you must not take it

Do not take this product if you are allergic to mebendazole, similar medicines or any other ingredients in the product.

- Before you start to use it

This product is unsuitable for phenylketonurics.

Not recommended to be used during pregnancy or whilst breastfeeding.

- Taking other medicines

Tell your doctor if you are taking medicines such as dexamethasone (relieve inflammation), praziquantel (treat infections) and cimetidine (relieve heartburn).

How to take M-zole Chewable Tablet

- How much to take

For adults and children above 2 years old: Take 1 tablet as a single dose.

Can be chewed, crushed or mixed with juice or food.

- When to take it

Take as a single dose after food or directed by your doctor or pharmacist.

- How long to take it

A second dose can be taken after 2 to 3 weeks if reinfestation occurs.

- If you forget to take it

If you forget to take M-zole Chewable Tablet, take it as soon as you remember. Do not take a double dose to make up for the missed dose.

- If you take too much (overdose)

If you think you or anyone else may have taken too much of this medicine, contact your doctor immediately or go to the Emergency Department of your nearest hospital. Do this even if there are no sign of discomfort or poisoning. You may need urgent medical attention. Taking too much medicine may cause stomach and bowel disturbances and diarrhoea.

While you are using M-zole Chewable Tablet

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell your doctor immediately if you become pregnant while taking this medication.

If symptoms persist, please consult a doctor.

- Things you must not do

Do not take any new medicines without consulting your doctor.

- Things to be careful of

All family members in close contact should be treated at the same time to prevent reinfestation.

Side Effects

Side effects including stomach and bowel disturbances, diarrhoea, headache and dizziness may occasionally occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website nptra.gov.my [Consumers Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of M-zole Chewable Tablet

- Storage

- Store below 30°C.
- Protect from light, heat and moisture.
- Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Light yellow coloured, round and flat uncoated tablet with banana and vanilla flavour.

Ingredients

- Active ingredient
Mebendazole

- Inactive Ingredients

Isomalt, microcrystalline cellulose, aspartame, colloidal silicon dioxide, precipitated silicon dioxide, banana flavour, vanilla flavour, quinoline yellow lake, magnesium stearate.

- MAL Number:

MAL06061546X

Manufacturer and Product Registration Holder

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

22/02/2023

Serial Number:

NPRA(R4/I)150223/00013

X.TMZ-PIL.01