

RIDSIL OINTMENT

Terbinafine Hydrochloride (1% w/w)

What is in this leaflet

1. What Ridsil Ointment is used for
2. How Ridsil Ointment works
3. Before you use Ridsil Ointment
4. How to use Ridsil Ointment
5. While you are using it
6. Side effects
7. Storage and Disposal of Ridsil Ointment
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Ridsil Ointment is used for

It is used to treat fungal infections of the skin and nails, including the groin area.

How Ridsil Ointment works

Ridsil Ointment contains the active ingredient terbinafine hydrochloride, which is an antifungal medication. Terbinafine works by killing the fungi that cause infections.

Before you use Ridsil Ointment

- When you must not use it

Do not use Ridsil Ointment if:

- you are allergic to terbinafine hydrochloride or any other ingredients of this medicine (listed in Product Description);

- Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

Terbinafine must only be used during pregnancy if the benefits are more than the risks to the fetus.

Terbinafine is secreted into breast milk, therefore its use in nursing mothers is not recommended, unless the benefits are more than the risks.

- Taking other medicines

Inform your doctor or pharmacist before using this medicine, including any that you get without a prescription from your pharmacy, supermarket, or health food store.

Please tell your doctor or pharmacist if you are currently using any topical medication.

How to use Ridsil Ointment

Always use this medicine exactly as your doctor or pharmacist has told you.

You should check with your doctor or pharmacist if you are not sure.

- How much to use

Clean and dry the infected area thoroughly with warm water.

Apply a thin layer onto the infected area. Rub in gently. Wash hands after every use.

- When to use it

Apply the ointment once daily or as directed by your doctor or pharmacist.

- How long to use it

Apply the medicine for the duration as the doctor recommends.

The treatment period is usually between 1-2 weeks. Even after symptoms are gone, keep treating the area for another 1 to 2 weeks to prevent the infection from coming back. You may repeat the treatment if needed.

If your infection worsens or no improvement seen after using for 1 week, please see your doctor again.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much of this ointment may cause more severe or increased side effects. In case of accidental ingestion, it may cause gastrointestinal symptoms such as nausea and vomiting.

While you are using it

- Things you must do

Tell all the doctors and pharmacists treating you that you are using Ridsil Ointment.

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Ridsil Ointment to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

For external use only. Avoid contact with eyes, nose and mouth.

Be careful when it is used on skin lesions because it contains alcohol, which can irritate.

If you notice any allergy or skin reactions while using this ointment, stop using and see doctor immediately.

Side effects

Like all medicines, Ridsil Ointment can cause side effects, although not everybody gets them.

Occasionally, itching or redness may occur.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Ridsil Ointment

- Storage

- Keep out of the reach and sight of children.
- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White to off-white ointment.

- Ingredients

- Active ingredient
Terbinafine Hydrochloride

RIDSIL OINTMENT

Terbinafine Hydrochloride (1% w/w)

- Inactive ingredients

Cetomacrogol Emulsifying Wax,
Hard Paraffin Wax, Cetostearyl
Alcohol, White Soft Paraffin,
Lamellar Gel System, PEG-40
Hydrogenated Castor Oil,
Hydrogenated Polyisobutene,
Butylated Hydroxytoluene,
Isopropyl Myristate, Propylene
Glycol, N-Methylpyrrolidone,
Benzyl Alcohol, Menthol.

- MAL number:

MAL08021517X

Manufacturer and Product

Registration Holder

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of Revision

22/08/2024

Serial Number

NPRA(R4/1)190824/00108