

BENZ-C DOUBLE ACTION LOZENGE

Benzylamine HCl / Cetylpyridinium chloride (3mg / 1.33mg)

What is in the leaflet

1. What Benz-C Double Action Lozenge is used for
2. How Benz-C Double Action Lozenge works
3. Before you use Benz-C Double Action Lozenge
4. How to use Benz-C Double Action Lozenge
5. While you are using it
6. Side Effects
7. Storage and Disposal of Benz-C Double Action Lozenge
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What Benz-C Double Action Lozenge is used for

Benz-C Double Action Lozenge is used to relieve inflammation and pain in the mouth, denture ulcers and sore gums. Cetylpyridinium Chloride has antiseptic activities.

How Benz-C Double Action Lozenge works

Benz-C Double Action Lozenge works by reducing the pain, redness and swelling.

Before you use Benz-C Double Action Lozenge

- When you must not use it

Do not use this product if you are allergic to Benzylamine HCl, Cetylpyridinium Chloride or any other ingredients in this product.

Do not use this medicine if you are pregnant or plan to become pregnant. If you are not sure whether you should start using this medicine, talk to your pharmacist or doctor.

- Before you start to use it

Tell your pharmacist if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your pharmacist if you have impaired kidney or liver function.

It is not known if Benzylamine HCl or Cetylpyridinium Chloride passes into the breast milk.

Your pharmacist or doctor will discuss the potential benefits and risks of using the medicine if you are breastfeeding.

- Taking other medicines

Tell your pharmacist or doctor if you are taking any other medicines, including any that you get without a prescription or that are herbal from your pharmacy, supermarket or health food shop.

Your pharmacist or doctor will have more information on medicines to be careful with or avoid while using this medicine.

How to use Benz-C Double Action Lozenge

- How much to use

For oral use only. The lozenges should not be chewed. One lozenge should be sucked slowly every 1 to 2 hours as required up to a maximum of 12 lozenges per day.

Children under 6 years: Not recommended to use.

The use of Benz-C Double Action Lozenge should be considered in patients with renal impairment as well as liver disorders.

- When to use it

Use it every 1 to 2 hours as required or when necessary.

- How long to use it

Uninterrupted treatment should not exceed 7 days.

- If you forget to use it

If you forget to use Benz-C Double Action Lozenge, use it as soon as you remember. Never double up on the next dose to make up for the one missed.

- If you use too much (overdose)

Immediately telephone your doctor for advice, or go to the Accident and Emergency at the nearest hospital. If you think that you or anyone else may have use or accidental swallow too much Benz-C Double Action Lozenge, do this even if there are no signs of discomfort or poisoning.

While you are using Benz-C Double Action Lozenge

- Things you must do

Your pharmacist or doctor will assess your condition and decide if you should continue to use the medicine.

- Things you must not do

Do not use this product to treat any other conditions unless it has been prescribed by your doctor.

- Things to be careful of

If your condition does not improve, please see your doctor or pharmacist for further advice.

Side Effects

Some unwanted effects such as oral numbness, burning and stinging sensation, mouth dryness and thirst have been observed.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Benz-C Double Action Lozenge

- Storage

- Store below 30°C.
- Protect from light and moisture.
- Keep out of reach of children.

- Disposal

Do not use Benz-C Double Action Lozenge after the expiry date which is stated on the carton. Medicines should not be disposed off via wastewater or household waste. Ask your pharmacist how to dispose medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Benz-C Double Action Lozenge is an orange coloured lozenge, round and flat on both sides with fruity flavour. It is also sugar free.

BENZ-C DOUBLE ACTION LOZENGE

Benzydamine HCl / Cetylpyridinium chloride (3mg / 1.33mg)

- Ingredients

- Active Ingredients

Benzydamine HCl 3mg

Cetylpyridinium chloride 1.33mg

- Inactive Ingredients

Isomalt, copovidone, microcrystalline cellulose, ammonium glycyrrhizinate, sucralose, colloidal silicon dioxide, sodium citrate dihydrate, precipitated silicon dioxide, menthol, orange flavour, aniseed oil, mixed fruit flavour powder, sunset yellow lake, magnesium stearate.

- MAL Number:

MAL21036019AZ

**Manufacturer and Product
Registration Holder**

Winwa Medical Sdn Bhd
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

17/05/2023

Serial Number:

NPRA(R2)20/124

A.TBC-PIL.01