

HOCODIN LOZENGE

Dequalinium chloride (250mcg)

What is in the leaflet

1. What Hocodin Lozenge is used for
2. How Hocodin Lozenge works
3. Before you use Hocodin Lozenge
4. How to take Hocodin Lozenge
5. While you are using it
6. Side effects
7. Storage and disposal of Hocodin Lozenge
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Hocodin Lozenge is used for

Hocodin Lozenge is indicated for the relief of minor infections of the mouth and throat.

How Hocodin Lozenge works

Hocodin Lozenge contains dequalinium chloride as an active ingredient. dequalinium chloride has antiseptic effect and works by killing a wide range of bacteria, some fungi, and viruses.

Before you use Hocodin Lozenge

– When you must not take it

Do not take Hocodin Lozenge if you are:

- hypersensitive to dequalinium chloride or any other ingredients of this medicine (listed in Product Description).

– Before you start to use it

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

Tell your doctor if you are known to have phenylketonuria (a rare genetic disease).

– Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket, or health food shop.

How to take Hocodin Lozenge

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

– How much to take

Suck 1 lozenge slowly every 3 hours or when necessary. Maximum 8 lozenges per day.

– When to take it

Use as directed by your doctor or pharmacist.

– How long to take it

Use this medicine for the duration that your doctor recommends.

– If you forget to take it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

– If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Hocodin Lozenge

– Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Hocodin Lozenge.

Tell your doctor immediately if you become pregnant while using this medication.

– Things you must not do

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Hocodin Lozenge to anyone else, even if they have the same symptoms or condition as you.

– Things to be careful of

Be careful driving or operating machinery until you know how Hocodin Lozenge affects you.

Side Effects

Side effects and undesirable symptoms of the Hocodin Lozenge have not been observed.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Hocodin Lozenge

– Storage

- Store below 30°C.
- Protect from light, heat, and moisture.
- Keep out of reach of children.

– Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

– What it looks like

Peppermint flavoured, round green lozenge with 'W' on one side and score on the other.

– Ingredients

- Active ingredient
Dequalinium chloride
- Inactive Ingredients
Isomalt, ammonium glycyrrhizinate, aspartame, sodium citrate dihydrate, colloidal silicon dioxide, precipitated silicon dioxide, menthol, peppermint oil, aniseed oil, orange durarome, brilliant blue lake, quinoline yellow lake, magnesium stearate.

– MAL Number:

MAL11075007X

Manufacturer and Product Registration Holder

Winwa Medical Sdn. Bhd.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

05/12/2025

Serial Number:

NPRA(R4/1)310725/00060