

WINDEXCO PE SYRUP

Dextromethorphan HBr / Phenylephrine HCl / Chlorpheniramine Maleate (10mg/5ml / 5mg/5ml / 2mg/5ml)

What is in the leaflet

1. What Windexco PE Syrup is used for
2. How Windexco PE Syrup works
3. Before you use Windexco PE Syrup
4. How to take Windexco PE Syrup
5. While you are using it
6. Side Effects
7. Storage and Disposal of Windexco PE Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Windexco PE Syrup is used for

Windexco PE Syrup is indicated for cough and cold relief such as dry irritating cough, runny nose and blocked nose. It also helps to relieve sneezing, itchy and watery eyes associated with common cold, allergy and other respiratory infection.

How Windexco PE Syrup works

Windexco PE Syrup contains 3 active ingredients. Dextromethorphan is an antitussive (cough suppressant) drug of the morphinan class. It suppresses the urge to cough by dulling the sensitivity of the nerves that control the cough reflex. Phenylephrine acts as a decongestant by a direct effect on α -adrenergic receptors. Chlorpheniramine is a histamine antagonist of the H1 receptor that serves to reduce effects mediated by histamine.

Before you use Windexco PE Syrup

- When you must not take it

Do not take this product if you are with severe hypertension, ventricular tachycardia, peripheral or mesenteric vascular thrombosis. Do not take if you are allergic (hypersensitivity) to Dextromethorphan HBr, Phenylephrine HCl, Chlorpheniramine Maleate or any other ingredients in this product. This product should not be taken for persistent or chronic cough and patients receiving MAOI or for 2 weeks after discontinuing them. It should not be used in premature or full-term neonates.

- Before you start to use it

This product should be used with caution in atopic children, sedated or debilitated patients and in patients confined to the supine position. Caution should be taken in patients with angle-closure glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction or bladder neck obstruction.

- Taking other medicines

Should not take this product if you are taking any of these medicines:

Tricyclic antidepressants (TCAs), antipsychotics, anxiolytics and hypnotics, cimetidine, ciprofloxacin, domperidone, metoclopramide, mexiletine, CYP2D6 inhibitors, ritonavir, alcohol.

How to take Windexco PE Syrup

- How much to take

Adults and children above 12 years:

Take 10ml every 6 hours.

Children 6 to 12 years:

Take 5ml every 6 hours.

Children 2 to 5 years:

Take 2.5ml every 6 hours.

Do not exceed 4 doses in 24 hours.

Do not give more than the recommended dose unless advised by a doctor.

- When to take it

To be taken with food.

- How long to take it

Take as directed by your doctor or pharmacist.

- If you forget to take it

If you forget to take Windexco PE Syrup, take it as soon as you remember. Never double up on the next dose to make up for the one missed.

- If you take too much (overdose)

In case of overdose, call National Poison Centre (1800 888 099) or your doctor or go to the accident and emergency centre at your nearest hospital immediately.

While you are using Windexco PE Syrup

- Things you must do

Please consult your pharmacist / doctor if symptoms persist / worsen.

- Things you must not do

- Not to be used in children less than 2 years of age
- To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

- Things to be careful of

This medicine may cause drowsiness and may increase the effects of alcohol. If affected, do not drive a motor vehicle or operate machinery. Avoid alcoholic drinks.

Side Effects

Nausea, epigastric distress, anorexia, vomiting, diarrhea, constipation, other GI disturbances, slight drowsiness and

dizziness, restlessness, anxiety, nervousness, weakness, precordial pain, tremor, respiratory distress, pallor or blanching of the skin or a pilomotor response, sedation, hypotension may occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Windexco PE Syrup

- Storage

- Store below 30°C.
- Protect from light.
- Keep cap tightly closed.
- Keep out of reach of children.

- Disposal

Do not take this after the expiry date. An expiry date is printed on the label and box of the medicine.

Product Description

- How it looks like

Dark brown coloured syrupy solution with a pleasant taste. Alcohol free.

Ingredients

- Active Ingredients

Each 5ml contains Dextromethorphan HBr 10mg, Phenylephrine HCl 5mg, Chlorpheniramine Maleate 2mg.

- Inactive Ingredients

Methyl Hydroxybenzoate, Propyl Hydroxybenzoate, Sodium Saccharin, Sucralose, Xanthan Gum, Sorbitol Powder, Liquid Glucose, Sodium Citrate, Menthol, Peppermint Oil, Aniseed Oil, Hydrogenated Castor Oil, Liquid Caramel, Purified Water.

- MAL Number:

MAL16015070AZ

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

14/01/2021

Serial Number:

NPRA(R2/1)201015/533