

ORALIS PAIN RELIEF GEL

Lidocaine Hydrochloride 0.33%w/w, Cetylpyridinium Chloride 0.1%w/w

What is in the leaflet

1. What Oral Pain Relief Gel is used for
2. How Oral Pain Relief Gel works
3. Before you use Oral Pain Relief Gel
4. How to take Oral Pain Relief Gel
5. While you are using it
6. Side Effects
7. Storage and Disposal of Oral Pain Relief Gel
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Oral Pain Relief Gel is used for

Oral Pain Relief Gel is indicated to relieve pain and soothe the gum.

How Oral Pain Relief Gel works

Lidocaine hydrochloride is a local anaesthetic which works by stopping the sensation of pain. Cetylpyridinium chloride is an antiseptic which has bactericidal activity.

Before you use Oral Pain Relief Gel

- When you must not take it

Do not take this product if you are hypersensitivity to Lidocaine Hydrochloride, Cetylpyridinium Chloride and its derivatives. Talk to your pharmacist or doctor if you are not sure whether you should start taking this medicine.

- Before you start to use it

Tell your pharmacist or doctor if you have allergies to any other medicines.

- Taking other medicines

Do not use any other product on the affected area without consulting your doctor.

How to take Oral Pain Relief Gel

- How much to take

Place a small quantity of **Oral Pain Relief Gel** on a pad of cotton wool or a clean fingertip and rub gently on the gum. Repeat after 20 minutes if necessary. Use when necessary during the teething period.

- When to take it

You should follow the recommended dose and should not take more than the prescribed dose.

- How long to take it

Take as directed by your doctor or pharmacist.

- If you forget to take it

If you missed a dose, apply the next dose as soon as you remember. Do not double the amount applied than needed.

- If you take too much (overdose)

In case you accidentally ingest too much of Oral Pain Relief Gel, contact your doctor immediately or your nearest hospital for medical advice.

While you are using Oral Pain Relief Gel

- Things you must do

Check with your doctor if symptoms persist or getting worse.

- Things you must not do

Do not take this product to treat any other conditions unless it has been prescribed by your doctor.

- Things to be careful of

Never leave the medicine on counters, nightstands or tables where reachable by children.

Side Effects

None known.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers → Reporting).

Storage and Disposal of Oral Pain Relief Gel

- Storage

- Store below 30°C.
- Protect from light and moisture.
- Keep out of reach of children.

Disposal

Do not take this after the expiry date. An expiry date is printed on the label and box of the medicine.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- How it looks like

Oral Pain Relief Gel is a transparent and colourless gel with orange flavour.

- Ingredients

Active Ingredients

Lidocaine Hydrochloride 0.33%w/w, Cetylpyridinium Chloride 0.1%w/w.

Inactive Ingredients

Poloxamer 407, Carbomer, Sucralose, Propylene Glycol, Orange Flavour, 1M NaOH Solution, Purified Water.

- MAL Number:

MAL00000000AZ

Manufacturer and Product Registration Holder

WINWA MEDICAL SDN. BHD.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Date of revision

7 July 2017

A.IOP-PIL.01