

BUSPAN TABLET

Hyoscine butylbromide (10mg)

What is in this leaflet

1. What Buspan Tablet is used for
2. How Buspan Tablet works
3. Before you use Buspan Tablet
4. How to use Buspan Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Buspan Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Buspan Tablet is used for

Buspan Tablet is used to relieve cramps in the stomach and bowel. It is also used to help relax gallbladder and kidney spasms.

How Buspan Tablet works

Buspan Tablet contains a medicine called "Hyoscine butylbromide" which works by relaxing the muscle found in the walls of the organs.

Before you use Buspan Tablet

- When you must not use it

Do not take Buspan Tablet if:

- you are allergic (hypersensitive) to Hyoscine butylbromide or other ingredients listed at the end of the leaflet

If you are not sure, talk to your doctor or pharmacist before taking this medicine.

- Before you start to use it

You should check with your doctor or pharmacist before taking Buspan Tablet if you:

- have heart or lung disease
- have acid reflux
- have a condition where the bowel is blocked and does not work properly
- have glaucoma (increased pressure in the eye)
- have bleeding with unstable heart condition
- have high blood pressure or high blood pressure during pregnancy
- are allergic to similar medicines
- have urinary retention

- have overactive thyroid
- have dry mouth
- have kidney or liver function impairment
- are pregnant, breast-feeding or plan to start breast-feeding.
- have ulcers (sores) in your digestive tract or enlarged bowel

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Buspan Tablet.

- Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription and herbal medicines. This is because Buspan can affect the way some other medicines work. Also some other medicines can affect the way Buspan work.

In particular, tell your doctor or pharmacist if you are taking any of the following:

- Urine alkalizer: antacids, citrates, sodium bicarbonate
- Medicines for psychotic disorders: Haloperidol
- Medicines for diarrhea
- Medicines for nausea and vomiting: Metoclopramide
- Antifungal: Ketoconazole
- Opioid painkiller
- Medicine for anxiety disorder: Lorazepam
- Medicines for muscle weakness

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Buspan.

How to use Buspan Tablet

- How much to use

1 – 2 tablets

- When to use it

Take Buspan Tablet every 8 hourly at the same time every day. If you are not sure, check with your doctor or pharmacist.

- How long to use it

Always take Buspan Tablet exactly as your doctor has told you. If you are not

sure, check with your doctor or pharmacist.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

If you (or someone else) took a lot of tablets at the same time, or you think a child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately.

Signs of overdose symptoms are blurred vision, clumsiness, confusion, difficulty in breathing, dizziness, drowsiness, severe dryness of mouth, nose, or throat, very fast heartbeat, fever, hallucinations, muscle weakness, seizures, slurred speech, unusual excitement, restlessness, or irritability unusual warmth, dryness and flushing of skin.

While you are using it

- Things you must do

If the symptom still persists or not lighten after a few days, please consult your doctor.

- Things you must not do

- Do not give Buspan Tablet to anyone else, even if they have the same condition as you.
- Do not take Buspan Tablet to treat any other complaints unless your pharmacist or doctor tells you to.
- Do not take more than the recommended dose unless your pharmacist or doctor tells you to.

- Things to be careful of

- Buspan tablets contain lactose. If you have been told you have intolerance to some sugars, contact your doctor before taking this medicine.
- Buspan may cause drowsiness, if affected do not drive or operate machinery.

BUSPAN TABLET

Hyoscine butylbromide (10mg)

- Prolonged use of this medicine may decrease or inhibit salivary flow, thus contributing to the development of caries, gum disease, oral thrush and discomfort.

Side effects

Like all medicines, Buspan Tablet may cause side effects, although not everybody gets them.

Consult doctor or pharmacist for further information if you notice any of the following and they worry you:

- Dry mouth with difficulty in swallowing
- Increased pressure in the eye
- difficulty in urination
- constipation
- Allergic reaction such as skin rash or hives
- Confusion
- Dizziness, feeling faint
- Decreased flow of breast milk
- Decreased sweating
- Blurred vision
- Drowsiness
- Dryness of nose, throat or skin
- Hallucination
- Headache
- Loss of memory
- Dilatation of eye pupil
- Nausea or vomiting
- Trouble in sleeping
- Unusual tiredness or weakness

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Buspan Tablet

- Storage

Keep all medicines out of the reach of children.

Store below 30°C in a dry place protected from direct light.

- Disposal

Do not take this medicine after the expiry date stamped on the pack.

Return any unused tablets to your pharmacist to be disposed of. Only keep the tablets if your doctor tells you to. Do not throw them away in your normal household drainage or waste. This will help to protect the environment.

Product Description

- What it looks like

Round white tablet with score on one side and H 11 embossed on the other side with score.

- Ingredients

- Active ingredient(s)
Hyoscine Butylbromide.

- Inactive ingredients
Sodium benzoate, magnesium stearate, corn starch, explotab (sodium starch glycolate), lactose

- MAL number:

MAL19900043ACZ

Manufacturer and Product

Registration Holder

Terapeutics Sdn. Bhd. (590500-W)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat, S.P.S., Pulau Pinang,
Malaysia

Date of revision

30/11/2022

Serial Number

NPRA(R4/1)301122/00109