

PANAMOL SUSPENSION

Paracetamol (250mg/5ml)

What is in this leaflet

1. What Panamol Suspension is used for
2. How Panamol Suspension works
3. Before you use Panamol Suspension
4. How to use Panamol Suspension
5. While you are using it
6. Side Effects
7. Storage & Disposal of Panamol Suspension
8. Product Description
9. Manufacturer & Product Registration Holder
10. Date of Revision

Metoclopramide, Domperidone
- Antiseizure medication
- Blood thinning medication: Warfarin
- Cholesterol lowering medication: Cholestyramine
- Oral hormone contraceptive

How to use Panamol Suspension

- How much to use:

Children (6 - 12 years) : 5 ml - 10 ml
(1-2 tsp)
(1 - 5 years) : 2.5 ml - 5ml
(1/2-1 tsp)

- When to use it :

- To be given 3 or 4 times a day or as directed by your doctor or pharmacist.
- Do not give more than 4 doses in any 24 hour period.
- Leave at least 4 hours between doses

- How long to use it :

- Use Panamol Suspension for the duration as your doctor recommends.
- Do not give this medicine to your child for more than 3 days without speaking to your doctor or pharmacist.

- If you forget to use it :

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose) :

Contact your doctor or your pharmacist immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of paracetamol overdosage in the first 24 hours are pale, nausea, vomiting, eating disorder and abdominal pain. Liver damage may become

apparent 12 to 48 hours of ingestion. Abnormalities of sugar metabolism and metabolic acidosis (too much acid in the body) may occur.

In severe poisoning, it may progress to brain damage and death. Acute kidney failure with death of cells may develop. Irregular heartbeats, inflammation of pancreas and liver cell death have been reported.

While you are using it

- Things you must do :

- Use your medicine exactly as your doctor has told you.
- Tell the doctors, dentists and pharmacists treating you that you are taking Panamol Suspension.
- Tell your doctor immediately if you are pregnant while taking this medication.
- shake well before use of medication
- Always use the spoon supplied with the pack

- Things you must not do :

- Do not stop using the medicine unless advised by your doctor.
- Do not use any new medicines without consulting your doctor or pharmacist.
- Do not give Panamol Suspension to anyone else, even if they have the same symptoms or condition as you.
- This preparation contains PARACETAMOL. Do not use with any other paracetamol-containing products.
- Do not overfill the spoon.

- Things to be careful of :

This medicine may affect your ability to drive or use machines. If the medicine makes you feel sick, dizzy or tired, or gives you a headache, do not drive or use machines and contact your doctor immediately.

Side Effects

Like all medicines, Panamol Suspension can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

What Panamol Suspension is used for

Panamol Suspension is used for the relief of pain and fever.

How Panamol Suspension works

Paracetamol works to stop the pain messages from getting through to the brain. It also acts in the brain to reduce fever.

Before you use Panamol Suspension

- When you must not use it :

Do not take this medicine if:

- you are allergic to Paracetamol or to other ingredients in this product.

- Before you start to use it:

Please consult your doctor or pharmacist if you are/have:

- alcoholic liver disease.
- severe liver or kidney impairment.
- persistent symptoms
- pregnant, planning for pregnancy or breast-feeding. Paracetamol can be used during pregnancy if clinically needed at the lowest effective dose for the shortest possible time. Paracetamol is excreted in breast milk. Available data do not contraindicate breast feeding.

- Taking other medicines:

Tell the doctor if you are taking:

- Any other medicines including any medicine that you buy without prescription from a pharmacy, supermarket or health food shop.
- Medications that may interfere with Panamol Suspension such as:
 - Alcohol
 - Antidepressant medication
 - Antiemetic medication:

PANAMOL SUSPENSION

Paracetamol (250mg/5ml)

Adverse effects of paracetamol are rare, but Stevens Johnson Syndrome, Toxic Epidermal Necrolysis have been reported. There have been reports of blood disorder including low platelets and low white blood cell in the blood.

ALLERGY ALERT: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my
[Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Panamol Suspension

-Storage

Store below 30°C in a dry place, protected from direct sunlight.
Keep out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

Orange coloured viscous suspension with sweet taste and fruity odour.

- Ingredients

-Active Ingredient(s)

Paracetamol

-Inactive Ingredients

Tutti fruity, sorbitol, propylene glycol, sugar, purified water, sodium saccharin, sodium chloride, methyl paraben, propyl paraben, rhodigel, titanium dioxide, liquid glucose, silicone antifoam emulsion, sunset yellow, gum tragacanth.

-MAL Number:

MAL19940171XCZ

Manufacturer

Terapeutics Sdn. Bhd. (590500-W)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat S.P.S. Penang, Malaysia

Product Registration Holder

Zontron Pharmaceuticals Sdn. Bhd.
(445695-T)
Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat S.P.S. Penang, Malaysia

Date of Revision

14/10/2022

Serial Number

NPRA(R4/1)12102022/00086