

FLAUISEPT LOTION 0.1%

Acridlavine Hydrochloride (0.1% w/v)

What is in this leaflet

1. What Flavisept Lotion 0.1% is used for
2. How Flavisept Lotion 0.1% works
3. Before you use Flavisept Lotion 0.1%
4. How to use Flavisept Lotion 0.1%
5. While you are using Flavisept Lotion 0.1%
6. Side effects
7. Storage and Disposal of Flavisept Lotion 0.1%
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

What Flavisept Lotion 0.1% is used for

Flavisept Lotion is used for the treatment of infected wounds, burns, and skin disinfection.

How Flavisept Lotion 0.1% works

Flavisept Lotion 0.1% contains acridlavine which kills the bacteria causing infection in wounds or burns.

Before you use Flavisept Lotion 0.1%

- When you must not use it

Do not use if you are allergic to acridlavine, acridine derivatives, or any other ingredient in this product.

- Before you start to use it

- Caution when using near eyes as it may cause irritation
- Inform your doctor if you are pregnant or breastfeeding.
- Prolonged use may delay healing.

Taking other medicines

The medicine is for topical application, and will usually have minimal effect on other medications. However, please inform your doctor or pharmacist if you are taking other medicines.

How to use Flavisept Lotion 0.1%

- How much to use

Wash the affected area and dry carefully. Apply the lotion to the affected area or on suitable gauze dressing and renew daily if required.

- When to use it

Please follow the advice of your doctor or pharmacist. May be applied 2-3 times a day.

- How long to use it

You should check with your doctor or pharmacist.

- If you forget to use it

If you forget to use this medicine, use it as soon as you remember. Then, continue to use it at the usual time.

- If you use too much (overdose)

If you or anyone you know used too much Flavisept Lotion 0.1%, contact your doctor, an emergency department or a poisoning center immediately.

While you are using Flavisept Lotion 0.1%

- Things you must do

Inform your doctor or pharmacist of any other medications you are taking.

- Things you must not do

Do not stop using this medication although you feel better after a few days. Continue using for as long as your doctor or pharmacist has advised.

- Things to be careful of

If you are pregnant, planning to get pregnant or breastfeeding during treatment, you should not use Flavisept Lotion 0.1% unless advised by your doctor or pharmacist.

Side effects

Reported adverse effects caused by Flavisept Lotion 0.1% are skin problems such as contact dermatitis, rash and severe itching. If any of the side effects become serious or you experience any other side effects not listed in this leaflet, please inform your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Flavisept Lotion 0.1%

- Storage

Store in a cool, dry place below 30°C. protect from direct light. Keep out of reach of children.

- Disposal

Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like

Orange coloured clear lotion

- Ingredients

Active ingredient:

Acridlavine Hydrochloride

Inactive ingredients:

Chlorocresol, Purified water

MAL number:

MAL19940184XCZ

Manufacturer and Product Registration Holder:

TERAPUTICS SDN. BHD.

(590500-W)

Lot 10 & 11, PERDA Industrial Park, Lorong IKS Simpang Ampat B, 14100 Simpang Ampat, S.P.S., Pulau Pinang, Malaysia

Date of Revision

20 September 2019

Serial Number

NPRA(R4/1)190919/00260