

Ellqy Corns & Warts Treatment Solution

NAME AND STRENGTH OF ACTIVE SUBSTANCE : Salicylic Acid 17% w/v.

Other ingredients: Lactic Acid, Ethylcellulose, Acrylates/Octylacrylamide Copolymer, Isopropyl Alcohol.

PRODUCT DESCRIPTION : A clear and colourless liquid with characteristic odour.

INDICATIONS : For the topical treatment of corns, warts and calluses.

CONTRAINDICATIONS : Known hypersensitivity to Salicylic Acid or any of the ingredients.

ROUTE OF ADMINISTRATION : Topical.


RECOMMENDED DOSAGE : Apply a thin film of solution directly to the affected area using the plastic applicator from the bottle. Make sure the affected area is clean and dry before applying.

The successful use of Ellqy Corns & Warts Treatment Solution can be achieved by following these instructions :




STEP 1

Soak the corn/wart/callus in warm water for 5 minutes, then dry the area with your own towel.




STEP 3

Apply a thin film of the Ellqy Corns & Warts Treatment Solution directly to the affected area using the plastic applicator from the bottle. Avoid applying to the surrounding normal skin.



STEP 2

Using a pumice stone or manicure emery board, carefully rub along the surface of the corn/wart/callus.



STEP 4

Allow Ellqy Corns & Warts Treatment Solution to dry and cover with a plaster.

Continue treatment twice daily with Ellqy Corns & Warts Treatment Solution until the corn/wart/ callus has completely cleared and the ridge lines of the skin have been restored.

If any irritation develops then stop using the Ellqy Corns & Warts Treatment Solution and wait for the irritation to go away, then recommence as above. If it does not get better within 48 hours or if it is getting worse seek medical attention.

If Ellqy Corns & Warts Treatment Solution gets in the eyes or on mucus membranes then flush immediately with water for at least fifteen minutes and call your medical doctor.

Be careful to keep the lid tightly closed on the bottle otherwise it may evaporate.

DRUG INTERACTIONS : Unknown.

ADVERSE EFFECTS : Salicylic Acid preparation may cause dermatitis.

WARNINGS AND PRECAUTIONS :

- For external use only.
- Keep in a dry place below 30°C.
- Keep out of reach of children. Jauhi daripada kanak-kanak.
- Avoid contact with eyes.
- Do not use if you are a diabetic or have impaired circulation unless under a doctor's supervision or recommendation.
- Do not use on moles, birthmarks or unusual skin growths or near eyes or on mucous membranes.
- Do not treat corn/wart/callus over large areas at one time. Keep bottle tightly capped.

PERSISTENCE AND PATIENCE :

There is no immediate cure. The usual treatment time is 6 to 12 weeks. Persistence with the treatment and patience is essential!

PACKING : 10 ml glass bottle.

STORAGE : Store in a dry place below 30°C.

Recommended shelf-life : 3 years

MANUFACTURED BY :

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,

42000 Port Klang, Selangor, MALAYSIA.

Revision Date : November 2020