

OXY-NASE PAEDIATRIC NASAL DROPS

Oxymetazoline hydrochloride (0.025% w/v)

What is in this leaflet

1. What OXY-NASE PAEDIATRIC NASAL DROPS is used for
2. How OXY-NASE PAEDIATRIC NASAL DROPS works
3. Before you use OXY-NASE PAEDIATRIC NASAL DROPS
4. How to use OXY-NASE PAEDIATRIC NASAL DROPS
5. While you are using OXY-NASE PAEDIATRIC NASAL DROPS
6. Side effects
7. Storage and Disposal of OXY-NASE PAEDIATRIC NASAL DROPS
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

This leaflet answers some common questions about OXY-NASE PAEDIATRIC NASAL DROPS. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits they expect it will have for you.

Keep this leaflet with the medicine.

You may want to read it again.

What OXY-NASE PAEDIATRIC NASAL DROPS is used for

OXY-NASE PAEDIATRIC NASAL DROPS is indicated for the relief of nasal congestion.

How OXY-NASE PAEDIATRIC NASAL DROPS works

Oxymetazoline has a narrowing of the blood vessel effect. They act by

redistributing local blood flow to reduce swelling of the nasal lining thus improving ventilation, drainage and nasal stuffiness.

Before you use OXY-NASE PAEDIATRIC NASAL DROPS

-When you must not use it

Do not use OXY-NASE PAEDIATRIC NASAL DROPS if:

You are allergic (hypersensitive) to Oxymetazoline hydrochloride or any of the other ingredients in this product.

-Before you start to use it

Take special care with OXY-NASE PAEDIATRIC NASAL DROPS:

Tell your doctor if you have or have ever had:

- high blood pressure
- diabetes
- difficulty urinating due to an enlarged prostate gland
- enlarged thyroid
- heart disease.

Pregnancy and breast feeding:

- Safety in pregnancy and lactation has not been established. It is not to be used during pregnancy and lactation unless the potential benefits outweigh the risk.
- Ask your doctor or pharmacist for advice before taking any medicine.

-Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Combinations containing antidepressants such as tricyclic antidepressants or maprotiline or monoamine oxidase inhibitors, depending on the amount present,

may also interact with this medication.

How to use OXY-NASE PAEDIATRIC NASAL DROPS

-How much to use

Children above 1 year:

1 to 2 drops. Doses to be given twice to three times daily.

Apply up to four times daily until condition improves, then reduce frequency.

-When to use it

Use the medicine according to recommendation by doctor or pharmacist.

-How long to use it

Do not exceed the recommended dosage and do not use for longer than 3 days.

-If you forget to use it

Take the missing dose as soon as you remember.

If it is almost time for your next dose, skip the missed dose and take the medicine at your next regularly scheduled time.

Do not take a double dose to make up for the one you have missed.

-If you use too much (overdose)

If you accidentally take too much OXY-NASE PAEDIATRIC NASAL DROPS, tell your doctor immediately or go to your nearest hospital.

Oxymetazoline may produce profound central nervous system depression, leading to coma and marked reduction in body temperature, especially in infants. Consult a physician for more information on the management of overdose.

While you are using OXY-NASE PAEDIATRIC NASAL DROPS

-Things you must do

If you are about to start taking any other new medicine, tell your

pharmacist or doctor that you are taking OXY-NASE PAEDIATRIC NASAL DROPS.

-Things you must not do

Follow your doctor's instructions about any restrictions on food, beverages or activity while taking OXY-NASE PAEDIATRIC NASAL DROPS.

-Things to be careful of

During the first days of treatment with OXY-NASE PAEDIATRIC NASAL DROPS reactions may be impaired. Do not take charge of vehicles or machinery until the effect of OXY-NASE PAEDIATRIC NASAL DROPS treatment on the individual is known.

Use of the spray by more than one person may spread infection.

Side effects

OXY-NASE PAEDIATRIC NASAL DROPS may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away.

- stinging
- dryness of the mouth and throat
- sneezing
- worsen nasal congestion (prolonged use)

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of OXY-NASE PAEDIATRIC NASAL DROPS

-Storage

Keep container well closed. Protect from light. Store below 30 °C.

Keep out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

OXY-NASE PAEDIATRIC NASAL DROPS is a clear solution in 10ml and 100ml amber glass bottle respectively.

-Ingredients

-Active ingredients:

Oxymetazoline Hydrochloride
0.025% w/v.

-Inactive ingredients:

Benzalkonium Chloride, Citric Acid Anhydrous, Sodium Citrate, Disodium Edetate, Sorbitol Solution 70% and Purified Water.

-MAL number

MAL19870791AZ

Product Registration Holder

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang, Selangor,
Malaysia.

Manufacturer

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang, Selangor,
Malaysia.

Date of Revision

21/05/2026

Serial Number

NPRA(R2/1)260430/00056