beprogel®

BETAMETHASONE DIPROPIONATE TOPICAL SOLUTION

Avoid contact with eyes. Please refer to enclosed leaflet for precautions/ warnings / contraindications.

30 ml

HOE
Est. 1979

Wash and dry affected area. Apply sparingly to the affected areas once or twice daily or as directed by your doctor or pharmacist.
Route of administration: Topical

CONTROLLED MEDICINE

For external use only. Not for ophthalmic use. Flammable : Do not use or dry the hair near a fire or naked flame. Store below 30°C. Keep out of reach of children. Jauhi daripada kanak-kanak.

LOT:
MFG:
EXP:

beprogel®

BETAMETHASONE DIPROPIONATE TOPICAL SOLUTION

Avoid contact with eyes. Please refer to enclosed leaflet for precautions/ warnings / contraindications.

30 ml

BRU09020208P
MAL19913399AZ
SIN10242P
MM Reg: No. R2806A6085
Lao Reg. No.:10I2188/03

HOE
Est. 1979

UNTUK BUKA

Active Ingredient:
Betamethasone Dipropionate 0.064% w/v (Equivalent to Betamethasone 0.05% w/v)
Preservative:
Methyl Paraben 0.1% w/v

Product Registration Holder/Manufacturer:
HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang,
Selangor, MALAYSIA.

HOE
Est. 1979

beprogel

BETAMETHASONE TOPICAL SOLUTION

TO OPEN