

Note: Same design will be applied to all pack size.

Treatment of candidosis of the oropharyngeal cavity
and the gastrointestinal tract in adults and
pediatric patients 6 months and older.



Decozol oral gel 2% w/w

15g gel

HOE
Est. 1979

"Please refer to enclosed leaflet for Precautions/Warnings/Contraindications"

Decozol oral gel 2% w/w

15g gel

HOE
Est. 1979

Decozol oral gel 2% w/w

STORE BELOW 30°C.
KEEP OUT OF REACH OF CHILDREN.
JAUHI DARIPADA KANAK-KANAK.

MAL07010892AZ
BRU13021021P



HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,
42000 Port Klang, Selangor, MALAYSIA.

Est. 1979

Product Registration Holder/Manufacturer:

CONTROLLED MEDICINE

Route of Administration : Oral

Apply as directed by your doctor or pharmacist.

Please refer to enclosed leaflet.

RECOMMENDED DOSAGE:

Treatment of candidosis of the oropharyngeal cavity and the gastrointestinal tract in adults and pediatric

patients 6 months and older.

INDICATIONS:

Active Ingredient : Miconazole 2.0% w/w
Preservatives : Methyl Paraben 0.08% w/w,
Propyl Paraben 0.02% w/w

Decozol oral gel 2% w/w

Decozol
oral gel 2% w/w
HOE
Est. 1979

LOT:
MFG:
EXP: