



HOE ZINC CREAM

NAME AND STRENGTH OF ACTIVE SUBSTANCE

Zinc Oxide 32% w/w.

PRODUCT DESCRIPTION : A white cream.

Other ingredients: Palm Oil, Oleic Acid, Wool Fat, Sodium Chloride, Dimethicone Copolyol and Purified Water.

ROUTE OF ADMINISTRATION

Topical

PHARMACODYNAMICS

Zinc oxide is used externally in a cream base as a mild antiseptic and a mild astringent for the skin, as a soothing and protective application in eczema and as a protective for slight excoriations.

PHARMACOKINETICS

Zinc oxide is used externally and it is not absorbed in to the systemic blood. Acts as a mild astringent that precipitate proteins but have very little permeability that only the surface is affected.

INDICATIONS

A mild astringent for the skin, a soothing and protective application in eczema and a protective to slight excoriation.

CONTRAINDICATIONS

Known hypersensitivity to any component in the formulation.

PREGNANCY AND LACTATION

Unknown.

ADVERSE EFFECTS

Not known adverse effects studies on zinc oxide.

INTERACTIONS WITH OTHER MEDICAMENTS

Unknown.

WARNINGS AND PRECAUTIONS

For external use only. Discontinue use if irritation or sensitivity develops.

RECOMMENDED DOSAGE

Please stir before use. Apply two or three times daily. Wash and dry the affected area thoroughly before applying the cream.

SYMPTOMS AND TREATMENT FOR OVERDOSE

No information is available concerning overdosage in human beings following ingestion. In case of excessive ingestion, treat symptomatically.

PACKING

450g in polypropylene jar.

STORAGE

Keep container well closed. Protect from light. Store below 30°C. Keep out of the reach of children. For external use only.

Recommended shelf-life: 3 years.

MANUFACTURED BY

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,
42000 Port Klang, Selangor, MALAYSIA.

Date of revision : July 2023