

COUNTERPAIN ROLL ON TOPICAL LOTION

Glycol Salicylate 2.5% w/v, l-Menthol 3.0% w/v

What is in this leaflet

1. What COUNTERPAIN ROLL ON TOPICAL LOTION is used for
2. How COUNTERPAIN ROLL ON TOPICAL LOTION works
3. Before you use COUNTERPAIN ROLL ON TOPICAL LOTION
4. How to use COUNTERPAIN ROLL ON TOPICAL LOTION
5. While you are using it
6. Side effects
7. Storage and Disposal of COUNTERPAIN ROLL ON TOPICAL LOTION
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What COUNTERPAIN ROLL ON TOPICAL LOTION is used for

COUNTERPAIN ROLL ON TOPICAL LOTION is used to provide temporary relief of stiff neck and shoulder pain, backache, muscular pain, bruises, sprain and joint pain.

How COUNTERPAIN ROLL ON TOPICAL LOTION works

COUNTERPAIN ROLL ON TOPICAL LOTION contains the active ingredients Glycol Salicylate 2.5% w/v and l-Menthol 3.0% w/v.

Glycol Salicylate is a salicylic acid derivative used similarly to methyl salicylate in topical rubefacient preparations for the relief of muscular and rheumatic pain.

l-Menthol is an organic compound that occurs naturally in mint plants, but it can also be made synthetically. The primary action of menthol is stimulating sensory receptors and acts as counter-irritants. When applied to the skin, menthol dilates the blood vessels, causing a sensation of coldness followed by an analgesic effect.

Before you use COUNTERPAIN ROLL ON TOPICAL LOTION

- When you must not use it

Do not use COUNTERPAIN ROLL ON TOPICAL LOTION:

- In patients with aspirin or salicylate sensitivity.
- If you are hypersensitive to any of the ingredients in the product.
- In children below 2 years of age.

- Before you start to use it

Tell your pharmacist or doctor:

- If you are allergic to Glycol Salicylate or l-Menthol; or if you have any other allergies.

Pregnancy and breast feeding:

- Consult your doctor or pharmacist for advice before taking this medicine if you are pregnant or breast-feeding.

- Taking other medicines

Tell your pharmacist or doctor if you are taking any other products (such as over the counter drug, prescribed drug and herbal products).

How to use COUNTERPAIN ROLL ON TOPICAL LOTION

Apply a proper amount to affected area. Massage until thoroughly absorbed into skin.

- How much to use

Adult: Apply 2 or 3 times daily.

Consult with your doctor if you want to apply this medicine to children under 12 years of age.

- When to use it

When the symptom of aches and pains in muscles and joints appear.

- How long to use it

Consult your doctor if your pain persists for more than 7 days.

- If you forget to use it

If you miss a dose, use it as soon as you remember.

If you use too much (overdose)

No case of overdose has been reported. In the event of overdose with obvious clinical manifestations, stop use and consult a doctor or pharmacist.

While you are using it

- Things you must do

If you are about to start taking on any other new medicine, tell your pharmacist or doctor that you are using COUNTERPAIN ROLL ON TOPICAL LOTION.

- Things you must not do

Do not apply this medication to skin that is injured or irritated (e.g., cut, rashes and sunburned).

Do not apply this medication around the eyes, mucous membranes, or eczema.

Do not apply this product and other external remedies together on the same part.

Avoid bandage tightly or refrain warming the applied part with heating appliances such as electric blanket and other device.

- Things to be careful of

If your skin is sensitive, avoid repeated use at the same site.

Side effects

COUNTERPAIN ROLL ON TOPICAL LOTION may have unwanted side effects in some people.

Stop using this medication and tell your doctor immediately if any of these side effects occur:

- Rash
- Itching
- Redness
- Eruption

If you notice other effects not listed above, contact your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

- Ingredients

Active ingredients:

Glycol Salicylate 2.5% w/v, l-Menthol 3.0% w/v

Inactive ingredients:

Benzyl Nicotinate, Vanillyl Butyl Ether, Carbomer copolymer, Isopropyl Myristate, Alcohol, Triethanolamine 99%, Purified water

- MAL number

MAL25086029X

Manufacturer

HOE Pharmaceuticals Sdn. Bhd.

Under Authority of Taisho

Pharmaceutical Co., Ltd., Tokyo, Japan

Product Registration Holder

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang, Selangor,
Malaysia.

Date of revision

05/06/2026

Serial Number

NPRA (R2) 25/078

Storage and Disposal of COUNTERPAIN ROLL ON TOPICAL LOTION

- Storage

Store below 30°C.
Avoid direct sunlight.
Keep away from fire.
Keep out of reach of children.

- Disposal

Dispose the medicine after the expiry date. Consult your pharmacist on how to safely dispose of any medicines no longer required.

Product description

- What it looks like

COUNTERPAIN ROLL ON TOPICAL LOTION is colourless liquid and available in 100ml bottle.