

QURIC EFFERVESCENT GRANULES

Sodium Bicarbonate/ Sodium Citrate Anhydrous/ Anhydrous Citric Acid/ Tartaric Acid (1.76g/ 630mg/720mg/890mg)

What is in this leaflet

1. What QURIC is used for
2. How QURIC works
3. Before you use QURIC
4. How to use QURIC
5. While you are using QURIC
6. Side effects
7. Storage and Disposal of QURIC
8. Product description
9. Manufacturer
10. Product Registration Holder
11. Date of revision

What QURIC is used for

- Relieve of discomfort in mild UTI .
- Symptomatic relief of dysuria.
- Enhance the action of certain antibiotics especially some sulphonamides.
- In gout as urinary alkalinisers to prevent crystallisation of urates and dissolution of uric acid stones

How QURIC works

QURIC works as urinary alkaliser raising pH level of urine promoting dissolution of uric acid stones and providing relief of hyperacidity symptoms.

Before you use QURIC

- When you must not use it

If you are allergic to citric acid, sodium bicarbonate, sodium citrate, sodium carbonate or any other ingredients of QURIC . If you have severe symptoms or cloudy urine.

If you are pregnant or breastfeeding.

If you are suffer from heart disease, high blood pressure, diabetes or kidney disease.

If your diet only allows you to have a small amount of salt.

- Before you start to use it

Tell your healthcare professional if you are allergic to any other medicines, foods, dyes or preservatives.

Tell your healthcare professional if you are pregnant or plan to become pregnant.

Your healthcare professional will discuss the risks and benefits of taking QURIC during pregnancy.

Tell your healthcare professional if you are breastfeeding or wish to breastfeed.

Tell your healthcare professional if you have or have had any of the following medical conditions:

- Peptic ulceration
- Kidney problems
- Taking quinolone antibiotics

Tell your healthcare professional if you are on sodium restricted diet.

- Taking other medicines

QURIC should be avoided while taking antacid, ciprofloxacin, ofloxacin, ofloxacin and salicylates.

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use QURIC

- How much to use

You should check with your healthcare professional if you are not sure how to take QURIC .

The usual dose is 4-8g (1 to 2 sachets) dissolved in cold water four times daily

- When to use it

Sachet to be taken 4 times a day at regular interval throughout the day.

- How long to use it

Continue taking QURIC for as long as your healthcare professional tells you to. The length of treatment with QURIC will depend on how quickly your symptoms improve.

- If you forget to use it

Do not take a double dose to make up for a forgotten dose. If you have further questions on the use of this product, ask your healthcare professional.

- If you use too much (overdose)

Overdosage with sodium salts may cause serious electrolyte disturbances. Ingestion of large amounts of sodium citrate irritates the gastrointestinal mucosa & may result in diarrhea, nausea, vomiting, swelling and convulsions. If you accidentally take too much QURIC you should consult a healthcare professional or hospital as soon as possible.

QURIC EFFERVESCENT GRANULES

Sodium Bicarbonate/ Sodium Citrate Anhydrous/ Anhydrous Citric Acid/ Tartaric Acid (1.76g/ 630mg/720mg/890mg)

While you are using QURIC

- Things you must do

Drink plenty of water.

Urinate frequently. Don't delay going to the bathroom.

Wipe from front to back after a bowel movement.

- Things you must not do

Do not take QURIC to treat any other complaints unless your healthcare professional tells you to do so.

Do not give QURIC to anyone else, even if they have the same condition as you.

Do not stop taking QURIC or lower the dose, without checking with your healthcare professional.

- Things to be careful of

If symptoms persist after the 48 hour treatment is completed please consult your healthcare professional.

Side effects

Like all medicines, QURIC can cause side effects, although not everybody gets them. Taking QURIC may cause flatulence. If you notice this or any other effects, stop taking the product and tell your healthcare professional or pharmacist. Large doses of QURIC may cause diarrhea, nausea, vomiting and exert laxative effects. You may report any side effects or adverse drug reactions directly to the

National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers - > Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of QURIC

- Storage

Store below 30°C. Protect from moisture. Keep out of the reach of children. Do not use QURIC after the expiry date which is stated on the sachet. The expiry date refers to the last day of the month.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist /healthcare professional how to dispose of medicines no longer required.

These measures will help to protect the environment.

Product description

- What it looks like

White granular free flowing powder, which dissolve in 200 ml water gives clear colourless solution.

- Ingredients

- Active ingredient:

Sodium Bicarbonate, Sodium Citrate Anhydrous, Anhydrous Citric Acid, Tartaric Acid

- Inactive ingredient:

Flavor Lemon Dry
(Fresh Lemon Lime)

MAL number:

MAL25046045XC

Manufacturer

Akums Drugs & Pharmaceuticals Ltd.
19,20 and 21, Sector-6A, IIE,
SIDCUL, Ranipur, Haridwar- 249403,
Uttarakhand, India.

Product Registration Holder

Regulatory Dynamic Enterprise
12 Jalan Tekukur 2 , Bandar
Puchong Jaya
47170 Puchong, Selangor

Date of Revision

26/03/2026

Serial Number

NPRA (R2)24/175