

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only

(Metformin Hydrochloride Sustained Release and
Glimepiride Tablets)

GLIMET 1 mg/500 mg SUSTAINED RELEASE TABLETS
GLIMET 2 mg/1000 mg SUSTAINED RELEASE TABLETS

GLIMET 1 mg/500 mg SUSTAINED RELEASE TABLETS

Metformin hydrochloride USP [In sustained release form]	500 mg
Glimepiride USP	1 mg

Metformin hydrochloride USP [In sustained release form]	1000 mg
Glimepiride USP	2 mg

White to off white flat oblong shaped uncoated tablet having one blue round inlay tablet near the center of one side.

White to off white flat oblong shaped uncoated tablet having one pink round inlay tablet near the center of one side.

The primary mechanism of action of Glimepiride in lowering blood glucose appears to be dependent on stimulating the release of insulin from functioning pancreatic beta cells. In addition, extra pancreatic effects may also play a role in the activity of sulfonylurea such as Glimepiride.

Metformin is an antihyperglycemic agent, which improves glucose tolerance in patients with, type 2 diabetes, lowering both basal and post-prandial plasma glucose. Its pharmacologic mechanisms of action are different from other classes of oral antihyperglycemic agents. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing the peripheral glucose uptake and utilization. Unlike sulfonylurea, Metformin does not cause hypoglycemia in either with patients with type 2 diabetes or normal individuals.

Glimepiride is an oral glucose-lowering drug of the sulphonylureas class, which acts by increasing the release of insulin from the pancreatic beta cells. Metformin, a biguanides class, reduces the output of glucose from the liver and thus reduces the requirement of insulin at the level of liver. Thus, sulphonylureas and biguanides have additive action. Moreover, Metformin and Glimepiride are often co prescribed in clinical practice, and have proven to be efficacious in achieving the target glycaemic control and also in preventing the multiple metabolic defects, which are often present in diabetic subjects.

After oral administration, Glimepiride is completely (100%) absorbed from the GI tract. After oral administration of Glimepiride, maximum plasma concentration was observed after 2-3 hours.

The oral bioavailability of Metformin is about 50-60% and 30% of it is excreted via feces. Decrease in oral bioavailability could be due to first pass metabolism. Concomitant food intake may slightly impair the absorption.

The protein binding of Glimepiride was greater than 99.5%. The Volume of distribution was 113ml/kg and clearance was 47.8ml/min after an intravenous dose of Glimepiride in normal subjects.

Binding of Metformin to plasma proteins is negligible. Distribution is rapid. Metformin accumulates in kidneys, salivary glands and walls of oesophagus, stomach and duodenum. Binding to blood cells increases progressively. It is excreted into breast milk in small quantities.

Glimepiride:
Glimepiride is completely metabolized by oxidative biotransformation after either an IV or oral dose. The major metabolites are the cyclohexyl hydroxy methyl derivative (M1) and the carboxyl derivative (M2).

Cytochrome P450 II C9 has been shown to be involved in the biotransformation of Glimepiride to M1 (active). M1 is further metabolized to M2 (inactive) by one or several cytosolic enzymes. M1, but not M2, possesses about 1/3rd of the pharmacological activity as compared to its parent in an animal model.

Metformin is considered to be eliminated unchanged but some studies have indicated that some metabolic (about 20%) transformation may occur. No metabolites have been identified.

After single oral dose of Glimepiride $t_{1/2}$ was found to be 5.35 hours. No parent drug was recovered from urine or feces.

Metformin is excreted through the kidney by active tubular secretion though the renal clearance of the drug can be correlated with creatinine clearance. Elimination half-life is 1.5 to 4.5 hours.

GLIMET is indicated as an adjunct to diet and exercise in non-insulin dependent diabetes mellitus (NIDDM) (Type 2) patients who are unable to achieve sufficient glycaemic control with monotherapy of Metformin or Glimepiride alone or who are already treated with the combination of Metformin and Glimepiride as separate tablets.

The dosage of antidiabetic drugs should be individualized based on the patient's blood glucose levels. Generally, it should be recommended to initiate the lowest effective dose and increase the dose depending on the patient's blood glucose levels. Adequate monitoring of blood glucose levels should be performed.

GLIMET should be administered once per day during breakfast or the first main meal. The highest recommended dose per day should be 8mg of glimepiride and 2000mg of metformin.

When switching from combination therapy of Glimepiride plus Metformin as separate tablets, Glimet should be administered on the basis of dosage currently being taken.

A GFR should be assessed before initiation of treatment with metformin-containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months.

The maximum daily dose of metformin should preferably be divided into 2-3 daily doses. Factors that may increase the risk of lactic acidosis should be reviewed before considering initiation of metformin in patients with GFR <60 ml/min.

If no adequate strength of GLIMET 1 mg / 500 mg SUSTAINED RELEASE TABLETS and GLIMET 2 mg / 1000 mg SUSTAINED RELEASE TABLETS is available, individual monocomponents should be used instead of the fixed dose combination.

CONTRAINDICATIONS

1. Known hypersensitivity to the drug.
2. Diabetic ketoacidosis, with or without coma. This condition should be treated with insulin.

1. Renal disease or renal dysfunction (e.g., as suggested by serum creatinine levels 1.5 mg/dL [males], 1.4 mg/dL [females] or abnormal creatinine clearance), which may also result from conditions such as cardiovascular collapse (shock), acute myocardial infarction, and septicemia.
2. Congestive heart failure requiring pharmacological treatment.
3. Known hypersensitivity to Metformin hydrochloride.
4. Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma. Diabetic ketoacidosis should be treated with insulin.
5. Severely reduced kidney function (GFR <30 mL/min)
6. Any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis)

The administration of oral hypoglycemic drugs has been reported to be associated with increased cardiovascular mortality as compared to treatment with diet alone or diet plus insulin. This warning is based on the study conducted by the University Group Diabetes Program (UGDP), a long-term, prospective clinical trial designed to evaluate the effectiveness of glucose-lowering drugs in preventing or delaying vascular complications in patients with non-insulin-dependent diabetes. Results of the study suggested that there is increased incidence of cardiovascular mortality in patients taking sulfonylurea class of drugs.

All sulfonylureas are capable of producing severe hypoglycemia. Proper patient selection, dosage, and instructions are important to avoid hypoglycemic episodes. Hypoglycemia is more likely to occur when caloric intake is deficient, after severe or prolonged exercise, when alcohol is ingested, or when more than one glucose-lowering drug is used.

When a patient stabilized on any diabetic regimen is exposed to stress such as fever, trauma, infection, or surgery, a loss of control may occur. At such times, it may be necessary to add insulin in combination with Glimepiride or even use insulin monotherapy. The effectiveness of any oral hypoglycemic drug, including Glimepiride, in lowering blood glucose to a desired level decreases in many patients over a period of time, which may be due to progression of the severity of the diabetes or to diminished responsiveness to the drug. This phenomenon is known as secondary failure, to distinguish it from primary inactivity which the drug is ineffective in an individual patient when first given. Should a secondary failure occur with Glimepiride or metformin monotherapy, combined therapy with Glimepiride and metformin or Glimepiride and insulin may result in a response. Should secondary failure occur with combined Glimepiride/metformin therapy, it may be necessary to initiate insulin therapy.

Risk Related to Diabetes: Abnormal blood glucose levels during pregnancy are associated with a higher incidence of congenital abnormalities and perinatal

PRODUCT NAME	: Glimet	COUNTRY : Malaysia	LOCATION : Indrad	Supersedes A/W No.:			
ITEM / PACK	: Insert	NO. OF COLORS : 1	REMARK :				
DESIGN STYLE	:	PANTONE SHADE NOS.:	SUBSTRATE :				
CODE	: xxxxxxxx-5253		Activities	Department	Name	Signature	Date
DIMENSIONS (MM) (LxWxH)	: 180 x 300		Prepared By	Pkg.Dev			
ART WORK SIZE	: S/S	 Black	Reviewed By	Pkg.Dev			
DATE	: 18-11-2017		Approved By	Quality			