xxxxxxxxx-5253

TOLANZ ODT TABLETS 10 MG TOLANZ ODT TABLETS 5 MG

(Olanzapine Orodispersible Tablets 10 mg / 5 mg)

BRAND OR PRODUCT NAME TOLANZ ODT TABLETS 10 MG TOLANZ ODT TABLETS 5 MG

Each uncoated orodispersible tablet

NAME AND STRENGTH OF ACTIVE SUBSTANCE(S) TOLANZ ODT TABLETS 5 MG

TOLANZ ODT TABLETS 10 MG

Each uncoated orodispersible tablet contains

Olanzapine PRODUCT DESCRIPTION

TOLANZ ODT TABLETS 5 MG: Yellow colored, round, flat, bevel edged uncoated tablets with break line on one side and debossed "5" on other side.

TOLANZ ODT TABLETS 10 MG: Yellow colored, round, flat, bevel edged uncoated tablets with break line on one side and debossed "10" on other side

DOSAGE FORM

PHARMACODYNAMICS

Olanzapine is an antipsychotic agent that demonstrates a broad pharmacologic profile across a number of receptor systems. Olanzapine exhibited a range of receptor affinities (Ki: < 100 nM) for serotonin 5 HT2A/2C, 5 HT3, 5 HT6; dopamine D1, D2, D3, D4, D5; cholinergic muscarinic receptors. It is reported that olanzapine reduced a conditioned avoidance response, a test indicative of antipsychotic activity, at doses below those producing catalepsy, an effect indicative of motor side-effects. Unlike some other antipsychotic agents, olanza,

increases responding in an "anxiolytic" test.

Olanzapine produced a higher 5-HT₂A than dopamine D₂ receptor occupancy. In addition, olanzapine-responsive patients had lower striatal D2 occupancy than some other antipsychotic and risperidone-responsive patients, while being comparable to clozapine-responsive patients. It is reported that olanzapine was associated with statistically significantly greater improvements in negative as well as positive symptoms.

It is reported that olanzapine demonstrated statistically significant superiority over placebo on the primary endpoint of bipolar recurrence. Olanzapine also reported a statistically significant advantage over placebo in terms of preventing either recurrence into mania or

Olanzapine is well absorbed after oral administration reaching peak plasma concentrations within 5 to 8 hours. The absorption is not

Olanzapine is metabolized in the liver by conjugative and oxidative pathways. The major circulating metabolite is the 10-N-glucuronide which does not pass the blood brain barrier. Cytochromes P450- CYP1A2 and P450-CYP2D6 contribute to the formation of the Which does not pass the blood brain barrier. Sportholders 1430-647-1742 and 1430-747-20 controlled to the foliable of the Middle of the Middle

33.8 hr) and the clearance was reduced (17.5 versus 18.2 l/hr). The pharmacokinetic variability reported in the elderly is within the range for the non-elderly. It is reported that in 44 patients with schizophrenia >65 years of age, dosing from 5 to 20 mg/day was not associa

to the overall variability between individuals.

the average olanzapine exposure was approximately 27% higher in adolescents. Demographic differences between the adolescents and adults include a lower average body weight and fewer adolescents were smokers. Such factors likely contribute to the higher average exposure reported in adolescents.

The plasma protein binding of olanzapine was about 93 % over the concentration range of about 7 to about 1000 ng/mL. Olanzapine is

Olanzapine is indicated for the treatment of schizophrenia. Olanzapine is effective in maintaining the clinical improvement during continuing therapy in patients who have shown initial treatment response. Olanzapine is indicated for short-term treatment of acute manic episode associated with Bipolar I Disorder. Olanzapine is indicated for preventing recurrence of manic, mixed or depressive

Schizophrenia: The recommended starting dose for olanzapine is 10 mg/day.

Manic episode: The starting dose is 15 mg as a single daily dose in monotherapy or 10 mg daily in combination therapy.

Preventing recurrence in bipolar disorder: The recommended starting dose is 10 mg/day. For patients who have been receiving olanzapine for treatment of manic episode, continue therapy for preventing recurrence at the same dose. If a new manic, mixed, or depressive episode occurs, olanzapine treatment should be continued (with dose optimisation as needed), with supplementary therapy to treat mood symptoms, as clinically indicated.

During treatment for schizophrenia, manic episode, and recurrence prevention in bipolar disorder, daily dosage may subsequently be ted on the basis of individual clinical status within the range 5-20 mg/day. An increase to a dose greater than the recom starting dose is advised only after appropriate clinical reassessment and should generally occur at intervals of not less than 24 hours. Olanzapine can be given without regard for meals, as absorption is not affected by food. Gradual tapering of the dose should be

Olanzapine orodispersible tablet should be placed in the mouth, where it will rapidly disperse in saliva, so it can be easily swallowed. Removal of the intact orodispersible tablet from the mouth is difficult. Since the orodispersible tablet is fragile, it should be taken immediately on opening the blister. Alternatively, it may be dispersed in a full glass of water or other suitable beverage (orange juice,

Olanzapine orodispersible tablet has the same dosage and frequency of administration as olanzapine coated tablets. Olanzapine orodispersible tablets may be used as an alternative to olanzapine coated tablets.

Olanzapine is not recommended for use in children and adolescents below 18 years. A greater magnitude of weight gain, lipid and prolactin alterations has been reported adolescent patients than of adult patients

A lower starting dose (5 mg/day) is not routinely indicated but should be considered for those 65 and over when clinical factors warrant.

class A or B), the starting dose should be 5 mg and only increased with caution.

The starting dose and dose range need not be routinely altered for female patients relative to male patients

The starting dose and dose range need not be routinely altered for non-smokers relative to smokers.

When more than one factor is present which might result in slower metabolism (female gender, geriatric age, non-smoking status), consideration should be given to decreasing the starting dose. Dose escalation, when indicated, should be conservative in such patients. Mode of Administration Olanzapine orodispersible tablet should be placed in the mouth, where it will rapidly disperse in saliva, so it can be easily swallowed.

Removal of the intact orodispersible tablet from the mouth is difficult. Since the orodispersible tablet is fragile, it should be taken immediately on opening the blister. Alternatively, it may be dispersed in a full glass of water or other suitable beverage (orange juice, apple juice, milk or coffee) immediately before administration. CONTRAINDICATIONS

Hypersensitivity to olanzapine or any of the excipients. Patients with known risk for narrow-angle glaucoma. WARNINGS AND PRECAUTIONS

closely monitored during this period.

Dementia-related psychosis and/or behavioural disturbances

Olanzapine is not approved for the treatment of dementia-related psychosis and/or behavioural disturbances and is not recommended for use in this particular group of patients because of an increase in mortality and the risk of cerebrovascular accident. There was a 2-fold increase in the incidence of death reported in olanzapine-treated patients. The higher incidence of death was not associated with olanzapine dose (mean daily dose 4.4 mg) or duration of treatment. Risk factors that may predispose this patient population to increased mortality include age > 65 years, dysphagia, sedation, malnutrition and dehydration, pulmonary conditions (e.g., pneumonia, with or without aspiration), or concomitant use of benzodiazepines. However, the incidence of death was higher in olanzapine-treated

olanzapine. All olanzapine patients who experienced a cerebrovascular event had pre-existing risk factors. Age> 75 years and vascular/mixed type dementia were reported as risk factors for CVAE in association with olanzapine treatment.

recommended. Worsening of Parkinsonian symptomatology and hallucinations were reported with use of olanzapine

signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. If a patient develops signs and symptoms indicative of NMS, or presents with unexplained high fever without additional clinical manifestations of NMS, all

patients treated with atypical antipsychotics. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atvoical an increased risk of treatment-emergent hyperglycaemia-related adverse events in patients treated with the atypical antipsychotics.

Precise risk estimates for hyperglycaemia-related adverse events in patients treated with atypical antipsychotics are not available. Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycaemia including polydipsia, polyuria; polyphagia, and weakness. Patients who develop symptoms of hyperglycaemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycaemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of antidiabetic treatment despite discontinuation of the suspect drug.

appropriate, particularly in dyslipidemic patients and in patients with risk factors for the development of lipids disorders. Patients treated with any antipsychotic agents, including olanzapine, should be monitored regularly for lipids in accordance with utilised antipsychotic guidelines, e.g., at baseline, 12 weeks after starting olanzapine treatment and every 5 years thereafter. Anticholinergic activity

Olanzapine had anticholinergic activity. As clinical experience with olanzapine in patients with concomitant illness is limited, caution is advised when prescribing for patients with prostatic hypertrophy, or paralytic ileus and related conditions.

Caution should be exercised and follow-up organised in patients with elevated ALT and/or AST, in patients with signs and symptoms of hepatic impairment, in patients with pre-existing conditions associated with limited hepatic functional reserve, and in patients who are being treated with potentially hepatotoxic medicines. In cases where hepatitis (including hepatocellular, cholestatic or mixed liver injury) has been diagnosed, olanzapine treatment should be discontinued.

Caution should be exercised in patients with low leucocyte and/or neutrophil counts for any reason, in patients receiving medicines known to cause neutropenia, in patients with a history of drug- induced bone marrow depression/toxicity, in patients with bone marrow depression caused by concomitant illness, radiation therapy or chemotherapy and in patients with hypereosinophilic conditions or with myeloproliferative disease. Neutropenia has been reported commonly when olanzapine and valproate are used concomitantly.

Discontinuation of treatment Acute symptoms such as sweating, insomnia, tremor, anxiety, nausea, or vomiting have been reported when olanzapine is stopped

As with other antipsychotics, caution should be exercised when olanzapine is prescribed with medicines known to increase QTc interval, especially in the elderly, in patients with congenital long QT syndrome, congestive heart failure, heart hypertrophy, hypokalaemia or hypomagnesae

<u>Thromboembolism</u> Temporal association of olanzapine treatment and venous thromboembolism has been reported with use of olanzapine. A causal relationship between the occurrence of venous thromboembolism and treatment with olanzapine has not been reported. However, since patients with schizophrenia often present with acquired risk factors for venous thromboembolism, all possible risk factors of VTE e.g.,

Given the primary CNS effects of olanzapine, caution should be used when it is taken in combination with other centrally acting

medicines and alcohol. Olanzapine may antagonise the effects of direct and indirect dopamine agonists.

Olanzapine should be used cautiously in patients who have a history of seizures or are subject to factors which may lower the seizure threshold. Seizures have been reported to occur rarely in patients when treated with olanzapine. In most of these cases, a history of

Olanzapine was reported a lower incidence of treatment-emergent dyskinesia. However, the risk of tardive dyskinesia increases with long-term exposure, and therefore if signs or symptoms of tardive dyskinesia appear in a patient on olanzapine, a dose reduction or discontinuation should be considered. These symptoms can temporally deteriorate or even arise after discontinuation of treatment.

patients over 65 years.

The event of sudden cardiac death has been reported in patients with plantagine. The risk of presumed sudden cardiac death in patients treated with olanzapine was reported approximately twice the risk in patients not using antipsychotics. The risk of olanzapine was reported comparable to the risk of atypical antipsychotics.

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with clanzapine exposure. DRESS consists of a combination of three or more of the following: cutaneous reaction (such as rash or exfoliative dermatitis), eosinophilia, fever, lymphadenopathy and one or more systemic complications such as hepatitis, nephritis, pneumonitis, myocarditis, and pericarditis. Discontinue olanzapine if DRESS is suspected.

Paediatric population
Various adverse reactions, including weight gain, changes in metabolic parameters and increases in prolactin levels were reported in patients aged 13-17. Long-term outcomes associated with these events have not been reported and remain unknown

This formulation contains aspartame, which is a source of phenylalanine. May be harmful for people with phenylketonuria.

<u>Mannitol</u>

mean increase in olanzapine Cmax following fluvoxamine was 54 % in female nonsmokers and 77 % male smokers. The mean increase in olanzapine AUC was 52 % and 108 % respectively. A lower starting dose of olanzapine should be considered in patients who are using

hours before or after olanzapine, Fluoxetine (a CYP2D6 inhibitor), single doses of antacid (aluminum, magnesium) or cimetidine have not been found to significantly affect the pharmacokinetics of olanzapine.

expected as verified through in vivo studies where no inhibition of metabolism of the following active substances was found: tricyclic antidepressant (representing mostly CYP2D6 pathway), warfarin (CYP2C9), theophyline (CYP1A2) or diazepam (CYP3A4 and

concomitant olanzapine.

Effects on Ability to Drive and Use Machines.

No studies on the effects on the ability to drive and use machines have been performed. Because olanzapine may cause somnolence

self-limited, in other cases neonates have required intensive care unit support and prolonged hospitalisation.

TOLANZ ODT TABLETS should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus

Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during treatment with

Blood and the lymphatic system disorders

Immune system disorders

Metabolism and nutrition disorders

Weight gain, Elevated cholesterol levels, Elevated glucose levels, Elevated triglyceride levels, Glucosuria, Increased appetite, Development or exacerbation of diabetes occasionally associated with ketoacidosis or coma, including some fatal cases,

symptoms, Restless legs syndrome Cardiac disorders

Vascular disorders Orthostatic hypotension, Thromboembolism (including pulmonary embolism and deep vein thrombosis)

Hepato-biliary disorders

Transient, asymptomatic elevations of hepatic aminotransferases (ALT, AST), especially in early treatment, Hepatitis (including hepatocellular, cholestatic or mixed liver injury)

Rash. Photosensitivity reaction, Alopecia

Musculoskeletal and connective tissue disorders

Renal and urinary disorders

Urinary incontinence, urinary retention, Urinary hesitation

roductive system and breast disorders

Erectile dysfunction in males, Decreased libido in males and females, Amenorrhea, Breast enlargement, Galactorrhea in females, Gynaecomastia/breast enlargement in males, Priapism

Elevated plasma prolactin levels, High creatine phosphokinase, Increased total bilirubin, Increased alkaline phosphatase

Respiratory, Thoracic and Mediastinal Disorders:

*Atypical antipsychotic drugs, such as Olanzapine, have been associated with cases of sleep apnoea, with or without concomitan

The proportion of patients who had adverse, clinically significant changes in weight gain, glucose, total/LDL/HDL cholesterol or triglycerides reported increased over time. In adult patients who completed 9-12 months of therapy, the rate of increase in mean blood glucose slowed after approximately 6 months.

Additional information on special populations n elderly patients with dementia, olanzapine treatment was associated with a higher incidence of death and cerebrovascular adverse eactions. Very common adverse reactions reported with the use of olanzapine in this patient group were abnormal gait and falls

patients with bipolar mania, valproate combination therapy with olanzapine reported an incidence of neutropenia; a potential contributing factor could be high plasma valproate levels. Olanzapine administered with lithium or valproate reported in increased levels

of tremor, dry mouth, increased appetite, and weight gain. Speech disorder was also reported commonly. During treatment with olanzapine in combination with lithium or divalproex, an increase body weight occurred in patients during acute treatment (up to 6 weeks). Long-term planzapine treatment (up to 12 months) for recurrence prevention in patients with bipolar disorder was reported with an increase body weight.

Hepato-biliary disorders

Elevations of hepatic aminotransferases (ALT/AST)

Decreased total bilirubin, increased GGT, elevated plasma prolactin levels. OVERDOSE AND TREATMENT

Management of Overdose There is no specific antidote for olanzapine. Induction of emesis is not recommended. Standard procedures for management of overdose may be indicated (i.e. gastric lavage, administration of activated charcoal). The concomitant administration of activated charcoal was shown to reduce the oral bioavailability of olanzapine by 50 to 60%. Symptomatic treatment and monitoring of vital organ function should be instituted according to clinical presentation, including treatment of hypotension and circulatory collapse and support of respiratory function. Do not use epinephrine, dopamine, or other sympathomimetic agents with beta-agonist activity since beta stimulation may

Store below 30°C, Protected from light and moisture PACKAGING AVAILABLE

Indrad-382 721, Dist. Mehsana, INDIA

TOLANZ Tablets are packed in Alu-Alu blister of 10 tablets. Such blisters containing 10 Tablets are packed into boxes of 10's, 30's, 60's

100's. Not all presentations may be available locally. DATE OF REVISION OF PACKAGE INSERT

NAME AND ADDRESS OF MANUFACTURER

PHARMA

matching. colour actual for reference Department
Pkg.Dev
Pkg.Dev
Quality **Pantone** Follow colour binding. COUNTRY: N
NO. OF COLC
PANTONE SH
Black not <u>s</u>

Front/Back xxxxxxxx-52 150 x 460 m S/S 09-08-2019 proof colour

This

PHARMACOKINETICS affected by food.

elimination half-life of olanzapine in healthy subjects varied on the basis of age and gender.

It is reported that in healthy elderly (65 and over) versus non-elderly subjects, the mean elimination half-life was prolonged (51.8 versus

with any distinguishing profile of adverse events. The plasma clearance of olanzapine is lower in elderly versus young subjects, in females versus males, and in non-smokers versus smokers. However, the magnitude of the impact of age, gender, or smoking on olanzapine clearance and half-life is small in comparison

Adolescents (ages 13 to 17 years) - The pharmacokinetics of olanzapine are similar between adolescents and adults. It is reported that

RECOMMENDED DOSAGE AND MODE OF ADMINISTRATION

considered when discontinuing olanzapine.

apple juice, milk, or coffee) immediately before administration

Elderly patients Patients with renal and/or hepatic impairment

A lower starting dose (5 mg) should be considered for such patients. In cases of moderate hepatic insufficiency (cirrhosis, Child-Pugh

During antipsychotic treatment, improvement in the patient's clinical condition may take several days to some weeks. Patients should be

independent of these risk factors. Cerebrovascular adverse events (CVAE e.g., stroke, transient ischaemic attack), including fatalities, were also reported with use of

The use of olanzapine in the treatment of dopamine agonist associated psychosis in patients with Parkinson's disease is not Neuroleptic Malignant Syndrome (NMS) NMS is a potentially life-threatening condition associated with antipsychotic medicinal products. Rare cases reported as NMS have also been reported in association with olanzapine. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional

antipsychotic medicines, including olanzapine must be discontinued. Hyperglycemia and diabetes mellitus Hyperglycaemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in antipsychotic use and hyperglycaemia-related adverse events is not completely understood. However, epidemiological studies suggest

Lipid alterations Undesirable alterations in lipids have been reported in olanzapine-treated patients. Lipid alterations should be managed as clinically

Hepatic function Transient, asymptomatic elevations of hepatic aminotransferases, ALT, AST have been reported commonly, especially in early treatment,

abruptly.

tion of patients, should be identified and preventive measures undertaken General CNS activity

Tardive dyskinesia

Postural hypotension was reported in the elderly in olanzapine. It is recommended that blood pressure is measured periodically in

Sudden cardiac death

INTERACTIONS WITH OTHER MEDICAMENTS Caution should be exercised in patients who receive medicinal products that can cause central nervous system depression.

Potential interactions affecting olanzapine: Since olanzapine is metabolized by CYP1A2, substances that can specifically induce or

inhibit this isoenzyme may affect the pharmacokinetics of olanzapine.

Induction of CYP1A2: The metabolism of olanzapine may be induced by smoking and carbamazepine, which may lead to reduced olanzapine concentrations. Only slight to moderate increase in olanzapine clearance has been observed. The clinical consequences are likely to be limited, but clinical monitoring is recommended and an increase of olanzapine dose may be considered if necessary. *Inhibition of CYP1A2*: Fluvoxamine a specific CYP1A2 inhibitor has been shown to significantly inhibit the metabolism of olanzapine. The

fluvoxamine or any other CYP1A2 inhibitors, such as ciprofloxacin or ketoconazole. A decrease in the dose of olanzapine should be considered if treatment with an inhibitor of CYP1A2 is initiated.

Decreased bioavailability: Activated charcoal reduces the bioavailability of oral olanzapine by 50 to 60% and should be taken at least 2

Potential for olanzapine to affect other medicinal products: Olanzapine may antagonize the effects of direct and indirect dopamine Olanzapine does not inhibit the main CYP450 isoenzymes in vitro (e.g. 1A2, 2D6, 2C9, 2C19.3A4). Thus no particular interaction is

2C19). Olanzapine showed no interaction when co-administered with lithium or biperiden. Therapeutic monitoring of valproate plasma levels did not indicate that valproate dosage adjustment is required after the introduction of

and dizziness, patients should be cautioned about operating machinery, including motor vehicles. STATEMENT ON USAGE DURING PREGNANCY AND LACTATION Neonates exposed to antipsychotic drugs during the third trimester of pregnancy are at risk for extrapyramidal and/or withdrawal symptoms following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been

TOLANZ ODT TABLETS was excreted in breast milk. Patients should be advised not to breast feed an infant if they are taking

ADVERSE EFFECTS / UNDESIRABLE EFFECTS

Hypothermia Nervous system disorders Somnolence, Dizziness, Akathisia, Parkinsonism, Dyskinesia, Seizures where in most cases a history of seizures or risk factors for seizures were reported, Neuroleptic malignant syndrome, Dystonia (including oculogyration), Tardive dyskinesia, Discontinuation

Bradycardia, QTc prolongation, entricular tachycardia/fibrillation, sudden death

Gastrointestinal disorders Mild, transient anticholinergic effects including constipation and dry mouth, Pancreatitis

Skin and subcutaneous tissue disorder Very rare: Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS).

Pregnancy, puerperium and perinatal conditions
Drug withdrawal syndrome neonatal

General disorders and administration site conditions Asthenia, Fatigue, Oedema

weight gain. In patients who have a history of or are at risk for sleep apnoea, Tolanz ODT Tablets should be prescribed with caution. Long-term exposure (at least 48 weeks)

Pneumonia, increased body temperature, lethargy, erythema, visual hallucinations and urinary incontinence were reported commonly. Patients with drug-induced (dopamine agonist) psychosis associated with Parkinson's disease, worsening of Parkinsoniar symptomatology and hallucinations were reported very commonly.

Paediatric population Olanzapine is not indicated for the treatment of children and adolescent patients below 18 years Metabolism and nutrition disorders Weight gain, elevated triglyceride levels¹⁰, increased appetite, Elevated cholesterol levels Nervous system disorders Sedation (including: hypersomnia, lethargy, somnolence). Gastrointestinal disorders

Very common symptoms in overdose include tachycardia, agitation, aggressiveness, dysarthria, various extrapyramidal symptoms, and reduced level of consciousness ranging from sedation to coma. Other medically significant sequelae of overdose include delirium, convulsion, coma, possible neuroleptic malignant syndrome, respiratory depression, aspiration, hypertension or hypotension, cardiac arrhythmias and cardiopulmonary arrest. Fatal outcomes have been reported for acute overdoses as low as 450mg but survival has also been reported following acute overdose of 1,500mg.

worsen hypotension. Cardiovascular monitoring is necessary to detect possible arrhythmias. Close medical supervision and monitoring should continue until the patient recovers. STORAGE CONDITIONS

