

LION BALL CORNCAP PLASTER 50%w/w

LION BALL CORNCAP PLASTER 50% w/w is band type corn plaster that accurately applies to affected area to effectively remove corn and wart.

Drug Facts: Among 1 Sheet (9.46cm², 64mg) salicylate acid (KP) 32.00 mg


Appearance: Surface side of the product is similar to that of general band, plastic adhesive tape of sulfur white. On the internal side, salicylic acid adhesive tape is attached roundly on circular shape of felt. Adhesive side is milky white, and it sticks well on the skin.

Indication: Corn, callus, wart removal.

Dosage: Adults and children 16 years old and above, put central part having active ingredient on affected area and change to a new one everyday for up to 2 days. Repeat treatment if necessary and treatment should not continue for more than 6 weeks.

For areas such as sole where adhering is hard, adhere with disposable band or bandage.

When exchanging, tear off whitened dead skin with disinfected tweezers until not feel the pain.



Caution for use:

1. Do not administer to following patients and part
 - 1) Patient having a history of hypersensitivity.
 - 2) Diabetes patient.
 - 3) Peripheral blood patient.
 - 4) Around eye and eye membrane.
 - 5) On the part of infection, inflammation, and rash.
 - 6) Mole, mole since birth, wart with hair and wart around genital area.
 - 7) Infants and children below 16 years old.
2. Carefully administer to following patient
 - 1) Allergic to drug, cosmetics, or others (ex: rash, redness, itching, dermatitis caused by lacquer)
 - 2) Renal insufficiency patient (When large volume of salicylic acid is absorbed to skin, it might cause poisoning, so it shall not be extensively applied to renal insufficiency patient.)
3. Side effects
 - 1) Rash, redness, erythema, and itching may appear, and if these symptoms arise, stop the treatment.
 - 2) Continuous use of this drug may cause dermatitis, skin ulcers. If these symptoms arise, stop the treatment.
 - 3) When this drug is in contact with normal skin, local irritating may appear. When this happens, stop the treatment.
 - 4) Rarely irritation, contact allergic reactions may arise.
4. General caution
 - 1) Follow regulated dosage.
 - 2) When there is no improvement after using the drug many times, stop the treatment and consult pharmacist or doctor.
 - 3) Pyodermitis, wetting, inflammation shall be treated in advance of applying this drug.
 - 4) When going into water with this drug on may cause the activate site to be dissolved.
 - 5) When applied extensively side effects may arise. Use with caution.
 - 6) Patient who had used this drug by a mistake shall consult with doctor or a pharmacist.

5. Interaction

- 1) When used with MTX (methothrexate), it may increase toxicity of MTX.
- 2) This drug might stimulate absorption of other topical application drug.

6. Administration on pregnant women and lactating women

- 1) Pregnancy: There is no data on the use of topical salicylic acid in pregnant women. Therefore the use of this product during pregnancy is contraindicated.
- 2) Breast-feeding: Salicylates should be given with caution to breast-feeding mothers because of the possible risk of Reye's syndrome in nursing infants and there is no data on the use of topical salicylic acid in breast feeding women. Therefore the use of this product during pregnancy is contraindicated.

7. Precautions for application

- 1) This drug is only for external use.
- 2) Do not apply to near eye, nose, mouth, and other mucosa. When got into eye by mistake, wash completely with enough volume of water and follow direction from doctor.
- 3) Be careful not to apply to skin other than affected area.
- 4) Do not reuse.

8. Symptoms and treatment of overdose

Skin irritation and redness around the application site may become intolerable. Stop the medication and seek advice from your doctor/pharmacist immediately.

Storage conditions

Store below 30°C. Avoid storing in direct sunlight.

Keep out of reach from children. Jauhi daripada kanak-kanak.

Packaging: 6 and 8 sheets per pack

Manufacturer:

SINSIN PHARMACEUTICAL CO.,LTD.

9, Sojeongsandandong-ro, Sojeong-myeon, Sejong-si, South Korea

Date of revision: 19/3/2025