

SANGOBION COMPLETE CAPSULE

Ferrous gluconate / manganese sulfate / copper sulfate / vitamin C / folic acid / vitamin B12 / sorbitol (250mg / 0.2mg / 0.2 mg / 50mg / 1mg / 7.5µg / 25mg)

What is in this leaflet

1. What Sangobion Complete Capsule is used for
2. How Sangobion Complete Capsule works
3. Before you use Sangobion Complete Capsule
4. How to use Sangobion Complete Capsule
5. While you are using it
6. Side effects
7. Storage and Disposal of Sangobion Complete Capsule
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Sangobion Complete Capsule is used for

Sangobion Complete Capsule is used to treat anemia which is a condition in which the blood doesn't have enough healthy red blood cells. It is also used to prevent anemia in blood donors.

How Sangobion Complete Capsule works

Sangobion Complete Capsule helps with absorption of iron and involves in the formation of red blood cells.

Before you use Sangobion Complete Capsule

- When you must not use it
Do not use Sangobion Complete Capsule if you are allergic (hypersensitive) to any of the ingredients in this medicine.
- Before you start to use it
Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.
- Taking other medicines
Inform your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicine obtained without prescription such as supplements or traditional medicines.

How to use Sangobion Complete Capsule

- How much to use
1 to 2 capsules daily.

Sangobion Complete Capsule should be taken with plenty of water.

- When to use it
Take Sangobion Complete Capsule during or after meals or as directed by your doctor or pharmacist.
- How long to use it
Continue using Sangobion Complete Capsule for as long as your doctor recommends.

- If you forget to use it
Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)
Contact your doctor immediately or go to the Emergency Department of your nearest hospital if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. Urgent medical attention may be required.

While you are using it

- Things you must do
Tell your doctor and pharmacist treating you that you are using Sangobion Complete Capsule. Tell your doctor immediately if you become pregnant while using this medication.
- Things you must not do
Do not use the product after its expiry date.

Do not take any new medicines without consulting your doctor or pharmacist first.

- Things to be careful of
Driving and using machines

This medicine may affect your ability to drive or use machines. If the capsules make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, Sangobion Complete Capsule can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Sangobion Complete Capsule

- Storage
Keep out of the reach and sight of children.

Store below 25°C and protect from moisture.

- Disposal
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like
Dark brown hard gelatin capsules, printed with 'Merck' logo on the cap and body containing greenish/brown grey powder.

- Ingredients

- Active ingredient(s)

Ferrous gluconate
Manganese sulfate
Copper sulfate
Vitamin C
Folic acid
Vitamin B12
Sorbitol

- Inactive ingredients

Talc
Aerosil 200
Ethyl vanillin
Lactose

- MAL number:

MAL19911002X

Manufacturer

PT. Merck Tbk.,
Jl T.B. Simatupang No. 8,
Pasar Rebo,
Jakarta 13760, Indonesia.

Product Registration Holder

Procter & Gamble (Malaysia) Sdn. Bhd.
(8842-P),
10th Floor Surian Tower,
1 Jalan PJU 7/3,
Mutiar Damansara, 47810,
Petaling Jaya, Selangor D.E.
Malaysia.

Date of revision

15/09/2022

Serial Number

NPRA(R4/1)010922/00062