



## PRIDRINATE SYRUP 25mg/5ml

Description: Plain, blackcurrant flavoured syrup.

### Contents:

Pridrinate Syrup: Dimenhydrinate 25mg/5ml  
Methylparaben 0.1%w/v  
Propylparaben 0.01%w/v

### Indications

For prevention and treatment of motion sickness and symptomatic treatment of vertigo due to Meniere's disease and other labyrinthine disturbances.

### Pharmacology

Its exact pharmacological mechanism is not precisely known but may be related to its action on the medullary chemoreceptor trigger zone. It diminishes vestibular stimulation and depresses labyrinthine function.

### Adverse Effects

Drowsiness, gastrointestinal disturbances, thickening of bronchial secretion, dryness of the mouth, nose, and throat and skin rash.

### Precautions

- Caution in patients with acute asthma, bladder neck obstruction, cardiovascular disease, narrow-angled glaucoma, hypertension, hyperthyroidism, peptic ulcer, prostatic hypertrophy, pyloroduodenal obstruction, renal function impairment and in geriatric patients.
- Caution when driving or doing jobs requiring alertness because of possible drowsiness. Avoid alcoholic beverages.
- Safety for use in pregnancy and lactation has not been established.

### Contraindications

- Avoid in patients known to be hypersensitive to antihistamines.
- Use is not recommended in nursing mothers and in newborn or premature infants.

### Durg interactions

- Co-administration of monoamine oxidase (MAO) inhibitors may prolong and intensify the antimuscarinic effects of antihistamines.
- Concurrent use with alcohol, other CNS depressants, tricyclic antidepressants or atropine may potentiate the effects of either these medications or antihistamines.

### Dosage and Administration

Adults : Oral, 50mg every 4 hours as needed; not exceeding 400mg daily.  
Children 2 to 6 years : Oral, 12.5 to 25mg every 6 to 8 hours as needed; not exceeding 75mg daily.  
Children 7 to 12 years : Oral, 25 to 50mg every 6 to 8 hours as needed; not exceeding 300mg daily.

Note: For prophylaxis of motion sickness, it should be taken at least 30 minutes before departure.

### Overdosage

Clinical signs include unsteadiness, flushing, shortness of breath, hallucinations, trouble sleeping, severe drowsiness, hypotension, and convulsions.

Treatment consists of gastric lavage and convulsions may be controlled with diazepam given intravenously.

Emetics should not be used.

**Storage conditions** : Keep container tightly closed.  
Store in a cool, dry place below 30°C. Protect from light. Keep out of reach of children.

**Presentation/ Packing** : Syrup 25mg/5ml x 60ml

**Manufactured &** PRIME PHARMACEUTICAL SDN. BHD.

**Distributed by** : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,  
14000 Bukit Mertajam, Penang, Malaysia.