

KEEP OUT OF REACH OF CHILDREN
JAUHKAN DARI KANAK-KANAK


Store in a cool, dry place, below 30°C.
Protect from light.
Do not use if you notice any change in colour or odour.
Keep away from children.
Simpan dalam tempat yang sejuk dan kering.
Elakkan cahaya matahari langsung dan cahaya.
Jauhkan dari kanak-kanak.

SPASMIN TABLET


Each tablet contains:
DICYCLOMINE HCL20mg

Indications:
For treatment of irritable bowel syndrome, spastic constipation, spastic colitis, pylorospasm, and biliary dyskinesia

Dosage: Adult : 10 to 20 mg 3 or 4 times a day.
(ORAL) Children over 2 years : 10 mg 3 or 4 times a day.

1,000 tablets  **Controlled Medicine**

WARNING
Dicyclomine is not recommended for use in infants under the age of six months.

 **Manufacturer & Distributed by:**
PRIME PHARMACEUTICAL SDN. BHD.
150, Lorong Perumahan Usma 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

Batch No. / No. Batch :
Mfg. Date/ Tarikh Dibuat :
Exp. Date/ Tarikh Luput :
Reg. No. / No. Pendaftaran: MAL-19830132AZ