# **RASITIN TABLET**

Cloperastine hydrochloride (10mg)

#### What is in this leaflet

- 1. What *Rasitin* is used for
- 2. How Rasitin works
- 3. Before you use *Rasitin*
- 4. How to use Rasitin
- 5. While you are using it
- 6. Side effects
- 7. Storage and Disposal of Rasitin
- 8. Product description
- 9. Manufacturer and Product Registration holder

10.Date of revision

### What Rasitin is used for

For the relief of dry cough.

# How Rasitin works

Cloperastine is a medicine with effect of reducing cough similar to that of codeine. It While you are using Rasitin also has some antihistaminic effect.

# Before vou use Rasitin

- When you must not use it

Do not take *Rasitin* if allergic to its active ingredient or other ingredients in it.

# - Before you start to use it

Inform the doctor or pharmacist if you are pregnant or breast feeding. Safety for use has not been established. Ask your doctor or pharmacist for advice before taking any medicine.

# - Taking other medicines

Tell your doctor if you are taking any other - Things to be careful of medicines. Also, including any medicine that you buy without a prescription from a pharmacy, supermarket or health food shop.

# How to use Rasitin

#### - How much to use

 $\overline{\text{Adults: oral, } 10 - 20\text{mg } (1 - 2 \text{ tablets})}$ . Children: 4 - 6 years: oral, 2.5 - 5mg. 7 - 15 years: oral, 5 to 10mg ( $\frac{1}{2}$  to 1 tablet) or as directed by physician. Follow all directions given by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or calling Tel: 03-78835550, or visiting the pharmacist for help.

### - When to use it

To be taken three times a day or as directed by the doctor or pharmacist.

### - How long to use it

Continue taking Rasitin for as long as your doctor recommends.

# - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - <u>If you use too much (overdose)</u>

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

# Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Rasitin.

# - Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist. Do not give Rasitin to anyone else, even if they have the same symptoms or condition as you.

Caution when driving or doing jobs requiring alertness because of possible drowsiness or dizziness. Avoid alcoholic beverages.

#### **Side effects**

Rasitin can cause side effects, although not everybody gets them. Visit the doctor or pharmacist immediately if you experience any side effects after taking this medicine. Drowsiness and dryness of mouth may occur. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by website npra.moh.gov.my

(Public → Reporting Medicinal Problems / Side Effects / AEFI / Vaccine Safety).

# Storage and Disposal of Rasitin

#### - <u>Storage</u>

Store in cool, dry place below 30°C. Protect from light. Keep out of reach of children.

#### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

# **Product description**

- What it looks like

White, oval, biconvex, top-scored tablet.

#### - Ingredients

- Active ingredient: cloperastine hydrochloride.
- Inactive ingredients: lactose monohydrate, cornstarch, polyvinylpyrrolidone and magnesium stearate.

# - MAL number:

MAL19930045AZ

# **Manufacturer and Product Registration** Holder

Prime Pharmaceutical Sdn. Bhd. 1505, Lorong Perusahaan Utama 1 Taman Perindustrian Bukit Tengah 14000 Bukit Mertajam, Penang, Malaysia.

# Date of revision

17/08/2017

# Serial Number

NPRA(R4/1)160817/00216