

Indication: For the relief of discomfort and pain, reduction of inflammation and to promote healing in the following conditions; ulcer, lesions and soreness in the gums, tongue, lips or palate; trauma or lesions in the oral cavity caused by new denture, denture sore spots and denture stomatitis; and teething disorders.

Contents: Triamcinolone acetoneide..... 0.1%w/v
Lignocaine HCl..... 2.5%w/v
Chlorhexidine HCl..... 0.5%w/v

Re-liv Oral Lotion contains chlorhexidine. Inform your healthcare provider if you have a known allergy to chlorhexidine.
Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.
Please refer enclosed leaflet for further details.

Manufactured & Distributed by:
PRIME PHARMACEUTICAL SDN. BHD.
1505, Lrg. Per Utara 1, Tmn. Perind. Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.



RE-LIV oral lotion

12 x 5ml

Store in a cool, dry place below 30°C.
Protect from light.
Keep away from children.

Simpan di tempat yang dingin dan kering bawah 30°C.
Jauhi dari cahaya.
Jauhkan dari kanak-kanak.

Controlled Medicine

Batch No. :

Mfg. Date :

Exp. Date :

Reg. No. : MAL06051304AZ



9 555701 007748

12 x 5ml



RE-LIV oral lotion