

RE-LIV ORAL LOTION

Lignocaine hydrochloride/ chlorhexidine hydrochloride/triamcinolone acetonide (2.5%w/v / 0.5%w/v / 0.1%w/v)

What is in this leaflet

1. What *Re-liv* is used for
2. How *Re-liv* works
3. Before you use *Re-liv*
4. How to use *Re-liv*
5. While you are using it
6. Side effects
7. Storage and disposal of *Re-liv*
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

What *Re-liv* is used for

For the relief of discomfort and pain, reduction of inflammation and promote healing in ulcers, lesions and soreness in the gums, tongue, lips or palate; trauma or lesions in the oral cavity caused by new denture, denture sore spots and denture stomatitis; and teething disorders.

How *Re-liv* works

Re-liv contains triamcinolone acetonide which is responsible for its anti-inflammatory effects, chlorhexidine hydrochloride which kills bacteria and lignocaine hydrochloride which blocks pain.

Before you use *Re-liv*

- When you must not use it

- Do not take *Re-liv* if you are
- allergic to its active ingredients or any other ingredients in it.
- having viral and other infection.

Pregnancy and lactation

Pregnancy is not a contraindication, but you should be aware of possible toxic effects to the fetus if there is systemic absorption. Ask your doctor or pharmacist for advice before using it. Risk-benefit must be considered in women of childbearing potential and particularly during early pregnancy or nursing mothers.

- Before you start to use it

- Inform your doctor or pharmacist if you have any of the following:
- tuberculosis (bacterial infection that mainly affects the lungs), intestinal ulcer or diabetes, especially long term usage.
- use in children, elderly, ill or debilitated individuals especially when extensive areas are treated.

- Taking other medicines

- Tell your doctor if you are taking:
- beta-adrenergic blocking agents (treat high blood pressure)
- cimetidine (treat stomach ulcer)
- pain killer such as NSAIDS, aspirin
- potassium depleting medicines (e.g. thiazides, frusemide, ethacrynic acid)
- ambinonium, neostigmine, pyridostigmine

How to use *Re-liv*

- How much to use

Instill 2 - 4 drops of lotion to the lesion using the tip of a clean cotton bud or finger. Rub gently to cover the lesion with lotion. Application may be repeated after 1 - 2 hours, preferably after meals and bedtime. Follow all directions given by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue using *Re-liv* for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using *Re-liv*

- Things you must do

- *Re-liv* contains chlorhexidine. Inform your healthcare provider if you have a known allergy to chlorhexidine. Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.
- Use your medicine exactly as your doctor has told you.
- Tell the doctors, dentists and pharmacists treating you that you are using *Re-liv*.
- If improvement does not occur in 7 days, please consult your doctor. It should be borne in mind that normal defensive responses of oral tissues are depressed in people receiving topical corticosteroid therapy.
- Tell your doctor immediately if you are pregnant while using this medicine.

- Things you must not do

- Do not chew gum or food while numbness persists because of risk of biting tongue or sides of the mouth.
- Do not give *Re-liv* to anyone else, even if they have the same symptoms or condition as you.
- Do not eat for one hour following use of medication because it may impair

swallowing leading to risk of aspiration.

- Things to be careful of

This product contains chlorhexidine hydrochloride and do not instill into ear.

Side effects

Like all medicines, *Re-liv* can cause side effects, although not everybody gets them. Visit the doctor or pharmacist immediately if you experience any side effects after taking this medicine. Signs of infection such as pain, redness or pus-containing blister and/or signs of irritation such as burning, itching, blistering or peelings not present before therapy, may require medical attention. Systemic absorption due to long term use may cause effects including Cushing's syndrome, high blood glucose levels and presence of glucose in the urine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nprra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of *Re-liv*

- Storage

Store in a cool dry place below 30°C. Protect from light. Keep out of reach and sight of children.

- Disposal

Medicines should not be disposed via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like

White flavoured oral lotion.

- Ingredients

- Active ingredients: lignocaine hydrochloride, chlorhexidine hydrochloride and triamcinolone acetonide
- Inactive ingredients: glycerin, propylene glycol, sodium carboxymethylcellulose, sodium saccharin, blackcurrant flavor and deionized water.

- MAL number:

MAL06051304AZ

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong Perusahaan Utama 1
Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

15/07/2024

Serial number

BPFK(R4/1)220617/00152