



CARBOSOLTABLET

CARBOSOL SYRUP

Contents

Carbosol Tablet	- Carbocysteine	375mg
Carbosol Syrup	- Carbocysteine	125mg/ 5ml
	Methylparaben	0.1% w/v
	Propylparaben	0.01%w/v

Indications:

For treatment of acute and chronic respiratory tract diseases accompanied by excessive or viscous mucus.

Pharmacology:

Carbocysteine, a mucoregulator, reduces sputum viscosity as well as protects the mucous membrane against infection and irritation.

Contraindications:

Contraindicated in children below 2 years of age.

Adverse Effects:

Nausea, headache, gastric discomfort, gastrointestinal bleeding and skin rash may occur occasionally.

Precautions:

- Caution in patients with history of peptic ulcer.
- Safety for use in pregnancy has not been established.

Dosage and Administration:

Adults - Oral, 750mg (2 tablets) three times a day, reduced to 375mg (1 tablet) three times a day when a response is obtained.

Children 2 to 5 years - Oral, 62.5 to 125mg (1/2 to 1 teaspoonful) four times a day.

6 to 12 years - Oral, 250mg (2 teaspoonfuls) three times a day.

Overdosage:

Clinical features include gastrointestinal discomfort, nausea and vomiting. Seek medical treatment immediately if symptoms persist.

Storage	: Keep container tightly closed. Protect from light. Store in a cool, dry place, below 30°C.
Presentation/ Packing	: Syrup : 125mg/5ml X 60 ml, 120ml Tablet 375mg x 50 x10's/aluminium/PVC blister pack
Shelf-life	: Tablet: 3 years Syrup: 2 years
Manufactured & Distributed by :	Prime Pharmaceutical Sdn. Bhd. Penang, Malaysia.