



# 安咳嗽

## AMCOSOL SYRUP 30mg/5ml



### DESCRIPTION:

Plain, cherry flavoured syrup.

### CONTENTS

Amcosol Syrup : Ambroxol HCL 30mg/5ml

Preservatives : Methylparabens 0.10% w/v and Propylparabens 0.01%w/v

### INDICATIONS

For treatment of acute or chronic bronchitis accompanied by abnormal bronchial secretions, and silicosis.

*Untuk bronkitis jangka panjang dan pendek dengan mukus berlebihan dan silikosis.*

**主治急性/慢性支应气管炎、多痰。**

### PHARMACOLOGY

Ambroxol, a metabolite of bromhexine, produces a marked reduction in coughing and dyspnoea as well as improvement in expectoration for patients with bronchitis. Its twofold action as a secretolytic and surfactant stimulant accounts for its activity in silicosis patients.

Elimination is mainly by renal excretion.

*Ambroxol mengurangkan batuk dan melegakan pemapasan dengan berkesan. Ia menyenangkan pengeluaran kahak berlebihan.*

**能有效治疗咳嗽,使呼吸顺畅及有化痰作用。**

### ADVERSE EFFECTS

Hypersensitivity reaction to ambroxol may occur.

Immune System Disorders: Frequency not known: Anaphylactic reaction including anaphylactic shock.

Skin and Subcutaneous Skin Disorders: Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

### WARNINGS AND PRECAUTIONS

Safety for use in pregnancy and lactation has not been established.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis(TEN), Erythema Multiforme(EM)and Acute Generalized Exanthematous Pustulosis(AGEP) have been reported.

In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

### DOSAGE AND ADMINISTRATION

For oral use

Adults and children above 12 yrs: 10ml (2 tsp.) twice daily.

Child 6-12 yrs: 5ml (1 tsp.) 2-3 times daily.

Child 2-6 yrs: 2.5ml (1/2 tsp.) 3 times daily.

Child 1-2 yrs: 2.5ml (1/2 tsp.) twice daily.

Dewasa: 10ml (2 sudu teh) dua kali sehari.

Kanak-kanak 6-12 tahun: 5ml (1 sudu teh) 2-3 kali sehari.

Kanak-kanak 2-6 tahun: 2.5ml (1/2 sudu teh) 3 kali sehari.

Kanak-kanak 1-2 tahun: 2.5ml (1/2 sudu teh) 2 kali sehari.

成人或儿童12岁以上: 10ml (2茶匙), 一天两次

儿童6-12岁: 5ml (1茶匙), 一天两~三次

儿童2-6岁: 2.5ml (半茶匙), 一天三次

儿童1-2岁: 2.5ml (半茶匙), 一天两次

Storage : Keep container tightly closed. Store in cool, dry place below 30°C. Protect from light.  
Keep out of reach of children.

Presentation / Packing : Syrup 30mg/ 5ml x 120ml ; 3.6L (EXPORT)

Shelflife : 3 years

Product Registration Holder : **PRIME PHARMACEUTICAL Sdn. Bhd.**

& Manufacturer : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,  
14000 Bukit Mertajam, Penang, Malaysia

12/08/2020