



PRIHEXINE TABLET 8mg/tab

Description: Yellow, round with top-scored tablet. Diameter = 7.00mm

PRIHEXINE SYRUP 4mg/5ml

Description: Yellow, banana flavoured syrup.

Contents

Prihexine Tablet 8mg	: Bromhexine HCl	8mg
Prihexine syrup 4mg/5ml	: Bromhexine HCl	4mg
	Methylparabens	0.1%w/v
	Propylparabens	0.01%w/v

Pharmacology:

Bromhexine changes the structure of bronchial secretions and increase the volume and reduce the viscosity of sputum. It loosen sputum and help expectoration.

Indications:

As mucolytic agent used in the treatment of acute and chronic respiratory tract diseases accompanied by excessive or viscous mucus.

Side Effects/Adverse Effects:

Nausea, vomiting, gastric irritation may occur occasionally.

Immune System Disorders: Frequency not known: Anaphylactic reactions including anaphylactic shock.
Skin and Subcutaneous Skin Disorders: Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

Warnings and Precautions:

Bromhexine should be given cautiously to patients with gastric ulceration.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Contraindications:

First trimester of pregnancy.

Recommended dosage:

For oral use

- Adults : 1 to 2 tablets to be taken 3 times daily.
Children : 1/2 - 1 teaspoonfuls 3 times daily.

Storage	: Keep container tightly closed. : Store in a cool, dry place, below 30°C. Protect from light. Keep out of reach of children.
Presentation / Packing	: Tablet 8mgx1000's ; 4mg/5mlx60ml
Shelf Life	: Tablet : 3years : Syrup : 2years
Product registration holder & Manufacturer:	: PRIME PHARMACEUTICAL SDN. BHD. 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah, 14000 Bukit Mertajam, Penang, Malaysia.

12/08/2020