



## **PRITARAX CREAM 10%**

### **Contents:**

Crotamiton B.P. 10% w/w  
Chlorocresol 0.1% w/w

### **Indications:**

For treatment of scabies and relief of pruritus associated with various dermatoses.

### **Pharmacology:**

Crotamiton, a scabidical and antipruritic agent, is highly effective in eradicating scabies infestations and relief of pruritus ani, vulvae, and scroti, senile pruritus, and pruritus associated with various dermatoses.

One application is usually effective for 6 to 10 hours.

### **Adverse Effects:**

Allergicsensitivity or primary irritation reactions may occur in some patients.

### **Precautions:**

Avoid contact with the eyes and mucous membranes.

Check with physician if condition does not improve or if it becomes worse.

Treatment with crotamiton should be discontinued if severe irritation or sensitization develops.

Safety for use in pregnancy and lactation has not been established.

### **Contraindications:**

Avoid in patients who are hypersensitive to crotamiton.

It should not be used in the presence of acute exudative or versicular dermatitis.

### **Dosage and Administrations:**

For pruritus - Topical, to the skin. Repeat as needed.

For scabies - Topical, to the skin, as a 10% cream for two applications at 24-hour intervals. (Do not wash off the first coat of medication)

The cream should be rubbed well and enough to cover the entire skin surface from the chin down, especially to body folds and creases. A cleansing bath should be taken 48 hours after the second application. Patient should put on freshly washed or dry-cleaned clothing and change bedding to prevent reinfestation.

Note: In resistant cases of scabies, the course of treatment may be repeated one week later.

### **Overdosage:**

If accidentally ingested, gastric lavage should be performed.

**Storage** : Keep container tightly closed. Store in a cool, dry place, below 30C.  
Protect from light. Keep away from children.

**Presentation/Packing** : Cream 10% x 10gm

**Shelflife** : 3 years

**Manufactured &** : PRIME PHARMACEUTICAL SDN.BHD.

**Distributed by** : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,  
14000 Bukit Mertajam, Penang, Malaysia.