

HEXIDERM CREAM

Chlorhexidine hydrochloride/tolnaftate (0.1%w/w / 1%w/w)

What is in this leaflet

1. What Hexiderm is used for
2. How Hexiderm works
3. Before you use Hexiderm
4. How to use Hexiderm
5. While you are using Hexiderm
6. Side effects
7. Storage and Disposal of Hexiderm
8. Product description
9. Manufacturer and Product Registration holder
10. Date of revision

What Hexiderm is used for

It is used for skin disinfection and treatment of fungal infections of the skin of the foot (athlete's foot), groin, arms, legs and hand.

How Hexiderm works

It contains tolinaftate used for the treatment of ringworm and skin infections caused by certain fungi, and chlorhexidine which kills bacteria.

Before you use Hexiderm

- When you must not use it
Hexiderm contains chlorhexidine. Inform your healthcare provider if you have a known allergy to chlorhexidine.
- Before you start to use it
Avoid contact with the eyes.
- Taking other medicines
The medicine is for topical application and will usually have minimal effect on other medication. However, please inform the doctor or pharmacist if using other medicines.

How to use Hexiderm

- How much to use
Adults and children - apply to the skin.
- When to use it
Apply two times a day or follow the advice of the doctor or pharmacist.
- How long to use it
You should check with your doctor or pharmacist.
- If you forget to use it
If you forget to use this medicine, use it as soon as you remember. Then continue using it at your usual time.

- If you use too much (overdose)

If you have used too much Hexiderm, contact your doctor, an emergency department or a poison center immediately.

While you are using Hexiderm

- Things you must do
Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.
- Things you must not do
Do not stop using medication although you feel better after a few days, until the prescribed course is finished.
- Things to be careful of
If you are pregnant, become pregnant, or breastfeeding, please inform the doctor or pharmacist. Safety for use in pregnancy and breastfeeding has not been established.

Side effects

Skin reactions including irritation and itchiness may occur. Skin inflammation after contact with Hexiderm has been reported. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Hexiderm

- Storage
Store in a dry place below 30°C. Protect from light. Keep out of reach of children.
- Disposal
Ask your pharmacist how to dispose of medicines no longer required.

Product description

- What it looks like
White colour cream.

- Ingredients:

- Active ingredients: tolinaftate and chlorhexidine hydrochloride.
- Inactive ingredients: white soft paraffin, liquid paraffin, cetomacrogol emulsifying wax and deionised water.

- MAL number:

MAL19930279X

Manufacturer and Product Registration Holder

Prime Pharmaceutical Sdn. Bhd.
1505, Lorong Perusahaan Utama 1
Taman Perindustrian Bukit Tengah
14000 Bukit Mertajam, Penang, Malaysia.

Date of revision

23/09/2020

Serial Number

NPRA(R4/1)221118/00349