



Trabest 37.5mg/325mg Tablets

Each film coated tablet contains:
Tramadol Hydrochloride37.5 mg
Paracetamol325 mg

Pharmaceutical dosage form and Description
Film Coated Tablet
Pale yellow, capsule shaped, film-coated tablet, debossed with "TP" on one side and plain on the other side.

Clinical particulars
Therapeutic indications
Tramadol and Paracetamol tablets are indicated for the management of moderate to severe pain.

Posology and method of administrations
Posology:
Tramadol and Paracetamol should under no circumstances be administered for longer than is strictly necessary. If repeated use or long term treatment with Tramadol and Paracetamol is required as a result of the nature and severity of the illness, then careful, regular monitoring should take place (with breaks in the treatment, where possible), to assess whether continuation of the treatment is necessary.
Adults and Children Over 16 Years
The maximum single dose of Tramadol and Paracetamol is 1 to 2 tablets every 4 to 6 hours as needed for pain relief up to a maximum of 8 tablets per day.
Tramadol and Paracetamol can be administered without regard to food.
Pediatric (Children Below 16 Years):
The effective and safe use of Tramadol and Paracetamol has not been established in children below the age of 16 years. Treatment is therefore not recommended in this population.
Elderly (Geriatric) patients:
The usual dosages may be used, although it should be noted that in volunteers aged over 75 years, the elimination half life of tramadol was increased by 17% following oral administration. In patients over 75 years old, it is recommended that the minimum interval between doses should be not less than 6 hours, due to the presence of tramadol.
Method of administration:
Oral use.
Tablets must be swallowed whole, with a sufficient quantity of liquid. They must not be broken or chewed.

Contra-indications
Hypersensitivity to tramadol, paracetamol or to any of the excipients of the medicinal product.
Acute intoxication with alcohol, hypnotic drugs, centrally-acting analgesics, opioids or psychotropic drugs.
Co-administration to patients who are receiving monoamine oxidase inhibitors or within two weeks of their withdrawal.
Severe hepatic impairment.
Epilepsy not controlled by treatment.

Special warning and precautions for use
Warnings:

This preparation contains PARACETAMOL.
Do not take any other paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away. In adults and adolescents 16 years and older, the maximum dose of 8 tablets of Tramadol and Paracetamol should not be exceeded. In order to avoid inadvertent overdose, patients should be advised not to exceed the recommended dose and not to use any other paracetamol (including over the counter) or tramadol hydrochloride containing products concurrently without the advice of a physician.

In severe renal insufficiency (creatinine clearance <10 ml/min), Tramadol and Paracetamol is not recommended.
In patients with severe hepatic impairment Tramadol and Paracetamol should not be used. The hazards of paracetamol overdose are greater in patients with non-cirrhotic alcoholic liver disease. In moderate cases, prolongation of dosage interval should be carefully considered.
In severe respiratory insufficiency, Tramadol and Paracetamol is not recommended.
Tramadol is not suitable as a substitute in opioid-dependent patients. Although it is an opioid agonist, tramadol cannot suppress morphine withdrawal symptoms.
Convulsions have been reported in tramadol-treated patients susceptible to seizures and/or taking other medications that lower the seizure threshold, especially selective serotonin re-uptake inhibitors, tricyclic antidepressants, antipsychotics, centrally acting analgesics or local anaesthesia. Epileptic patients controlled by a treatment or patients susceptible to seizures should be treated with Tramadol and Paracetamol only if there are compelling circumstances. Convulsions have been reported in patients receiving tramadol at the recommended dose levels. The risk may be increased when doses of tramadol exceed the recommended upper dose limit.
Concomitant use of opioid agonists-antagonists (nalbuphine, buprenorphine, pentazocine) is not recommended.

Precautions for use:
Tramadol and Paracetamol should be used with caution in opioid dependent patients, or in patients with cranial trauma, in patients prone to convulsive disorder, biliary tract disorders, in a state of shock, in an altered state of consciousness for unknown reasons, with problems affecting the respiratory centre or the respiratory function, or with an increased intracranial pressure.
Paracetamol in overdosage may cause hepatic toxicity in some patients.
Tolerance and physical and/or psychological dependence may develop, even at therapeutic doses. The clinical need for analgesic treatment should be reviewed regularly. In opioid-dependent patients and patients with a history of drug abuse or dependence, treatment should only be for short period and under medical supervision.
Symptoms of withdrawal reaction, similar to those occurring during opiate withdrawal, may occur even at therapeutic doses and for short term treatment. Withdrawal symptoms may be avoided by taper it at the time of discontinuation especially after long treatment periods. Rarely, cases of dependence and abuse have been reported.

Interaction with other drugs, other forms of interactions
Concomitant use is contraindicated with:

- Non-selective MAO Inhibitors
- Risk of serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
- Selective A-MAO Inhibitors
- Extrapolation from non-selective MAO inhibitors.
- Risk of serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
- Selective B-MAO Inhibitors
- Central excitation symptoms evocative of a serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
- In case of recent treatment with MAO inhibitors, a delay of two weeks should occur before treatment with tramadol.

Concomitant use is not recommended with:

- Alcohol

Alcohol increases the sedative effect of opioid analgesics.
The effect on alertness can make driving of vehicles and the use of machines dangerous.
Avoid intake of alcoholic drinks and of medicinal products containing alcohol.
Carbamazepine and other enzyme inducers.
Risk of reduced efficacy and shorter duration due to decreased plasma concentrations of tramadol.

- Opioid agonists-antagonists (buprenorphine, nalbuphine, pentazocine)

Decrease of the analgesic effect by competitive blocking effect at the receptors, with the risk of occurrence of withdrawal syndrome.
Concomitant use which needs to be taken into consideration:
In isolated cases, there have been reports of Serotonin Syndrome in a temporal connection with the therapeutic use of tramadol in combination with other serotonergic medicines such as selective serotonin re-uptake inhibitors (SSRIs) and triptans. Signs of Serotonin Syndrome may be for example, confusion, agitation, fever, sweating, ataxia, hyperreflexia, myoclonus and diarrhoea.
Other opioid derivatives (including antitussive drugs and substitutive treatments), barbiturates and benzodiazepines.
Increased risk of respiratory depression which can be fatal in cases of overdose
Other central nervous system depressants, such as other opioid derivatives (including antitussive drugs and substitutive treatments), barbiturates, benzodiazepines, other anxiolytics, hypnotics, sedative antidepressants, sedative antihistamines, neuroleptics, centrally-acting antihypertensive drugs, thalidomide and baclofen.
These drugs can cause increased central depression. The effect on alertness can make driving of vehicles and the use of machines dangerous.
As medically appropriate, periodic evaluation of prothrombin time should be performed when Tramadol and Paracetamol and warfarin like compounds are administered concurrently due to reports of increased INR.
Other drugs known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation), and probably also the metabolism of the active O-demethylated metabolite. The clinical importance of such an interaction has not been studied.
Medicinal products reducing the seizure threshold, such as bupropion, serotonin reuptake inhibitor antidepressants, tricyclic antidepressants and neuroleptics. Concomitant use of tramadol with these drugs can increase the risk of convulsions. The speed of absorption of paracetamol may be increased by metoclopramide or domperidone, and absorption reduced by cholestyramine.
In a limited number of studies the pre- or postoperative application of the antiemetic 5-HT3 antagonist ondansetron increased the requirement of tramadol in patients with postoperative pain.

Use in pregnancy and lactation:
Pregnancy:
Since Tramadol and Paracetamol is a fixed combination of active ingredients including tramadol, it should not be used during pregnancy.
Data regarding paracetamol:
Epidemiological studies in human pregnancy have shown no ill effects due to paracetamol used in the recommended dosages.
Data regarding tramadol:
Tramadol should not be used during pregnancy as there is inadequate evidence available to assess the safety of tramadol in pregnant women. Tramadol administered before or during birth does not affect uterine contractility. In neonates, it may induce changes in the respiratory rate which are usually not clinically relevant. Long-term treatment during pregnancy may lead to withdrawal symptoms in the newborn after birth, as a consequence of habituation.

Breastfeeding:
Since Tramadol and Paracetamol is a fixed combination of active ingredients including tramadol, it should not be ingested during breast feeding.
Data regarding paracetamol:
Paracetamol is excreted in breast milk but not in a clinically significant amount. Available published data do not contraindicate breast feeding by women using single ingredient medicinal products containing only paracetamol.
Data regarding tramadol:
Tramadol and its metabolites are found in small amounts in human breast milk. An infant could ingest about 0.1% of the dose given to the mother. Tramadol should not be taken during breast feeding.

Side effects:
The most commonly reported undesirable effects with the paracetamol/tramadol combination were nausea, dizziness and somnolence.

Cardiovascular system disorders:
Uncommon: hypertension, palpitations, tachycardia, arrhythmia
Central and peripheral nervous system disorders:
Very common: dizziness, somnolence
Common: headache, trembling
Uncommon: involuntary muscular contractions, paraesthesia, tinnitus
Rare: ataxia, convulsions.

Psychiatric disorders:
Common: confusion, mood changes (anxiety, nervousness, euphoria), sleep disorders
Uncommon: depression, hallucinations, nightmares, amnesia
Rare: drug dependence
Post marketing surveillance:
Very rare: abuse.

Vision disorders:
Rare: blurred vision

Respiratory system disorders:
Uncommon: dyspnoea

Gastro-intestinal disorders:
Very common: nausea
Common: vomiting, constipation, dry mouth, diarrhoea abdominal pain, dyspepsia, flatulence
Uncommon: dysphagia, melaena.

Liver and biliary system disorders:
Uncommon: hepatic transaminases increase.

Skin and appendages disorders:
Common: sweating, pruritus
Uncommon: dermal reactions (e.g. rash, urticaria).

Urinary system disorders:
Uncommon: albuminuria, micturition disorders (dysuria and urinary retention).

Body as a whole:
Uncommon: shivers, hot flushes, thoracic pain.
The occurrence of the following undesirable effects known to be related to the administration of tramadol or paracetamol cannot be excluded:

Market/Customer :	Malaysia	Location :	Pithampur
Minimum Font Size :	6 pt	Tracking No. :	04
Product name :	LEAFLET TRABEST TAB 37.5/325MG (MY)		
Material code :	XXXXXX	Supersedes code :	265634
Open Size :	190 x 390 mm (L X W)	Pharmacode value :	XXXX
Folded Size :	50 x 30 mm (L x W)	Component :	Leaflet
No. of folds :	----	GSM :	40 gsm Bible Paper
Pantone Colours :	Black Black	Gluing :	YES
Reason for Change:	Shelf life changed to 36months from 24 months. Importer is change from Biocare to Unimed SDN BHD		
Unicorn Creation D/Lupin/Export/Malaysia/Trabest/ Date: 18/08/2022			

50 mm

390 mm

190 mm

Tramadol
Postural hypotension, bradycardia, collapse (tramadol).
Post-marketing surveillance of tramadol has revealed rare alterations of warfarin effect, including elevation of prothrombin times.
Rare cases: allergic reactions with respiratory symptoms (e.g. dyspnoea, bronchospasm, wheezing, angioneurotic oedema) and anaphylaxis.
Rare cases: changes in appetite, motor weakness, and respiratory depression.
Psychic side-effects may occur following administration of tramadol which vary individually in intensity and nature (depending on personality and duration of medication). These include changes in mood, (usually elation, occasionally dysphoria), changes in activity (usually suppression, occasionally increase) and changes in cognitive and sensorial capacity (e.g. decision behaviour perception disorders).
Worsening of asthma has been reported though a causal relationship has not been established.
Symptoms of withdrawal reactions, similar to those occurring during opiate withdrawal, may occur as follows: agitation, anxiety, nervousness, insomnia, hyperkinesia, tremor and gastrointestinal symptoms. Other symptoms that have very rarely been seen if tramadol hydrochloride is discontinued abruptly include: panic attacks, severe anxiety, hallucinations, paraesthesia, tinnitus and unusual CNS symptoms.

Paracetamol
Adverse effects of paracetamol are rare. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported. There have been reports of blood dyscrasias including thrombocytopenia and agranulocytosis, but these were not necessarily causally related to paracetamol. There have been several reports that suggest that paracetamol may produce hypoprothrombinemia when administered with warfarin-like compounds. In other studies, prothrombin time did not change.

Overdoses:
Tramadol and Paracetamol is a fixed combination of active ingredients. In case of overdose, the symptoms may include the signs and symptoms of toxicity of tramadol or paracetamol or of both these active ingredients.
Symptoms of overdose from tramadol:
In principle, on intoxication with tramadol, symptoms similar to those of other centrally acting analgesics (opioids) are to be expected. These include in particular, miosis, vomiting, cardiovascular collapse, consciousness disorders up to coma, convulsions and respiratory depression up to respiratory arrest.
Symptoms of overdose from paracetamol:
An overdose is of particular concern in young children. Symptoms of paracetamol overdose in the first 24 hours are pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, coma and death. Acute renal failure with tubular necrosis may develop, even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.
Liver damage is possible in adults who have taken 7.5-10 g or more of paracetamol. It is considered that excess quantities of a toxic metabolite (usually adequately detoxified by glutathione when normal doses of paracetamol are ingested) become irreversibly bound to liver tissue.
Emergency treatment:
Transfer immediately to a specialised unit.
Maintain respiratory and circulatory function.
Prior to starting treatment, a blood sample should be taken as soon as possible after overdose in order to measure the plasma concentration of paracetamol and tramadol and in order to perform hepatic tests
Perform hepatic tests at the start (of overdose) and repeat every 24 hours. An increase in hepatic enzymes (ASAT,ALAT) is usually observed, which normalizes after one or two weeks.
Empty the stomach by causing the patient to vomit (when the patient is conscious) by irritation or gastric lavage.
Supportive measures such as maintaining the patency of the airway and maintaining cardiovascular function should be instituted; naloxone should be used to reverse respiratory depression; fits can be controlled with diazepam.
Tramadol is minimally eliminated from the serum by haemodialysis or haemofiltration. Therefore, treatment of acute intoxication with Tramadol and Paracetamol with haemodialysis or haemofiltration alone is not suitable for detoxification.
Immediate treatment is essential in the management of paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention. Any adult or adolescent who had ingested around 7.5 g or more of paracetamol, or any child who has ingested 150 mg/kg of paracetamol in the preceding 4 hours, should undergo gastric lavage. Paracetamol concentrations in blood should be measured later than 4 hours after overdose in order to be able to assess the risk of developing liver damage (via the paracetamol overdose nomogram). Administration of oral methionine or intravenous N-acetylcysteine (NAC), which may have a beneficial effect up to at least 48 hours after the overdose, may be required. Administration of intravenous NAC is most beneficial when initiated within 8 hours of overdose ingestion. However, NAC should still be given if the time to presentation is greater than 8 hours after overdose and continued for a full course of therapy. NAC treatment should be started immediately when massive overdose is suspected. General supportive measures must be available.
Irrespective of the reported quantity of paracetamol ingested, the antidote for paracetamol, NAC, should be administered orally or intravenously, as quickly as possible, if possible, within 8 hours following the overdose.

Pharmacological properties:
Pharmaco-kinetic properties:
Tramadol is administered in racemic form, and the [-] and [+] forms of tramadol and its metabolite M1 are detected in the blood. Although tramadol is rapidly absorbed after administration, its absorption is slower (and its half-life longer) than that of paracetamol.
After a single oral administration of a tramadol/paracetamol (37.5 mg + 325 mg) tablet, peak plasma concentrations of 64.3/55.5 ng/ml [(+)-tramadol/(-)-tramadol] and 4.2 µg/ml (paracetamol) are reached after 1.8 h [(+)-tramadol/(-)-tramadol] and 0.9 h (paracetamol) respectively. The mean elimination half-lives t1/2 are 5.1/4.7 h [(+)-tramadol/(-)-tramadol] and 2.5 h (paracetamol).
During pharmacokinetic studies in healthy volunteers after single and repeated oral administration of Tramadol and Paracetamol, no clinical significant change was observed in the kinetic parameters of each active ingredient compared to the parameters of the active ingredients used alone.
Absorption:
Racemic tramadol is rapidly and almost completely absorbed after oral administration. The mean absolute bioavailability of a single 100 mg dose is approximately 75%. After repeated administration, the bioavailability is increased and reaches approximately 90%.
After administration of Tramadol and Paracetamol, the oral absorption of paracetamol is rapid and nearly complete and takes place mainly in the small intestine. Peak plasma concentrations of paracetamol are reached in 1 hour and are not modified by concomitant administration of tramadol.
The oral administration of Tramadol and Paracetamol with food has no significant effect on the peak plasma concentration or extent of absorption of either tramadol or paracetamol so that Tramadol and Paracetamol can be taken independently of meal times.
Distribution:
Tramadol has a high tissue affinity (Vd,β = 203 ± 40 l). It has a plasma protein binding of about 20%.
Paracetamol appears to be widely distributed throughout most body tissues except fat. Its apparent volume of distribution is about 0.9 l/kg. A relatively small portion (~20%) of paracetamol is bound to plasma proteins.
Metabolism:
Tramadol is extensively metabolized after oral administration. About 30 % of the dose is excreted in urine as unchanged drug, whereas 60% of the dose is excreted as metabolites. Tramadol is metabolised through O-demethylation (catalysed by the enzyme CYP2D6) to the metabolite M1, and through N-demethylation (catalysed by CYP3A) to the metabolite M2. M1 is further metabolised through N-demethylation and by conjugation with glucuronic acid. The plasma elimination half-life of M1 is 7 hours. The metabolite M1 has analgesic properties and is more potent than the parent drug. The plasma concentrations of M1 are several-fold lower than those of tramadol and the contribution to the clinical effect is unlikely to change on multiple dosing.
Paracetamol is principally metabolized in the liver through two major hepatic routes: glucuronidation and sulphation. The latter route can be rapidly saturated at doses above the therapeutic doses. A small fraction (less than 4%) is metabolized by cytochrome P 450 to an active intermediate (the N-acetyl benzoquinoneimine) which, under normal conditions of use, is rapidly detoxified by reduced glutathione and excreted in urine after conjugation to cysteine and mercapturic acid. However, during massive overdose, the quantity of this metabolite is increased.
Elimination:
Tramadol and its metabolites are eliminated mainly by the kidneys. The half-life of paracetamol is approximately 2 to 3 hours in adults. It is shorter in children and slightly longer in the newborn and in cirrhotic patients. Paracetamol is mainly eliminated by dose-dependent formation of glucuro- and sulpho-conjugate derivatives. Less than 9 % of paracetamol is excreted unchanged in urine. In renal insufficiency, the half-life of both compounds is prolonged.

Pharmaco-dynamic properties:
Pharmacotherapeutic group: Tramadol, combinations.
ATC code: N02AX 52

Analgesics
Tramadol is an opioid analgesic that acts on the central nervous system. Tramadol is a pure non-selective agonist of the µ, δ, and κ opioid receptors with a higher affinity for the µ receptors. Other mechanisms which contribute to its analgesic effect are inhibition of neuronal reuptake of noradrenaline and enhancement of serotonin release. Tramadol has an antitussive effect. Unlike morphine, a broad range of analgesic doses of tramadol has no respiratory depressant effect. Similarly, the gastro-intestinal motility is not modified. The cardiovascular effects are generally slight. The potency of tramadol is considered to be one-tenth to one-sixth that of morphine.
The precise mechanism of the analgesic properties of paracetamol is unknown and may involve central and peripheral effects.
Tramadol and Paracetamol is positioned as a step II analgesic in the WHO pain ladder and should be utilised accordingly by the physician

Incompatibilities
Concomitant use is contraindicated with:
Non-selective MAO Inhibitors
Risk of serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
Selective A-MAO Inhibitors
Extrapolation from non-selective MAO inhibitors.
Risk of serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
Selective B-MAO Inhibitors
Central excitation symptoms evocative of a serotonergic syndrome: diarrhoea, tachycardia, sweating, trembling, confusion, even coma.
In case of recent treatment with MAO inhibitors, a delay of two weeks should occur before treatment with tramadol.
Concomitant use is not recommended with:
Alcohol
Alcohol increases the sedative effect of opioid analgesics.
The effect on alertness can make driving of vehicles and the use of machines dangerous.
Avoid intake of alcoholic drinks and of medicinal products containing alcohol.
Carbamazepine and other enzyme inducers.
Risk of reduced efficacy and shorter duration due to decreased plasma concentrations of tramadol.
Opioid agonists-antagonists (buprenorphine, nalbuphine, pentazocine)
Decrease of the analgesic effect by competitive blocking effect at the receptors, with the risk of occurrence of withdrawal syndrome.
Concomitant use which needs to be taken into consideration:
In isolated cases, there have been reports of Serotonin Syndrome in a temporal connection with the therapeutic use of tramadol in combination with other serotonergic medicines such as selective serotonin re-uptake inhibitors (SSRIs) and triptans. Signs of Serotonin Syndrome may be for example, confusion, agitation, fever, sweating, ataxia, hyperreflexia, myoclonus and diarrhoea.
Other opioid derivatives (including antitussive drugs and substitutive treatments), barbiturates and benzodiazepines.
Increased risk of respiratory depression which can be fatal in cases of overdose.
Other central nervous system depressants, such as other opioid derivatives (including antitussive drugs and substitutive treatments), barbiturates, benzodiazepines, other anxiolytics, hypnotics, sedative antidepressants, sedative antihistamines, neuroleptics, centrally-acting antihypertensive drugs, thalidomide and baclofen.
These drugs can cause increased central depression. The effect on alertness can make driving of vehicles and the use of machines dangerous.
As medically appropriate, periodic evaluation of prothrombin time should be performed when Tramadol and Paracetamol and warfarin like compounds are administered concurrently due to reports of increased INR.
Other drugs known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation), and probably also the metabolism of the active O-demethylated metabolite. The clinical importance of such an interaction has not been studied.
Medicinal products reducing the seizure threshold, such as bupropion, serotonin reuptake inhibitor antidepressants, tricyclic antidepressants and neuroleptics. Concomitant use of tramadol with these drugs can increase the risk of convulsions. The speed of absorption of paracetamol may be increased by metoclopramide or domperidone, and absorption reduced by cholestyramine.
In a limited number of studies the pre- or postoperative application of the antiemetic 5-HT3 antagonist ondansetron increased the requirement of tramadol in patients with postoperative pain.

Shelf-life
Three years

Special precautions for storage
Store below 30° C.

Nature and contents of container
10's PVC/PVdC Blister

Date of Revision of Text
August 2022

Product Registration Holder:
Generic Health Sdn Bhd,
Upper Penthouse, Wisma RKT, No. 2, Jalan Raja Abdullah,
Off Jalan Sultan Ismail, 50300 Kuala Lumpur.

Imported & Marketed By:
UNIMED SDN BHD,
NO.53, JALAN TEMBAGA SD5/2B, BANDAR SRI DAMANSARA,
52200, KUALA LUMPUR, MALAYSIA.

 Manufactured by:
LUPIN Limited
Plot No. M-2 & M-2-A, SEZ Phase-II, MISC Zone,
Apparel Park, Pithampur, Distt. Dhar – 454775.

Market / Customer :	Malaysia	Location :	Pithampur
Minimum Font Size :	6 pt	Tracking No. :	04
Product name :	LEAFLET TRABEST TAB 37.5/325MG (MY)		
Material code :	XXXXXX	Supersedes code :	265634
Open Size :	190 x 390 mm (L X W)	Pharmacode value :	XXXX
Folded Size :	50 x 30 mm (L x W)	Component :	Leaflet
No. of folds :	----	GSM :	40 gsm Bible Paper
Pantone Colours :	Black Black	Gluing :	YES
Reason for Change:	Shelf life changed to 36months from 24 months. Importer is change from Biocare to Unimed SDN BHD		
Unicorn Creation D/Lupin/Export/Malaysia/Trabest/ Date: 18/08/2022			