DISOL TABLETS

Bromhexine hydrochloride (8mg)

What is in this leaflet

1. What DISOL is used for
2. How DISOL works
3. Before you use DISOL
4. How to use DISOL
5. While you are using it
6. Side effects
7. Storage and disposal of DISOL
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What DISOL is used for

DISOL is used for the relief of respiratory conditions with stubborn or excessive phlegm.

How DISOL works

Bromhexine Hydrochloride helps to liquefy and loosen thick phlegm in the airway.

Before you use DISOL

- *When you must not use it*

  Do not take DISOL if you or your children are allergic to bromhexine or any other ingredients of this medicine (listed in Product Description).

- *Before you start to use it*

  Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine. It should not be used during first three months of pregnancy or lactation unless advised by the doctor.

- *Taking other medicines*

  Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

  Medications that may interfere with DISOL such as antibiotic.

How to use DISOL

- *How much to use*
  - Adults and children > 12 years: 1 tablet, to be taken three times daily.
  - Children between 6 – 12 years: ½ tablet, to be taken three times daily.
  - Children between 2 – 6 years: ½ tablet, to be taken two times daily.
  - It may be necessary to increase the total daily dose up to 6 tablets in adults.

- *When to use it*

  DISOL should be taken 2-3 times daily or as directed by your doctor or pharmacist.

- *How long to use it*

  Use DISOL for the duration your doctor recommends.

- *If you forgot to use it*

  Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- *If you use too much (overdose)*

  Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- *Things you must do*

  Tell all the doctors, dentists and pharmacists treating you that you are using DISOL.

- *Things you must not do*

  Do not use any new medicines without consulting your doctor or pharmacist.

  Do not give DISOL to anyone else, even if they have the same symptoms or condition as you.

- *Things to be careful of*

  This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

DISOL may cause severe allergy and serious skin reactions. Stop using DISOL and seek medical assistance immediately if you experience any of the following symptoms:

1. Severe allergy: breathing difficulties, light headedness, skin swellings or rash
2. Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicine (ConSERF) or Vaccines (AEFI)]

DISOL TABLETS

Bromhexine hydrochloride (8mg)

Storage and disposal of DISOL

- *Storage*
  Store below 30°C.

- *Disposal*
  Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*
  Light yellow, round and convex tablet with Siam logo on one side and scored on the other.

- *Ingredients*
  - Active ingredient
    Bromhexine hydrochloride

  - Inactive ingredient
    Lactose monohydrate, starch, sodium starch glycolate, magnesium stearate, quinolone yellow.

- *MAL number*
  MAL19913329XZ

Manufacturer

SIAM BHEASACH CO., LTD.
123 Soi Chokechairuammitr, Vibhavadi-Rangsit Road, Chomphon, Chatuchak, Bangkok 10900.

Product Registration Holder

SPG PHARMA (MALAYSIA) SDN. BHD.,
Suite 1701, Block A, Damansara Intan e-Business Park,
No. 1, Jalan SS20/27, 47400 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

Date of Revision

04/06/2024

Serial Number

NPRA(R4/1)040624/00065