

LORSEDIN[®] ORAL SOLUTION

Loratadine (1mg/ml)

What is in this leaflet

1. What LORSEDIN[®] is used for
2. How LORSEDIN[®] works
3. Before you use LORSEDIN[®]
4. How to use LORSEDIN[®]
5. While you are using it
6. Side effects
7. Storage and Disposal of LORSEDIN[®]
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What LORSEDIN[®] is used for

LORSEDIN[®] is used to relieve symptoms of sneezing, runny nose, itching; red, itchy, and watery eyes. It is also indicated for chronic urticaria (red, raised, itchy rash) and other allergic skin disorder.

How LORSEDIN[®] works

LORSEDIN[®] belongs to a class of medications called antihistamines. It works by blocking the action of histamines, a substance in the body that causes allergic reaction: sneezing, itching, watery eyes, and running nose.

Before you use LORSEDIN[®]

-When you must not use it

Do not use LORSEDIN[®] if:

- you are allergic to loratadine or any of the ingredients listed under the Ingredients section of this leaflet.

-Before you start to use it

You must tell your healthcare professional if you:

- have liver problems

Pregnancy and breastfeeding:

Talk to your doctor if you are pregnant, might become pregnant or are breastfeeding before taking this medicine.

Loratadine is excreted into breast milk and due to the increased risk of antihistamine for infants, particularly new born and premature infants, a decision should be made whether to discontinue nursing or discontinue the drug.

-Taking other medicines

Tell your healthcare professional if you are taking any other medicines, including medicines that you buy without a prescription from a pharmacy, supermarket or health food shop.

Medications which affect the liver should be used with caution.

How to use LORSEDIN[®]

-How much to take

Adults and Children 12 years of age and over:

Two 5 ml spoonfuls (10 mg) once daily.

Children 2 to 12 years of age:

Body Weight >30 kg -- two 5 ml spoonfuls (10 mg) syrup once daily.

Body Weight <30 kg -- One 5 ml spoonful (5 mg) syrup once daily.

-When to take it

Take LORSEDIN[®] as directed by the doctor or pharmacist.

-How long to take it

LORSEDIN[®] should be taken as directed by the doctor.

-If you forget to take it

Take the missed dose as soon as you remember. If it is approaching time for the next dose, skip the missed dose and take the medicine at your next regularly scheduled time. Do not take 2 doses at once.

-If you take too much (overdose)

Immediately inform your doctor or go to the Emergency Department of your nearest hospital if you suspect that you or anyone else may have taken too much of LORSEDIN[®].

Do this even if you are not experiencing signs of discomfort or poisoning as you may need urgent medical attention.

Symptoms of overdose: drowsiness or sleepiness, abnormally rapid heart rate, and headache.

LORSEDIN[®] ORAL SOLUTION

Loratadine (1mg/ml)

While you are taking LORSEDIN[®]

-Things you must do

- Inform your healthcare professional if you become pregnant while taking LORSEDIN[®].
- If you are about to start taking a new medicine, inform your healthcare professional that you are taking LORSEDIN[®].

-Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not give this medicine to anyone else, even if they have similar symptoms as you.

-Things to be careful of

Do not take charge of vehicles or machinery until the effect of the medicine treatment on the individual is known.

Side effects

If you get any of these serious side effects, stop taking this medicine. See a doctor at once:

- Difficulty in breathing, swelling of the face, neck, tongue or throat (severe allergic reactions)

If you get any of the following side effects see your pharmacist or doctor:

- Headache, nervousness, tiredness in children
- Drowsiness, headache, increased appetite,

difficulty in sleeping in adults

- Fast or irregular heart rate, dizziness, tiredness
- Feeling sick, dry mouth, stomach pain, abnormal liver function, rash, hair loss

This is not a complete list of all possible side effects. Inform your healthcare professional if you notice anything making you feel unwell when you are taking, or soon after you have finished taking LORSEDIN[®].

“You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website np.ra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI)”

Storage and Disposal of LORSEDIN[®]

-Storage

- Keep LORSEDIN[®] in a cool dry place below 30°C.
- Keep out of reach of children.

-Disposal

Consult your healthcare professional for disposal of leftover that have passed their expiry date or if you have been instructed to stop the medicine. Do not dispose via wastewater or household waste.

Product Description

- What it looks like

LORSEDIN[®] is a clear colorless solution with orange flavor.

-Ingredients

Active ingredient:
Loratadine

Inactive ingredients:
Purified water, orange sweet, sodium benzoate, sucralose, Edetate disodium (dihydrate) Citric acid monohydrate, hydroxypropyl cellulose, and propylene glycol.

- MAL number:
MAL18086161AZ

Manufacturer

Siam Bheasach Co., Ltd., 123 Soi Chokechairuammitr, Vibhavadi-Rangsit Road, Chomphon, Chatuchak, Bangkok 10900 and 9 Soi Chokechairuammitr 3, Vibhavadi-Rangsit Road, Dindang, Dindang, Bangkok 10400, Thailand.

Product Registration Holder

SPG Pharma (Malaysia) Sdn. Bhd., Suite 1701, Block A, Damansara intan e-Business Park, No.1, Jalan SS20/27, 47400 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

Date of revision

13/11/2024

Serial Number:
NPRA(R2/1)241105/1367