



Unvarnished area
48 x 35 mm



B. No. : 1234567
Mfg. date : MM/YYYY
Exp. date : MM/YYYY

50 x 50 x 92 mm
XXXXXXXXXX

50 ml vial

BEMUNAT 100

**Bendamustine Hydrochloride
100 mg Powder for Concentrate
for Solution for Infusion**

Sterile Single- Use Vial

For Intravenous infusion After Dilution

Reconstitution and Dilution Required.
Retain in carton until time of use

CAUTION : CYTOTOXIC AGENT



Each vial contains:
100 mg of Bendamustine Hydrochloride

Usual Dosage: As directed by the Physician.
For directions for use, see Package leaflet.

Unopened Vial: Store below 30°C.
Keep the vial in the outer carton in order to
protect from light.

After opening: The powder should be
reconstituted immediately with water for injection
(WFI) after opening of the vial.
The reconstituted concentrate should be diluted
immediately with 0.9% w/v sodium chloride
solution.

Solution for Infusion: After reconstitution and
dilution, the solution should be used immediately.
If not used immediately, in-use storage times and
conditions prior to use are the responsibility of
the user.

Keep out of sight and reach of children
Jauhkan daripada kanak - kanak
Controlled Medicine/Ubat Terkawal

Cytotoxic Agent – Use caution during handling
and preparation. Use of gloves and safety
glasses is recommended to avoid exposure
(see insert for details).

50 ml vial

BEMUNAT 100

**Bendamustine Hydrochloride
100 mg Powder for Concentrate
for Solution for Infusion**

Sterile Single- Use Vial

For Intravenous infusion After Dilution

Reconstitution and Dilution Required.
Retain in carton until time of use

CAUTION : CYTOTOXIC AGENT



M.L.: 164/MN/AP/95/F/R

Reg NO.: MAL XXXXXX



Manufactured By:
Natco Pharma Limited,
Pharma Division, Unit V,
Kothur, Rangareddy District,
Telangana 509228
India.



Product Registration Holder:
Perniagaan Normahas Sdn Bhd
1-5-10, Block G, Diamond Square,
Jalan 1/50, melalui Jalan Gombak,
53000 Kuala Lumpur.

50 ml vial

BEMUNAT 100

**Bendamustine Hydrochloride
100 mg Powder for Concentrate
for Solution for Infusion)**

For Intravenous Infusion Only

For Intravenous infusion After Dilution.

Reconstitution and Dilution Required.
Retain in carton until time of use

A/s: 50 x 50 x 92 mm

Bendamustine 100 mg Malaysia 20-11-2024

Pantone 265 C

072C, 032C, 1235C, Black