

UNVARNISHED AREA
40 x 30 mm

2D Barcode area

B. No. : 1234567
Mfg. date : MM/YYYY
Exp. date : MM/YYYY

42 x 42 x 72 mm
4030197

1 vial

BEMUNAT 25

Bendamustine Hydrochloride 25 mg Powder for Concentrate for Solution for Infusion

Sterile Single-Use Vial

For Intravenous infusion After Dilution

Reconstitution and Dilution Required.
Retain in carton until time of use

CAUTION : CYTOTOXIC AGENT

NATCO

Each vial contains:
25 mg of Bendamustine Hydrochloride

Usual Dosage: As directed by the Physician.
For directions for use, see Package leaflet.

Unopened Vial: Store below 30°C.
Keep the vial in the outer carton in order to protect from light.

After opening: The powder should be reconstituted immediately with water for injection (WFI) after opening of the vial.
The reconstituted concentrate should be diluted immediately with 0.9% w/v sodium chloride solution.
Solution for Infusion: After reconstitution and dilution, the solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

Keep out of sight and reach of children
Jauhkan daripada kanak - kanak

Controlled Medicine/Ubat Terkawal

Cytotoxic Agent – Use caution during handling and preparation. Use of gloves and safety glasses is recommended to avoid exposure (see insert for details).

1 vial

BEMUNAT 25

Bendamustine Hydrochloride 25 mg Powder for Concentrate for Solution for Infusion

Sterile Single-Use Vial

For Intravenous infusion After Dilution

Reconstitution and Dilution Required.
Retain in carton until time of use

CAUTION : CYTOTOXIC AGENT

NATCO

M.L.: 164/MN/AP/95/F/R

Reg NO.: MAL25046020AZ



Manufactured By:
Natco Pharma Limited,
Pharma Division, Unit V,
Kothur, Rangareddy District,
Telangana 509228
India.

NATCO

Product Registration Holder:
Perniagaan Normahas Sdn Bhd
1-5-10, Block G, Diamond Square,
Jalan 1/50, melalui Jalan Gombak,
53000 Kuala Lumpur.

BEMUNAT 25

Bendamustine Hydrochloride 25 mg Powder for Concentrate for Solution for Infusion

For Intravenous Infusion Only

For Intravenous infusion After Dilution.

Reconstitution and Dilution Required.
Retain in carton until time of use

A/s: 42 x 42 x 72 mm
Bendamustine 25 mg 06/10/2025

Pantone 032 C Reflex Blue, 7691C, 265 C black