

Diabulyn - 30/70

(Biphasic) Insulin Injection, Biphasic Isophane 100IU/mL

What is in This Leaflet:

- 1) What Diabulyn - 30/70 is used for
- 2) How Diabulyn - 30/70 works
- 3) Before you use Diabulyn - 30/70
- 4) How to use Diabulyn - 30/70
- 5) While you are using it
- 6) Side effects
- 7) Storage and disposal of Diabulyn - 30/70
- 8) Product description
- 9) Product Registration Holder and Manufacturer
- 10) Date of revision
- 11) Serial number

1) What Diabulyn - 30/70 is used for

Diabulyn - 30/70 contains the active substance human insulin, which is used to treat diabetes. Diabetes mellitus is condition in which a person has high blood sugar, either because the pancreas does not produce enough insulin or the cells do not respond to the insulin that is produced; therefore, extra insulin is needed to lower your blood sugar levels. Diabulyn - 30/70 is a premixed suspension of an intermediate-acting and fast-acting insulin.

2) How Diabulyn - 30/70 works

After administration of Insugen - 30/70, it will start to lower your blood sugar in about half an hour, and the effect will last for approximately 24 hours.

3) Before you use Diabulyn - 30/70

When you must not use it

- If you are allergic (hypersensitive) to any of the ingredients in Diabulyn - 30/70 (see section 9 for list of ingredients). Watch out for the signs of allergy in section 6 (*Possible side effects*) or ask your doctor for clarification.
- If you experience symptoms of low blood sugar levels (hypoglycaemia), or a hypoglycaemic attack (symptom of low blood sugar). See section 6 (*What to do in an emergency*) to know more about hypoglycaemia.

Before you start to use it

Inform your doctor if:

- You have any other medical conditions, such as infection, fever, kidney or liver

problems. Some medical conditions can alter your insulin requirements and your doctor will suggest a change in dose.

- You are pregnant, planning a pregnancy or breast-feeding. The amount of insulin you need usually falls during the first three months of pregnancy and increases for the remaining six months. If you are breast-feeding, you may need to alter your insulin intake or diet.

Site of injection

Diabulyn - 30/70 is for injection under the skin (subcutaneously). Never inject your insulin directly into a vein or muscle.

Skin changes at the injection site:

The injection site should be rotated to prevent skin changes such as lumps under the skin. The insulin may not work very well if you inject into a lumpy area. Contact your doctor if you are currently injecting into a lumpy area before you start injecting in a different area. Your doctor may tell you to check your blood sugar more closely, and to adjust your insulin or your other antidiabetic medications dose (see 6, *Possible side effects*).

Taking other medicines

Many medicines affect the way glucose works in your body and they may influence your insulin dose.

Talk to your doctor or pharmacist if you take or have recently taken any other medicines, even those not prescribed.

Listed below are the most common medicines which may affect your insulin treatment.

monoamine oxidase inhibitors (MAOI);
beta-blockers; ACE-inhibitors;
acetylsalicylic acid; anabolic steroids;
sulphonamides; oral contraceptives;
thiazides; glucocorticoids; thyroid hormone
therapy; beta-sympathomimetic; growth
hormones; danazol; octreotide or lanreotide.

Pioglitazone (oral antidiabetic medicine used for the treatment of type 2 Diabetes mellitus)

Some patients with long-standing type 2 diabetes mellitus and heart disease or

previous stroke who were treated with pioglitazone and insulin experienced the development of heart failure.

Inform your doctor as soon as possible. If you experience signs of heart failure such as unusual shortness of breath or rapid increase in weight or localise swelling (oedema).

4) How to use Diabulyn - 30/70

General instructions

Follow your doctor's advice about how to use, when to use and how long to use.

This leaflet is only a general guide. If your doctor has switched you from other type or brand of insulin, or if you are suffering from a medical condition, your dose may have to be adjusted by your doctor.

The best places to give yourself an injection are: the front of your waist (abdomen); your buttocks; the front of your thighs or upper arms. Your insulin will work more quickly if you inject it around the waist.

How to use Cartridge

Things to be checked before using

- Check the label to make sure it is the right type of insulin.
- Always check the cartridge, including the rubber plunger (stopper).
- Don't use it if any damage is seen.
- Take it back to your supplier. See your insulin delivery device manual for further instructions.
- Disinfect the rubber membrane with a medicinal swab.
- Always use a new needle for each injection to prevent contamination.

Do not use

- In insulin infusion pumps.
- If the compatible insulin injector pen is dropped, damaged or crushed, there is a risk of leakage of insulin.
- If it hasn't been stored correctly or been frozen.
- If it's not uniformly white and cloudy when it's re-suspended.

How much to use

Dosage

- Your doctor has told you which

Diabulyn - 30/70

(Biphasic) Insulin Injection, Biphasic Isophane 100IU/mL

insulin to use, how much, when and how often to inject. These instructions are only for you. Follow them exactly and visit your diabetes clinic regularly.

- If you change your insulin type (for example from animal to human), you may have to take more or less than before. This might just be for the first injection or it may be a gradual change over several weeks or months.
- Inject Diabulyn - 30/70 under the skin. You should not administer it using a different administration route. Under no circumstances should Diabulyn - 30/70 be given into a vein.

Instruction for use

- Inject the insulin under the skin.
- Use the injection technique advised by your doctor or diabetes nurse and described in your insulin delivery device manual.
- Keep the needle under your skin for at least 10 seconds to make sure that the full dose has been delivered.
- After each injection be sure to remove and discard the needle and store Diabulyn - 30/70 without the needle attached. Otherwise, the liquid may leak out which can cause inaccurate dosing.
- Do not mix any other insulin in an Diabulyn - 30/70 cartridge. Once the cartridge is empty, do not use it again.

Direction of Use of INSUPen Pro+ with Cartridges.

For use with INSUPen Pro+ (re-usable injector) only. Please refer to User Manual for more information.

When to use it

Use as directed by your doctor or pharmacist.

How long to use it

Continue taking Diabulyn - 30/70 for as long as your doctor recommends.

If you forget to use it

If you take less Diabulyn - 30/70 than you

need or if you forget to take a dose your blood sugar levels may increase. Check your blood sugar.

If you use too much (overdose)

If you take more Diabulyn- 30/70 than your requirement it may cause hypoglycaemia (low blood sugar).

If you stop using it

If you stop taking Diabulyn - 30/70 than you need, your blood sugar levels may become too high.

Do not change your insulin unless your doctor tells you to.

If you have any further questions on the use of this product, ask your diabetes specialist nurse, doctor or pharmacist.

5) While you are using it

Things you must do

- Eat a meal or snack containing carbohydrates within 30 minutes of the injection.
- It is recommended that you measure your blood glucose regularly and have regular check-ups.

Things you must not do

Driving and using machines: your ability to concentrate and react may be reduced if you have hypoglycaemia (low blood sugar). You should contact your doctor about the advisability of driving, if you have:

- Frequent episodes of hypoglycaemia.
- Reduced or absent warning signs of hypoglycaemia.

Things you must be careful of

- **If you have a habit of drinking:** watch for signs of hypoglycaemia and never drink alcohol on an empty stomach. The amount of insulin you need may also change if you drink alcohol.
- If you have trouble with your kidneys or liver, or with your adrenal, pituitary or thyroid glands
- If you are exercising more than usual or if you want to change your usual diet.
- **If you are going abroad:** travelling over

time zones may affect your insulin needs and the timing of your injections.

- If you had hypoglycaemia (low blood sugar) after switching from animal insulin to human insulin and reported that the early warning symptoms were less obvious or different.
- If you are ill: carry on taking your insulin
- If you often have hypoglycaemia or have difficulty recognising the symptoms, please discuss this with your doctor.

6) Side effects

The most common side effect is hypoglycaemia.

You get a hypoglycaemic attack if your blood sugar gets too low. This might happen:

- If you take too much insulin.
- If you eat too little or miss a meal.
- If you exercise more than usual.
- If you have an infection or illness (especially diarrhoea or vomiting);
- If you have a change in your need for insulin; or
- If you have trouble with your kidneys or liver which gets worse.

If you take less Diabulyn - 30/70 than your requirement it may cause hyperglycaemia (high blood sugar).

Common side effects

Symptoms resulting from release of adrenaline are common manifestations of mild to moderate hypoglycaemia. They include cold sweats, anxiety, shakiness, hunger, rapid heartbeat, headache, and nervousness. Weight gain is common when taking insulin.

Less common side effects

Allergic reactions and changes at the injection site (lipodystrophy) may occur at the injection site as a consequence of failure to rotate injection sites within an area. If you inject yourself too often at the same site, fatty tissue under the skin at this site may shrink (lipoatrophy) or thicken (lipohypertrophy). Lumps under the skin may also be caused by build-up of a protein called amyloid. Changes at the injection site (Cutaneous amyloidosis); how often this occurs is not known. Changing the site with each injection may help to

Diabulyn - 30/70

(Biphasic) Insulin Injection, Biphasic Isophane 100IU/mL

prevent such skin changes. If you notice your skin pitting or thickening at the injection site, tell your doctor because these reactions can become more severe, or they may change the absorption of your insulin if you inject in such a site.

Signs of allergy

Reactions (redness, swelling, itching) at the injection site may occur (local allergic reactions). These usually disappear after a few weeks of taking your insulin. If they do not disappear, see your doctor.

Seek medical advice immediately

- If signs of allergy spread to other parts of the body, or
- If you suddenly feel unwell and you start sweating; start being sick (vomiting); have difficulty in breathing; have a rapid heartbeat; feel dizzy; feel like fainting.

You may have a very rare serious allergic reaction to Diabulyn - 30/70 or one of its ingredients (called a systemic allergic reaction). See also warning section (*Before you use Diabulyn - 30/70*)

Abnormal accumulation of fluid (oedema):

It may occur upon initiation of insulin therapy. These symptoms are usually of transitory nature.

Vision problems: you might experience transient visual disturbances when you initiate treatment with insulin.

Painful neuropathy (nerve related pain): If your blood glucose levels improve very fast it may cause a burning, tingling or electric pain. This is called acute painful neuropathy and it usually disappears. If it does not disappear, see your doctor.

Diabetic retinopathy (eye background changes): If you have diabetic retinopathy and your blood glucose levels improve very fast, the retinopathy may get worse. Ask your doctor about this. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please inform your health care provider (doctor, nurse, or

pharmacist).

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03- 78835490, or visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

What to do in an emergency

If you get a hypoglycaemic attack
Hypoglycaemia means your blood sugar level is too low.

Causes of hypoglycaemic attack

- If you take too much insulin.
- If you eat too little or miss a meal.
- If you exercise more than usual.
- If you have an infection or illness (especially diarrhoea or vomiting);
- If you have a change in your need for insulin; or
- If you have trouble with your kidneys or liver which gets worse.

The warning signs of a hypoglycaemic attack may come on suddenly and can include: cold sweat; cool pale skin; headache; rapid heartbeat; feeling sick; feeling very hungry; temporary changes in vision; drowsiness; unusual tiredness and weakness; nervousness or tremor; feeling anxious; feeling confused; difficulty in concentrating.

If you get any of these signs:

- Eat glucose tablets or a high sugar snack (sweets, biscuits, fruit juice), then rest.
- Don't take any insulin if you feel a hypoglycaemic attack coming on.
- Carry glucose tablets, sweets, biscuits or fruit juice with you, just in case.
- Tell your relatives, friends and close colleagues that if you pass out (become unconscious), they must: turn you on your side and seek medical advice straight away. They must not give you any food or drink as it could choke you.
- If severe hypoglycaemia is not treated, it can cause brain damage (temporary or permanent) and even death.
- If you have a hypoglycaemic attack that

makes you pass out, or a lot of hypoglycaemic attack, talk to your doctor. The amount or timing of insulin, food or exercise may need to be adjusted.

Using glucagon

You may recover more quickly from unconsciousness with an injection of the hormone glucagon by someone who knows how to use it. If you are given glucagon you will need glucose or a sugary snack as soon as you are conscious. If you do not respond to glucagon treatment, you will have to be treated in a hospital. Seek medical advice after an injection of glucagon; you need to find the reason for your hypoglycaemic attack to avoid getting more.

If you get a hyperglycaemia

If your blood sugar gets too high. Your blood sugar may get too high (this is called hyperglycaemia).

Causes of hyperglycaemia

- Having forgotten to take your insulin.
- Repeatedly taking less insulin than you need.
- An infection or a fever.
- Eating more than usual.
- Less exercise than usual.

The warning signs appear gradually. They include: increased urination; feeling thirsty; losing your appetite; feeling sick (nausea or vomiting); feeling drowsy or tired; flushed, dry skin; dry mouth and a fruity (acetone) smell of the breath.

If you get any of these signs:

Test your blood sugar level and test your urine for ketones if you can. Then seek medical advice straight away.

These may be signs of a very serious condition called diabetic ketoacidosis. If you don't treat it, this could lead to diabetic coma and death.

7) Storage and Disposal of Diabulyn - 30/70

How to store

- Do not store Diabulyn - 30/70 cartridge in or too near the freezer or cooling element.

Diabulyn - 30/70

(Biphasic) Insulin Injection, Biphasic Isophane 100IU/mL

- b. Keep the Diabulyn - 30/70 cartridge in the outer carton in order to protect from light.
- c. Protect from excessive heat and direct sunlight
- d. Store in a refrigerator at temperature between 2°C and 8°C.
- e. Do not refrigerate. Do not freeze. Diabulyn - 30/70 cartridge which are in use can be kept at room temperature (up to 30°C) for up to 42 days.
- f. Diabulyn - 30/70 which have been frozen must not be used.
- g. Keep Diabulyn - 30/70 out of reach of children. *Jauhi daripada kanak-kanak.*

How to Dispose

Any unused product or waste material should be disposed of in accordance with local requirements. Diabulyn - 30/70 formulations should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

8) Product Description

What it looks like

The suspension for injection comes as cloudy, white, aqueous suspension.

Diabulyn - 30/70 Cartridge

It is supplied in 3 mL colourless glass cartridges (Type 1) sealed using lined seals, plugged with plunger stoppers and glass bead.

Ingredients

Active Ingredients

1 mL contains 100 IU of insulin human.

Inactive Ingredients

The other ingredients are Metacresol, Phenol, Glycerol, Protamine sulphate, Zinc oxide, Hydrochloric acid, Sodium hydroxide, Disodium hydrogen phosphate dodecahydrate and Water for Injections.

MAL Number

MAL24126043AZ

9) Product Registration holder

Biocon Sdn. Bhd.

No.1, Jalan Bioteknologi 1,
Kawasan Perindustrian SiLC,
79200 Iskandar Puteri, Johor, Malaysia.

Manufacturer and Batch Releaser:

Bioton S.A.

Macierzysz, ul. Poznańska 12
05-850 Ożarów Mazowiecki, Poland.

Importer and Distributor:

Duopharma Marketing Sdn. Bhd.

Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section
13, Bangi Industrial Estate, 43650 Bandar
Baru Bangi, Selangor Darul Ehsan,
Malaysia.

10) Date of Revision

March 2026

11) Serial Number

NPRA (R3)24/70