




4 x 5 x 2 ml

Cipla

budecort
Respirator
Suspension **0.5 mg**

Budesonide Respirator Suspension

budecort
respules 0.5mg / 2mL
Respirator Suspension **0.5 mg**

for inhalation only



Store Upright ↑

Directions for use:

The product should be used with a respirator or nebuliser only, under the direction of a physician. It is not to be injected or administered orally. Remove a pouch of combipack respule from the carton. Use a fresh respule for each dose.



Tear one respule from the combipack



To open the respule twist off the cap as shown in the figure.



Squeeze the content into the reservoir of the nebulizing chamber.



Discard the respule after use.

Store the respules in foil pouch at all times. Once the foil pouch is opened the respules should be used within three month.

Date: 19-12-19



Pantone 218 C



Pantone 425 C



Pantone 185 C



Black

Actual Size: 85 x 55 x 140 mm

Path: PC: F:\Vaibhav\Vinod\Malaysia\xxxxxxx Budecort 0.5mg 4x5x2ml Carton Malaysia

4 x 5 x 2 ml

Cipla

budecort
Respirator
Suspension **0.5 mg**

Budesonide Respirator Suspension

budecort
respules 0.5mg / 2mL
Respirator Suspension **0.5 mg**

for inhalation only



Store Upright ↑

Each 2 ml respule contains

Budesonide BP 0.5 mg

- Store below 30°C
- Protect from light
- Do not freeze.

SHAKE WELL BEFORE USE

Keep medicine out of reach of children/
Jauhi ubat dari kanak kanak
CONTROLLED MEDICINE /
UBAT TERKAWAL

Caution:

Even invisible damage to respule caused during storage or transit may result in contamination. Do not use, and return for replacement if the suspension is discoloured, leak found on squeezing or contains visible black particles.

MAL No.: MAL10100018AZ

M.L. 536

Barcode

Registration Holder in Malaysia
CIPLA MALAYSIA SDN BHD Suite 1101,
Amcorp Tower, Amcorp Trade Centre,
18 Persiaran Barat, 46 050 Petaling
Jaya, Selangor, Malaysia.

Mfd. by **CIPLA LTD.**
L-139 to L-146 Verna Industrial Estate,
Verna Goa 403 722 INDIA