



DOCETAX
20

Contains : 1 vial of 0.5ml Docetaxel Injection Concentrate +1 vial of 1.5ml Solvent

Docetaxel Injection
Concentrate **20mg**

DOCETAX
20

Cytotoxic Agent
Cipla

XXX XX

Each single dose vial contains
Docetaxel Trihydrate equivalent to
Anhydrous Docetaxel 20 mg
Polysorbate 80 BP 0.5 ml
Dosage: As directed by the physician.
Each vial contains
Alcohol BP (95%v/v) 13% w/v
(Absolute Alcohol
content..... 15.25% v/v)
Water for
injection BP q.s. 1.5 ml
Caution:
1) This is a potent Cytotoxic agent.
Use only after reading the directions
on the pack insert.
2) It is dangerous to take this
preparation except under medical
supervision.
Keep Medicine out of reach
of children.
Jauhi ubat dari kanak-kanak
Controlled Medicine/Ubat Terkawal
Reconstitution: Reconstitute using
Solvent for Docetaxel Injection
Concentrate 20 mg. Rotate the vial
gently and allow the premix to stand
for 5 minutes. Use the Docetaxel
premix for further dilution.
Storage: Docetaxel Injection
Concentrate: Store under refrigeration
between 2-8°C. Protect from light.
For Diluted Solution For Premix:
Store either between 2-8°C or at room
temperature (between 15-25 °C) for
8 hours. For Infusion Solution:
Use within 4 hours at room temperature.

Contains : 1 vial of 0.5ml Docetaxel Injection Concentrate +1 vial of 1.5ml Solvent

Docetaxel Injection
Concentrate **20mg**

DOCETAX
20

Cytotoxic Agent
Cipla

FOR I.V. INFUSION AFTER DILUTION.
ATTENTION: TO BE SOLD ON DRUG
PRESCRIPTION ONLY.
CAREFULLY READ THE INSTRUCTIONS
BEFORE USE. MUST NOT EXCEED THE
RECOMMENDED DOSE.
DO NOT USE FOR CHILDREN.
**Indication, Administration &
Contraindication:**
See enclosed insert before use.
Docetaxel concentrate
MAL No.:
M.L. No.:
Batch No.:
Mfg. Dt.
Exp. Dt.

Solvent for Docetaxel concentrate
MAL No.:
M.L. No.:
Batch No.:
Mfg. Dt.
Exp. Dt.

Registration Holder in Malaysia
Medispec (M) Sdn. Bhd,
55 & 57, Lorong Sempaden 2,
Off Boundary Road, 11400 Ayer Itam,
Penang, Malaysia.
Mfd. by **CIPLA LTD.**
Verna Industrial Estate,
Verna Goa, INDIA.