


Each single dose vial contains
Docetaxel Trihydrate equivalent to
Anhydrous Docetaxel 20 mg
Polycarbate 80 BP 2.5 mg
Dosage: Is directed by the physician.
Reconstitution: Reconstitute using
Solvent for Docetaxel Injection
Concentrate 20 mg. Rotate the vial gently
and allow the premix to stand for 5 minutes.
Use the Docetaxel premix for further dilution.
Storage: Docetaxel Injection Concentrate vials under
refrigeration between 2-8°C. Protect from light.
For Orale Solution For Parental Use after opening
2-8°C or at room temperature between 15-25°C for
6 hours. For Intravenous Solution Use within 4 hours at
room temperature.

Vial of 0.5ml
Docetaxel Injection
Concentrate (20mg)
**DOCETAX
20**
Cytotoxic Agent
Cipla

**FOR I.V. INFUSION AFTER DILUTION,
Indication, Administration & Contraindications:**
See the Brief Patient Leaflet and
M.M.L. No.
Batch No.
Mfg. Dt.
Exp. Dt.
Keep medicines out of reach of children
Jauh dari dari kanak-kanak
Controlled Medicine (obat Terkawal)
Registration Holder in Malaysia
Medipac (M) Sdn. Bhd.
Mfg. by Cipla Ltd.,
Verna Industrial Estate, Verna Goa, INDIA.

XXX XX



Devendra\Docetax\XXX XX Docetax 20 Solvent Lab.ai
Date :- 24-02-12