

CIATUFF 10 (TADALAFIL TABLETS 10MG)

CIATUFF 20 (TADALAFIL TABLETS 20MG)

Tadalafil Tablets (10mg, 20 mg)

What is in this leaflet

1. What CIATUFF is used for
2. How CIATUFF works
3. Before you use CIATUFF
4. How to use CIATUFF
5. While you are using it
6. Side effects
7. Storage and Disposal of CIATUFF
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What CIATUFF is used for

CIATUFF is a treatment for adult men with erectile dysfunction. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. CIATUFF has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.

How CIATUFF works

CIATUFF contains the active ingredient tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors (PDE 5 inhibitors).

Following sexual stimulation CIATUFF works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. CIATUFF will not help you if you do not have erectile dysfunction. It is important to note that CIATUFF for the treatment of erectile dysfunction does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Before you use CIATUFF

- When you must not use it

Do not take CIATUFF if you:

- are allergic to tadalafil or any of the other ingredients of this medicine
- are taking any form of organic nitrate or nitric oxide donors such as amyl

nitrite. This is a group of medicines ("nitrates") used in the treatment of angina pectoris ("chest pain"). CIATUFF has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor.

- have serious heart disease or recently had a heart attack within the last 90 days.
- recently had a stroke within the last 6 months.
- have low blood pressure or uncontrolled high blood pressure.
- ever had a loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as "stroke of the eye".
- are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as CIATUFF, have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

- Before you start to use it

Talk to your doctor before taking CIATUFF.

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Before taking the tablets, tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells).
- multiple myeloma (cancer of the bone marrow).
- leukaemia (cancer of the blood cells).
- any deformation of your penis.
- a serious liver problem.
- a severe kidney problem.

CIATUFF is not intended for use by women.

Children and adolescents: CIATUFF is not intended for use by children and adolescents under the age of 18.

- Taking other medicines

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do not take CIATUFF if you are already taking nitrates.

Some medicines may be affected by CIATUFF or they may affect how well CIATUFF will work. Tell your doctor or pharmacist if you are already taking:

- an alpha blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia [benign enlargement of the prostate]).
- other medicines to treat high blood pressure.
- riociguat.
- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia)
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection.
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines).
- rifampicin, erythromycin, clarithromycin or itraconazole.
- other treatments for erectile dysfunction.

How to use CIATUFF

- How much to use

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended starting dose is 10 or 20 mg taken before sexual activity. Your doctor will decide the suitable dose for you.

CIATUFF 10 (TADALAFIL TABLETS 10MG)

CIATUFF 20 (TADALAFIL TABLETS 20MG)

Tadalafil Tablets (10mg, 20 mg)

- When to use it

CIATUFF tablets are for oral use in men only. Swallow the tablet whole with some water. The tablets can be taken with or without food.

You may take a CIATUFF tablet at least 30 minutes before sexual activity. CIATUFF may still be effective up to 36 hours after taking the tablet.

- How long to use it

Do not take CIATUFF more than once a day.

- If you forget to use it

CIATUFF is intended for use prior to anticipated sexual activity and is not recommended for continuous daily use.

- If you use too much (overdose)

Contact your doctor. You may experience side effects

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

While you are using it

- Things you must do

If you experience sudden decrease or loss of vision, stop taking CIATUFF and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking CIATUFF and contact your doctor immediately.

- Things you must not do

This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.

- Things to be careful of

Driving and using machines

Some men taking CIATUFF have

reported dizziness. Check carefully how you react to the tablets before driving or using machines.

CIATUFF contains lactose

If you have an intolerance to some sugars, contact your doctor before taking this medicine.

CIATUFF with drinks and alcohol

Drinking alcohol may affect your ability to get an erection and may temporarily lower your blood pressure. If you have taken or are planning to take CIATUFF, avoid excessive drinking (blood alcohol level of 0.08% or greater), since this may increase the risk of dizziness when standing up.

Grapefruit juice may affect how well CIATUFF will work and should be taken with caution. Talk to your doctor for further information.

Fertility

A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects stop using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon).
- chest pain – do not use nitrates but seek immediate medical assistance (frequency uncommon).
- priapism, a prolonged and possibly painful erection after taking CIATUFF (frequency rare). If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare).

Other side effects have been

reported: **Common**

- headache, back pain, muscle aches, pain in arms and legs, facial flushing, nasal congestion, indigestion

Uncommon

- dizziness, stomach ache, feeling sick, being sick (vomiting), reflux, blurred vision, eye pain, difficulty in breathing, presence of blood in urine, prolonged erection, pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure, nose bleeds, ringing in the ears, swelling of the hands, feet or ankles and feeling tired.

Rare

- fainting, seizures and passing memory loss, swelling of the eyelids, red eyes, sudden decrease or loss of hearing, hives (itchy red welts on the surface of the skin), penile bleeding, presence of blood in semen and increased sweating.

Heart attack and stroke have also been reported rarely in men taking CIATUFF. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking CIATUFF that were not seen in clinical trials.

These include:

- migraine, swelling of the face, serious allergic reaction which causes swelling of the face or throat, serious skin rashes, some disorders affecting blood flow to the eyes, irregular heartbeats, angina and sudden cardiac death.

The side effect dizziness has been reported more frequently in men over 75 years of age taking CIATUFF.

Diarrhoea has been reported more frequently in men over 65 years of age taking CIATUFF.

If you get any side effects, talk to your doctor or pharmacist. This includes possible side effects not listed in this leaflet.

CIATUFF 10 (TADALAFIL TABLETS 10MG)

CIATUFF 20 (TADALAFIL TABLETS 20MG)

Tadalafil Tablets (10mg, 20 mg)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and disposal of CIATUFF

- Storage

Keep out of the reach and sight of children.

Do not use this medicine after the expiry date, which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture. Store below

30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines which are no longer required. These measures will help to protect the environment.

Product Description

What it looks like

CIATUFF 10 mg:

Light yellow colored, oval shaped, film-coated tablets debossed with '10' on one side and 'TL' on other side.

CIATUFF 20 mg:

Yellow colored, oval shaped, film-coated tablets debossed with '20' on one side and 'TL' on other side.

Presentation: Box of 1 blister x 4 tablets
OR 2 blister x 4 tablets.

Ingredients

Active ingredient:

CIATUFF 10 mg

Each film coated tablet contains 10 mg of tadalafil.

CIATUFF 20 mg

Each film coated tablet contains 20 mg

of tadalafil.

Inactive ingredients:

Copovidone, Polyoxyl 40 hydrogenated castor oil, Lactose Monohydrate, Silica Colloidal Anhydrous, Microcrystalline cellulose, Croscarmellose Sodium, Magnesium Stearate, Opadry Yellow

MAL number: MAL19096020AZ,
MAL19096019AZ

Manufacturer

Aurobindo Pharma Limited, Unit-VII (SEZ), TSIIC, Plot No. S1, Survey No. 411/P, 425/P, 434/P, 435/P & 458/P, Green Industrial Park, Polepally Village, Jedcherla Mandal, Mahaboobnagar District, Telangana State, India

Product Registration Holder

Synerrv Sdn Bhd, SO-29-2, Menara 1, KL ECO City, Jalan Bangsar, KG Haji Abdullah Hukum, 59200 Kuala Lumpur Wilayah Persekutuan Kuala Lumpur, Malaysia.

Date of revision

03/11/2021

Serial Number

NPRA (R2) 19/99