

Sofovir

Sofosbuvir Tablets 400 mg

Sofovir (Sofosbuvir Tablets 400 mg)

Each film-coated tablet contains Sofosbuvir 400 mg.

PRODUCT DESCRIPTION

Orange oval, bevel biconvex, film coated tablets debossed with 'H' on one side and 'S14' on the other side.

PHARMACODYNAMICS

Pharmacotherapeutic group: Direct-acting antiviral; ATC code: J05AP08

Mechanism of action

Sofosbuvir is a pan-genotypic inhibitor of the HCV NS5B RNA-dependent RNA polymerase, which is essential for viral replication. Sofosbuvir is a nucleotide prodrug that undergoes intracellular metabolism to form the pharmacologically active uridine analog triphosphate (GS-461203), which can be incorporated into HCV RNA by the NS5B polymerase and acts as a chain terminator. In a biochemical assay, GS-461203 inhibited the polymerase activity of the recombinant NS5B from HCV genotype 1b, 2a, 3a and 4a with a 50% inhibitory concentration (IC₅₀) value ranging from 0.7 to 2.6 µM. GS-461203 (the active metabolite of sofosbuvir) is not an inhibitor of human DNA and RNA polymerases nor an inhibitor of mitochondrial RNA polymerase.

Antiviral activity

In HCV replicon assays, the effective concentration (EC₅₀) values of sofosbuvir against full-length replicons from genotype 1a, 1b, 2a, 3a and 4a were 0.04, 0.11, 0.05, 0.05 and 0.04 µM, respectively, and EC₅₀ values of sofosbuvir against chimeric 1b replicons encoding NS5B from genotype 2b, 5a or 6a were 0.014 to 0.015 µM. The mean ± SD EC₅₀ of sofosbuvir against chimeric replicons encoding NS5B sequences from clinical isolates was 0.068 ± 0.024 µM for genotype 1a (n = 67), 0.11 ± 0.029 µM for genotype 1b (n = 29), 0.035 ± 0.018 µM for genotype 2 (n = 15) and 0.085 ± 0.034 µM for genotype 3a (n = 106). In these assays, the *in vitro* antiviral activity of sofosbuvir against the less common genotypes 4, 5 and 6 was similar to that observed for genotypes 1, 2 and 3.

The presence of 40% human serum had no effect on the anti-HCV activity of sofosbuvir.

Resistance

In cell culture

HCV replicons with reduced susceptibility to sofosbuvir have been selected in cell culture for multiple genotypes including 1b, 2a, 2b, 3a, 4a, 5a and 6a. Reduced susceptibility to sofosbuvir was associated with the primary NS5B substitution S282T in all replicon genotypes examined. Site-directed mutagenesis of the S282T substitution in replicons of 8 genotypes conferred 2- to 18-fold reduced susceptibility to sofosbuvir and reduced the replication viral capacity by 89% to 99% compared to the corresponding wild-type. In biochemical assays, recombinant NS5B polymerase from genotypes 1b, 2a, 3a and 4a expressing the S282T substitution showed reduced susceptibility to GS-461203 compared to respective wild-types.

In clinical studies

In a pooled analysis of 991 subjects who received sofosbuvir in Phase 3 studies, 226 subjects qualified for resistance analysis due to virologic failure or early study drug discontinuation and having HCV RNA >1,000 IU/mL. Post-baseline NS5B sequences were available for 225 of the 226 subjects, with deep sequencing data (assay cutoff of 1%) from 221 of these subjects. The sofosbuvir-associated resistance substitution S282T was not detected in any of these subjects by deep sequencing or

population sequencing. The S282T substitution in NS5B was detected in a single subject receiving Sofosbuvir monotherapy in a Phase 2 study. This subject harboured <1% HCV S282T at baseline and developed S282T (>99%) at 4 weeks post-treatment which resulted in a 13.5-fold change in sofosbuvir EC50 and reduced viral replication capacity. The S282T substitution reverted to wild-type over the next 8 weeks and was no longer detectable by deep sequencing at 12 weeks post-treatment.

Two NS5B substitutions, L159F and V321A, were detected in post-treatment relapse samples from multiple genotype 3 HCV infected subjects in the Phase 3 clinical studies. No shift in the phenotypic susceptibility to sofosbuvir or ribavirin of subject isolates with these substitutions was detected. In the study done in subjects with hepatocellular carcinoma awaiting liver transplantation where subjects received up to 48 weeks of sofosbuvir and ribavirin, the L159F substitution emerged in multiple subjects with GT1a or GT2b HCV who experienced virologic failure (breakthrough and relapse). Furthermore, the presence of substitutions L159F and/or C316N at baseline was associated with sofosbuvir breakthrough and relapse post-transplant in multiple subjects infected with GT1b HCV. In addition, S282R and L320F substitutions were detected on treatment by deep sequencing in a pre-transplant subject with GT1a HCV with a partial treatment response. The clinical significance of these findings is unknown.

Effect of baseline HCV polymorphisms on treatment outcome

Baseline NS5B sequences were obtained for 1,292 subjects from Phase 3 studies by population sequencing and the S282T substitution was not detected in any subject with available baseline sequence. In an analysis evaluating the effect of baseline polymorphisms on treatment outcome, no statistically significant association was observed between the presence of any HCV NS5B variant at baseline and treatment outcome.

Cross-resistance

HCV replicons expressing the sofosbuvir-associated resistance substitution S282T were fully susceptible to other classes of anti-HCV agents. Sofosbuvir retained activity against the NS5B substitutions L159F and L320F associated with resistance to other nucleoside inhibitors. Sofosbuvir was fully active against substitutions associated with resistance to other direct-acting antivirals with different mechanisms of actions, such as NS5B non-nucleoside inhibitors, NS3 protease inhibitors and NS5A inhibitors.

Clinical efficacy and safety

The efficacy of sofosbuvir was evaluated in five Phase 3 studies in a total of 1,568 subjects with genotypes 1 to 6 chronic hepatitis C. One study was conducted in treatment-naïve subjects with genotype 1, 4, 5 or 6 chronic hepatitis C in combination with peginterferon alfa 2a and ribavirin and the other four studies were conducted in subjects with genotype 2 or 3 chronic hepatitis C in combination with ribavirin including one in treatment-naïve subjects, one in interferon intolerant, ineligible or unwilling subjects, one in subjects previously treated with an interferon-based regimen, and one in all subjects irrespective of prior treatment history or ability to receive treatment with interferon. Subjects in these studies had compensated liver disease including cirrhosis. Sofosbuvir was administered at a dose of 400 mg once daily. The ribavirin dose was weight-based at 1,000-1,200 mg daily administered in two divided doses, and the peginterferon alfa 2a dose, where applicable, was 180 µg per week. Treatment duration was fixed in each study and was not guided by subjects' HCV RNA levels (no response guided algorithm).

Plasma HCV RNA values were measured during the clinical studies using the COBAS TaqMan HCV test (version 2.0), for use with the High Pure System. The assay had a lower limit of quantification (LLOQ) of 25 IU/mL. Sustained virologic response (SVR) was the primary endpoint to determine the HCV cure rate for all studies which was defined as HCV RNA less than LLOQ at 12 weeks after the end of treatment (SVR12).

Clinical studies in subjects with genotype 1 and 4 chronic hepatitis C

Treatment-naïve subjects - NEUTRINO (study 110)

NEUTRINO was an open-label, single-arm study that evaluated 12 weeks of treatment with sofosbuvir in combination with peginterferon alfa 2a and ribavirin in treatment-naïve subjects with genotype 1, 4, 5 or 6 HCV infection.

Treated subjects (n = 327) had a median age of 54 years (range: 19 to 70); 64% of the subjects were male; 79% were White; 17% were Black; 14% were Hispanic or Latino; mean body mass index was 29 kg/m² (range: 18 to 56 kg/m²); 78% had baseline HCV RNA greater than 6 log₁₀ IU/mL; 17% had cirrhosis; 89% had HCV genotype 1 and 11% had HCV genotype 4, 5 or 6. Table 1 presents the response rates for the treatment group of sofosbuvir + peginterferon alfa + ribavirin.

Table 1: Response rates in study NEUTRINO

	SOF+PEG+RBV 12 weeks (n = 327)
Overall SVR12	91% (296/327)
Outcome for subjects without SVR12	
On-treatment virologic failure	0/327
Relapse ^a	9% (28/326)
Other ^b	1% (3/327)

a. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

b. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

Response rates for selected subgroups are presented in Table 2.

Table 2: SVR12 rates for selected subgroups in NEUTRINO

	SOF+PEG+RBV 12 weeks (n = 327)
Genotype	
Genotype 1	90% (262/292)
Genotype 1a	92% (206/225)
Genotype 1b	83% (55/66)
Genotype 4	96% (27/28)
Cirrhosis	
No	93% (253/273)
Yes	80% (43/54)
Race	
Black	87% (47/54)
Non-Black	91% (249/273)

SVR12 rates were similarly high in subjects with baseline IL28B C/C allele [94/95 (99%)] and non-C/C (C/T or T/T) allele [202/232 (87%)].

27/28 patients with genotype 4 HCV achieved SVR12. A single subject with genotype 5 and all 6 subjects with genotype 6 HCV infection in this study achieved SVR12.

Clinical studies in subjects with genotype 2 and 3 chronic hepatitis C Treatment-naïve adults - FISSION (study 1231)

FISSION was a randomised, open-label, active-controlled study that evaluated 12 weeks of treatment with sofosbuvir and ribavirin compared to 24 weeks of treatment with peginterferon alfa 2a and ribavirin in treatment-naïve subjects with genotype 2 or 3 HCV infection. The ribavirin doses used in the sofosbuvir + ribavirin and peginterferon alfa 2a + ribavirin arms were weight-based 1,000-1,200 mg/day and 800 mg/day regardless of weight, respectively. Subjects were randomised in a 1:1 ratio and stratified by cirrhosis (presence *versus* absence), HCV genotype (2 *versus* 3) and baseline HCV RNA level (<6 log₁₀ IU/mL *versus* ≥6 log₁₀ IU/mL). Subjects with genotype 2 or 3 HCV were enrolled in an approximately 1:3 ratio.

Treated subjects (n = 499) had a median age of 50 years (range: 19 to 77); 66% of the subjects were male; 87% were White; 3% were Black; 14% were Hispanic or Latino; mean body mass index was 28 kg/m² (range: 17 to 52 kg/m²); 57% had baseline HCV RNA levels greater than 6 log₁₀ IU/mL; 20% had cirrhosis; 72% had HCV genotype 3. Table 3 presents the response rates for the treatment groups of sofosbuvir + ribavirin and peginterferon alfa + ribavirin.

Table 3: Response rates in study FISSION

	SOF+RBV 12 weeks (n = 256)^a	PEG+RBV 24 weeks (n = 243)
Overall SVR12	67% (171/256)	67% (162/243)
Genotype 2	95% (69/73)	78% (52/67)
Genotype 3	56% (102/183)	63% (110/176)
Outcome for subjects without SVR12		
On-treatment virologic failure	< 1% (1/256)	7% (18/243)
Relapse ^b	30% (76/252)	21% (46/217)
Other ^c	3% (8/256)	7% (17/243)

a. The efficacy analysis includes 3 subjects with recombinant genotype 2/1 HCV infection.

b. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

c. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

The difference in the overall SVR12 rates between sofosbuvir + ribavirin and peginterferon alfa + ribavirin treatment groups was 0.3% (95% confidence interval: -7.5% to 8.0%) and the study met the predefined non-inferiority criterion.

Response rates for subjects with cirrhosis at baseline are presented in Table 4 by HCV genotype.

Table 4: SVR12 rates by cirrhosis and genotype in study FISSION

	Genotype 2		Genotype 3	
	SOF+RBV 12 weeks (n = 73)^a	PEG+RBV 24 weeks (n = 67)	SOF+RBV 12 weeks (n = 183)	PEG+RBV 24 weeks (n = 176)
Cirrhosis				

No	97% (59/61)	81% (44/54)	61% (89/145)	71% (99/139)
Yes	83% (10/12)	62% (8/13)	34% (13/38)	30% (11/37)

a. The efficacy analysis includes 3 subjects with recombinant genotype 2/1 HCV infection.

Interferon intolerant, ineligible or unwilling adults - POSITRON (study 107)

POSITRON was a randomised, double-blinded, placebo-controlled study that evaluated 12 weeks of treatment with sofosbuvir and ribavirin (n = 207) compared to placebo (n = 71) in subjects who are interferon intolerant, ineligible or unwilling. Subjects were randomised in 3:1 ratio and stratified by cirrhosis (presence *versus* absence).

Treated subjects (n = 278) had a median age of 54 years (range: 21 to 75); 54% of the subjects were male; 91% were White; 5% were Black; 11% were Hispanic or Latino; mean body mass index was 28 kg/m² (range: 18 to 53 kg/m²); 70% had baseline HCV RNA levels greater than 6 log₁₀ IU/mL; 16% had cirrhosis; 49% had HCV genotype 3. The proportions of subjects who were interferon intolerant, ineligible, or unwilling were 9%, 44%, and 47%, respectively. Most subjects had no prior HCV treatment (81.3%). Table 5 presents the response rates for the treatment groups of sofosbuvir + ribavirin and placebo.

Table 5: Response rates in study POSITRON

	SOF+RBV 12 weeks(n = 207)	Placebo 12 weeks (n = 71)
Overall SVR12	78% (161/207)	0/71
Genotype 2	93% (101/109)	0/34
Genotype 3	61% (60/98)	0/37
Outcome for subjects without SVR12		
On-treatment virologic failure	0/207	97% (69/71)
Relapse ^a	20% (42/205)	0/0
Other ^b	2% (4/207)	3% (2/71)

a. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

b. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

The SVR12 rate in the sofosbuvir + ribavirin treatment group was statistically significant when compared to placebo ($p < 0.001$).

Table 6 presents the subgroup analysis by genotype for cirrhosis and interferon classification.

Table 6: SVR12 rates for selected subgroups by genotype in POSITRON

	SOF+RBV 12 weeks	
	Genotype 2 (n = 109)	Genotype 3 (n = 98)
Cirrhosis		
No	92% (85/92)	68% (57/84)
Yes	94% (16/17)	21% (3/14)
Interferon classification		
Ineligible	88% (36/41)	70% (33/47)
Intolerant	100% (9/9)	50% (4/8)
Unwilling	95% (56/59)	53% (23/43)

Previously treated adults - FUSION (study 108)

FUSION was a randomised, double-blinded study that evaluated 12 or 16 weeks of treatment with sofosbuvir and ribavirin in subjects who did not achieve SVR with prior interferon-based treatment (relapsers and nonresponders). Subjects were randomised in a 1:1 ratio and stratified by cirrhosis (presence *versus* absence) and HCV genotype (2 *versus* 3).

Treated subjects (n = 201) had a median age of 56 years (range: 24 to 70); 70% of the subjects were male; 87% were White; 3% were Black; 9% were Hispanic or Latino; mean body mass index was 29 kg/m² (range: 19 to 44 kg/m²); 73% had baseline HCV RNA levels greater than 6 log₁₀ IU/mL; 34% had cirrhosis; 63% had HCV genotype 3; 75% were prior relapsers. Table 7 presents the response rates for the treatment groups of sofosbuvir + ribavirin for 12 weeks and 16 weeks.

Table 7: Response rates in study FUSION

	SOF+RBV 12 weeks (n = 103)^a	SOF+RBV 16 weeks (n = 98)^a
Overall SVR12	50% (51/103)	71% (70/98)
Genotype 2	82% (32/39)	89% (31/35)
Genotype 3	30% (19/64)	62% (39/63)
Outcome for subjects without SVR12		
On-treatment virologic failure	0/103	0/98
Relapse ^b	48% (49/103)	29% (28/98)
Other ^c	3% (3/103)	0/98

a. The efficacy analysis includes 6 subjects with recombinant genotype 2/1 HCV infection.

b. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

c. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

Table 8 presents the subgroup analysis by genotype for cirrhosis and response to prior HCV treatment.

Table 8: SVR12 rates for selected subgroups by genotype in study FUSION

	Genotype 2		Genotype 3	
	SOF+RBV 12 weeks (n = 39)	SOF+RBV 16 weeks (n = 35)	SOF+RBV 12 weeks (n = 64)	SOF+RBV 16 weeks (n = 63)
Cirrhosis				
No	90% (26/29)	92% (24/26)	37% (14/38)	63% (25/40)
Yes	60% (6/10)	78% (7/9)	19% (5/26)	61% (14/23)
Response to prior HCV treatment				
Relapser	86% (25/29)	89% (24/27)	31% (15/49)	65% (30/46)
Nonresponder	70% (7/10)	88% (7/8)	27% (4/15)	53% (9/17)

Treatment-naïve and previously treated adults - VALENCE (study 133)

VALENCE was a Phase 3 study that evaluated sofosbuvir in combination with weight-based ribavirin for the treatment of genotype 2 or 3 HCV infection in treatment-naïve subjects or subjects who did not achieve SVR with prior interferon-based treatment, including subjects with compensated cirrhosis.

The study was designed as a direct comparison of sofosbuvir and ribavirin *versus* placebo for 12 weeks. However, based on emerging data, the study was unblinded and all HCV genotype 2 subjects continued to receive sofosbuvir and ribavirin for 12 weeks, whilst treatment for HCV genotype 3 subjects was extended to 24 weeks. Eleven HCV genotype 3 subjects had already completed treatment with sofosbuvir and ribavirin for 12 weeks at the time of the amendment.

Treated subjects (n = 419) had a median age of 51 years (range: 19 to 74); 60% of the subjects were male; median body mass index was 25 kg/m² (range: 17 to 44 kg/m²); the mean baseline HCV RNA level was 6.4 log₁₀ IU/mL; 21% had cirrhosis; 78% had HCV genotype 3; 65% were prior relapsers. Table 9 presents the response rates for the treatment groups of sofosbuvir + ribavirin for 12 weeks and 24 weeks.

Placebo recipients are not included in the tables since none achieved SVR12.

Table 9: Response rates in study VALENCE

	Genotype 2 SOF+RBV 12 weeks (n = 73)	Genotype 3 SOF+RBV 12 weeks (n = 11)	Genotype 3 SOF+RBV 24 weeks (n = 250)
Overall SVR12	93% (68/73)	27% (3/11)	84% (210/250)
Outcome for subjects without SVR12			
On-treatment virologic failure	0% (0/73)	0% (0/11)	0.4% (1/250)
Relapse ^a	7% (5/73)	55% (6/11)	14% (34/249)
Other ^b	0% (0/73)	18% (2/11)	2% (5/250)

a. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

b. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

Table 10 presents the subgroup analysis by genotype for cirrhosis and exposure to prior HCV treatment.

Table 10: SVR12 rates for selected subgroups by genotype in study VALENCE

	Genotype 2 SOF+RBV 12 weeks (n = 73)	Genotype 3 SOF+RBV 24 weeks (n = 250)
Treatment-naïve	97% (31/32)	93% (98/105)
Non-cirrhotic	97% (29/30)	93% (86/92)
Cirrhotic	100% (2/2)	92% (12/13)
Treatment-experienced	90% (37/41)	77% (112/145)
Non-cirrhotic	91% (30/33)	85% (85/100)
Cirrhotic	88% (7/8)	60% (27/45)

SVR12 to SVR24 concordance

The concordance between SVR12 and SVR24 (SVR 24 weeks after the end of the treatment) following treatment with sofosbuvir in combination with ribavirin or ribavirin and pegylated interferon demonstrates a positive predictive value of 99% and a negative predictive value of 99%.

Clinical efficacy and safety in special populations

HCV/HIV co-infected patients - PHOTON-1 (study 123)

Sofosbuvir was studied in an open-label clinical study evaluating the safety and efficacy of 12 or 24 weeks of treatment with sofosbuvir and ribavirin in subjects with genotype 1, 2 or 3 chronic hepatitis C co-infected with HIV-1. Genotype 2 and 3 subjects were either treatment-naïve or experienced, whereas genotype 1 subjects were naïve to prior treatment. Treatment duration was 12 weeks in treatment-naïve subjects with genotype 2 or 3 HCV infection, and 24 weeks in treatment-experienced subjects with genotype 3 HCV infection, as well as subjects with genotype 1 HCV infection. Subjects received 400 mg sofosbuvir and weight-based ribavirin (1,000 mg for subjects weighing <75 kg or 1,200 mg for subjects weighing ≥75 kg). Subjects were either not on antiretroviral therapy with a CD4+ cell count >500 cells/mm³ or had virologically suppressed HIV-1 with a CD4+ cell count >200 cells/mm³. 95% of patients received antiretroviral therapy at the time of enrolment. Preliminary SVR12 data are available for 210 subjects.

Table 11 presents the response rates by genotype and exposure to prior HCV treatment.

Table 11: Response rates in study PHOTON-1

	Genotype 2/3 treatment-naïve SOF+RBV 12 weeks (n = 68)	Genotype 2/3 treatment-experienced SOF+RBV 24 weeks (n = 28)	Genotype 1 treatment-naïve SOF+RBV 24 weeks (n = 114)
Overall SVR12	75% (51/68)	93% (26/28)	76% (87/114)
Outcome for subjects without SVR12			
On-treatment virologic failure	1% (1/68)	0/28	1% (1/114)
Relapse ^a	18% (12/67)	7% (2/28)	22% (25/113)
Other ^b	6% (4/68)	0/28	1% (1/114)

a. The denominator for relapse is the number of subjects with HCV RNA <LLOQ at their last on-treatment assessment.

b. Other includes subjects who did not achieve SVR12 and did not meet virologic failure criteria (e.g., lost to follow-up).

Table 12 presents the subgroup analysis by genotype for cirrhosis.

Table 12: SVR12 rates for selected subgroups by genotype in study PHOTON-1

	HCV genotype 2		HCV genotype 3	
	SOF+RBV 12 weeks TN (n = 26)	SOF+RBV 24 weeks TE (n = 15)	SOF+RBV 12 weeks TN (n = 42)	SOF+RBV 24 weeks TE (n = 13)
Overall	88% (23/26)	93% (14/15)	67% (28/42)	92% (12/13)
No cirrhosis	88% (22/25)	92% (12/13)	67% (24/36)	100% (8/8)
Cirrhosis	100% (1/1)	100% (2/2)	67% (4/6)	80% (4/5)

TN = treatment-naïve; TE = treatment-experienced.

In subjects with HCV genotype 1 infection, the SVR rate was 82% (74/90) in subjects with genotype 1a infection and 54% (13/24) in subjects with genotype 1b infection, with relapse accounting for the majority of treatment failures. SVR rates in subjects with HCV genotype 1 infection were 80% (24/30) in subjects with baseline IL28B C/C allele and 75% (62/83) in subjects with baseline IL28B non-C/C alleles.

In the 223 CHC subjects with HIV-1 co-infection, the percentage of CD4+ cells did not change during treatment. Median CD4+ cell count decreases of 85 cells/mm³ and 84 cells/mm³ were observed at the end of treatment with sofosbuvir +ribavirin for 12 or 24 weeks, respectively. HIV-1 rebound during sofosbuvir +ribavirin treatment occurred in 2 subjects (0.9%) on antiretroviral therapy.

Patients awaiting liver transplantation - Study 2025

Sofosbuvir was studied in HCV infected subjects prior to undergoing liver transplantation in an open-label clinical study evaluating the safety and efficacy of sofosbuvir and ribavirin administered pre-transplant to prevent post-transplant HCV reinfection. The primary endpoint of the study was post-transplant virologic response (pTVR, HCV RNA <LLOQ at 12 weeks post-transplant).

HCV infected subjects, regardless of genotype, with hepatocellular carcinoma (HCC) meeting the MILAN criteria received 400 mg sofosbuvir and 1,000-1,200 mg ribavirin daily for a maximum of 24 weeks, subsequently amended to 48 weeks, or until the time of liver transplantation, whichever occurred first. An interim analysis was conducted on 61 subjects who received sofosbuvir and ribavirin; the majority of subjects had HCV genotype 1, 44 subjects were CPT class A and 17 subjects were CPT class B. Of these 61 subjects, 44 subjects underwent liver transplantation following up to 48 weeks of treatment with sofosbuvir and ribavirin; 41 had HCV RNA <LLOQ at the time of transplantation. The virologic response rates of the 41 subjects transplanted with HCV RNA <LLOQ is described in Table 13. Duration of viral suppression prior to transplantation was the most predictive factor for pTVR in those who were HCV RNA <LLOQ at the time of transplantation.

Table 13: Virologic response post-transplant in subjects with HCV RNA <LLOQ at the time of liver transplantation

	Week 12 post-transplant (pTVR) ^b
Virologic response in evaluable subjects ^a	23/37 (62%)

a. Evaluable subjects are defined as those who have reached the specified time point at the time of the interim analysis.

b. pTVR: post-transplant virologic response (HCV RNA <LLOQ at 12 weeks post-procedure).

In patients that discontinued therapy at 24 weeks, according to protocol, the relapse rate was 11/15.

Liver transplant recipients - Study 0126

Sofosbuvir was studied in an open-label clinical study evaluating the safety and efficacy of 24 weeks of treatment with sofosbuvir and ribavirin in liver transplant recipients with chronic hepatitis C. Eligible

subjects were ≥ 18 years old and had undergone liver transplantation 6 to 150 months prior to screening. Subjects had HCV RNA $\geq 10^4$ IU/mL at screening and documented evidence of chronic HCV infection pre-transplantation. The starting dose of ribavirin was 400 mg given in a divided daily dose. If subjects maintained haemoglobin levels ≥ 12 g/dL, ribavirin dose was increased at weeks 2, 4, and up to every 4 weeks until the appropriate weight-based dose was reached (1,000 mg daily in subjects < 75 kg, 1,200 mg daily in subjects ≥ 75 kg). The median ribavirin dose was 600 mg-800 mg daily at weeks 4-24.

Forty subjects (33 with HCV genotype 1 infection, 6 with HCV genotype 3 infection, and 1 with HCV genotype 4 infection) were enrolled, 35 of whom had previously failed interferon-based treatment, and 16 of whom had cirrhosis. 28 out of 40 (70%) subjects achieved SVR12: 22/33 (73%) with HCV genotype 1 infection, 6/6 (100%) with HCV genotype 3 infection, and 0/1 (0%) with HCV genotype 4 infection. All subjects who achieved SVR12 achieved SVR24 and SVR48.

Overview of outcomes by therapeutic regimen and treatment duration, a comparison across studies The following tables (Table 14 to Table 17) present data from Phase 2 and Phase 3 studies relevant to the dosing to help clinicians determine the best regimen for individual patients.

Table 14: Outcomes by therapeutic regimen and treatment duration, a comparison across studies in genotype 1 HCV infection

Patient population (Study number/name)	Regimen/Duration	Subgroup	SVR12 rate % (n/N)
Treatment-naïve ^a (NEUTRINO)	SOF+PEG+RBV 12 weeks	Overall	90% (262/292)
		Genotype 1a	92% (206/225)
		Genotype 1b	83% (55/66)
		No cirrhosis	93% (253/273)
		Cirrhosis	80% (43/54)
Treatment-naïve and co-infected with HIV (PHOTON-1)	SOF+RBV 24 weeks	Overall	76% (87/114)
		Genotype 1a	82% (74/90)
		Genotype 1b	54% (13/24)
		No cirrhosis	77% (84/109)
		Cirrhosis	60% (3/5)
Treatment-naïve (QUANTUM ^b and 11-1-0258 ^b)	SOF+RBV 24 weeks	Overall ^c	65% (104/159)
		Genotype 1a ^c	69% (84/121)
		Genotype 1b ^c	53% (20/38)
		No cirrhosis ^c	68% (100/148)
		Cirrhosis ^c	36% (4/11)

n = number of subjects with SVR12 response; N = total number of subjects per group.

a. For previously treated patients with genotype 1 HCV infection, no data exists with the combination of sofosbuvir, peginterferon alfa and ribavirin. Consideration should be given to treating these patients, and potentially extending the duration of therapy with sofosbuvir, peginterferon alfa and ribavirin beyond 12 weeks and up to 24 weeks; especially for those subgroups who have one or more factors historically associated with lower response rates to interferon-based therapies (prior null response to peginterferon alfa and ribavirin therapy, advanced fibrosis/cirrhosis, high baseline viral concentrations, black race, IL28B non CC genotype).

b. These are exploratory or Phase 2 studies. The outcomes should be interpreted with caution, as subject numbers are small and SVR rates may be impacted by the selection of patients.

c. Summary data from both studies.

Table 15: Outcomes by therapeutic regimen and treatment duration, a comparison across studies in genotype 2 HCV infection

Patient population (Study number/name)	Regimen/Duration	Subgroup	SVR12 rate % (n/N)
Treatment-naïve (FISSION)	SOF+RBV 12 weeks	Overall	95% (69/73)
		No cirrhosis	97% (59/61)
		Cirrhosis	83% (10/12)
Interferon intolerant, ineligible or unwilling (POSITRON)	SOF+RBV 12 weeks	Overall	93% (101/109)
		No cirrhosis	92% (85/92)
		Cirrhosis	94% (16/17)
Treatment-experienced (FUSION)	SOF+RBV 12 weeks	Overall	82% (32/39)
		No cirrhosis	90% (26/29)
		Cirrhosis	60% (6/10)
Treatment-naïve (VALENCE)	SOF+RBV 12 weeks	Overall	97% (31/32)
		No cirrhosis	97% (29/30)
		Cirrhosis	100% (2/2)
Treatment-experienced (VALENCE)	SOF+RBV 12 weeks	Overall	90% (37/41)
		No cirrhosis	91% (30/33)
		Cirrhosis	88% (7/8)
Treatment-experienced (FUSION)	SOF+RBV 16 weeks	Overall	89% (31/35)
		No cirrhosis	92% (24/26)
		Cirrhosis	78% (7/9)
Treatment-naïve co-infected with HIV (PHOTON-1)	SOF+RBV 12 weeks	Overall	88% (23/26)
		No cirrhosis	88% (22/25)
		Cirrhosis	100% (1/1)
Treatment-experienced co-infected with HIV (PHOTON-1)	SOF+RBV 24 weeks	Overall ^a	93% (14/15)
		No cirrhosis ^a	92% (12/13)
		Cirrhosis ^a	100% (2/2)
Treatment-naïve (ELECTRON ^b)	SOF+PEG+RBV 12 weeks	Overall ^c	96% (25/26)
Treatment-experienced (LONESTAR-2 ^b)	SOF+PEG+RBV 12 weeks	Overall	96% (22/23)
		No cirrhosis	100% (9/9)
		Cirrhosis	93% (13/14)

n = number of subjects with SVR12 response; N = total number of subjects per group.

a. These data are preliminary.

b. These are exploratory or Phase 2 studies. The outcomes should be interpreted with caution, as subject numbers are small and SVR rates may be impacted by the selection of patients. In the ELECTRON study (N = 11), the duration of peginterferon alfa ranged from 4-12 weeks in combination with sofosbuvir + ribavirin.

c. All patients were non-cirrhotic in these two studies.

Table 16: Outcomes by therapeutic regimen and treatment duration, a comparison across studies in genotype 3 HCV infection

Patient population (Study number/name)	Regimen/Duration	Subgroup	SVR12 rate % (n/N)
Treatment-naïve (FISSION)	SOF+RBV 12 weeks	Overall	56% (102/183)
		No cirrhosis	61% (89/145)
		Cirrhosis	34% (13/38)
Interferon intolerant, ineligible or unwilling (POSITRON)	SOF+RBV 12 weeks	Overall	61% (60/98)
		No cirrhosis	68% (57/84)
		Cirrhosis	21% (3/14)
Treatment-experienced (FUSION)	SOF+RBV 12 weeks	Overall	30% (19/64)
		No cirrhosis	37% (14/38)
		Cirrhosis	19% (5/26)
Treatment-experienced (FUSION)	SOF+RBV 16 weeks	Overall	62% (39/63)
		No cirrhosis	63% (25/40)
		Cirrhosis	61% (14/23)
Treatment-naïve (VALENCE)	SOF+RBV 24 weeks	Overall	93% (98/105)
		No cirrhosis	94% (86/92)
		Cirrhosis	92% (12/13)
Treatment-experienced (VALENCE)	SOF+RBV 24 weeks	Overall	77% (112/145)
		No cirrhosis	85% (85/100)
		Cirrhosis	60% (27/45)
Treatment-naïve co-infected with HIV (PHOTON-1)	SOF+RBV 12 weeks	Overall	67% (28/42)
		No cirrhosis	67% (24/36)
		Cirrhosis	67% (4/6)
Treatment-experienced co-infected with HIV (PHOTON-1)	SOF+RBV 24 weeks	Overall ^a	92% (12/13)
		No cirrhosis ^a	100% (8/8)
		Cirrhosis ^a	80% (4/5)
Treatment-naïve (ELECTRON ^b)	SOF+PEG+RBV 12 weeks	Overall ^c	97% (38/39)
Treatment-experienced (LONESTAR-2 ^b)	SOF+PEG+RBV 12 weeks	Overall	83% (20/24)
		No cirrhosis	83% (10/12)
		Cirrhosis	83% (10/12)

n = number of subjects with SVR12 response; N = total number of subjects per group.

- These data are preliminary.
- These are exploratory or Phase 2 studies. The outcomes should be interpreted with caution, as subject numbers are small and SVR rates may be impacted by the selection of patients. In the ELECTRON study (N = 11), the duration of peginterferon alfa ranged from 4-12 weeks in combination with sofosbuvir + ribavirin.
- All patients were non-cirrhotic in these two studies.

Table 17: Outcomes by therapeutic regimen and treatment duration, a comparison across studies in genotype 4 HCV infection

Patient population (Study number/name)	Regimen/Duration	Subgroup	SVR12 rate % (n/N)
Treatment-naïve (NEUTRINO)	SOF+PEG+RBV 12 weeks	Overall	96% (27/28)
		No cirrhosis	100% (27/27)
		Cirrhosis	0% (0/1)

n = number of subjects with SVR12 response; N = total number of subjects per group.

PHARMACOKINETICS

Sofosbuvir is a nucleotide prodrug that is extensively metabolised. The active metabolite is formed in hepatocytes and not observed in plasma. The predominant (>90%) metabolite, GS-331007, is inactive. It is formed through sequential and parallel pathways to the formation of active metabolite.

Absorption

The pharmacokinetic properties of sofosbuvir and the predominant circulating metabolite GS-331007 have been evaluated in healthy adult subjects and in subjects with chronic hepatitis C. Following oral administration, sofosbuvir was absorbed quickly and the peak plasma concentration was observed ~0.5-2 hour post-dose, regardless of dose level. Peak plasma concentration of GS-331007 was observed between 2 to 4 hours post-dose. Relative to healthy subjects, the sofosbuvir and GS-331007 AUC₀₋₂₄ was 57% higher and 39% lower, respectively in HCV infected subjects.

Effects of food

Relative to fasting conditions, the administration of a single dose of sofosbuvir with a standardised high fat meal slowed the rate of absorption of sofosbuvir. The extent of absorption of sofosbuvir was increased approximately 1.8-fold, with little effect on peak concentration. The exposure to GS-331007 was not altered in the presence of a high-fat meal.

Distribution

Sofosbuvir is not a substrate for hepatic uptake transporters, organic anion-transporting polypeptide (OATP) 1B1 or 1B3, and organic cation transporter (OCT) 1. While subject to active tubular secretion, GS-331007 is not a substrate for renal transporters including organic anion transporter (OAT) 1 or 3, OCT2, MRP2, P-gp, BCRP or MATE1. Sofosbuvir and GS-331007 are not inhibitors of drug transporters P-gp, BCRP, MRP2, BSEP, OATP1B1, OATP1B3 and OCT1. GS-331007 is not an inhibitor of OAT1, OCT2, and MATE1.

Sofosbuvir is approximately 85% bound to human plasma proteins (ex vivo data) and the binding is independent of drug concentration over the range of 1 µg/mL to 20 µg/mL. Protein binding of GS-331007 was minimal in human plasma. After a single 400 mg dose of [14C]-sofosbuvir in healthy subjects, the blood to plasma ratio of 14C-radioactivity was approximately 0.7.

Biotransformation

Sofosbuvir is extensively metabolised in the liver to form the pharmacologically active nucleoside analog triphosphate GS-461203. The metabolic activation pathway involves sequential hydrolysis of the carboxyl ester moiety catalysed by human cathepsin A (CatA) or carboxylesterase 1 (CES1) and phosphoramidate cleavage by histidine triad nucleotide-binding protein 1 (HINT1) followed by phosphorylation by the pyrimidine nucleotide biosynthesis pathway. Dephosphorylation results in the formation of nucleoside metabolite GS-331007 that cannot be efficiently rephosphorylated and lacks anti-HCV activity in vitro. Sofosbuvir and GS-331007 are not substrates or inhibitors of UGT1A1 or CYP3A4, CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, and CYP2D6 enzymes.

After a single 400 mg oral dose of [¹⁴C]-sofosbuvir, sofosbuvir and GS-331007 accounted for approximately 4% and >90% of drug-related material (sum of molecular weight-adjusted AUC of sofosbuvir and its metabolites) systemic exposure, respectively.

Elimination

The data indicate that renal clearance is the major elimination pathway for GS-331007 with a large part actively secreted. The median terminal half-lives of sofosbuvir and GS-331007 were 0.4 and 27 hours respectively.

Linearity/non-linearity

Sofosbuvir and GS-331007 AUCs are near dose proportional over the dose range of 200 mg to 400 mg.

Pharmacokinetics in special populations Gender and race

No clinically relevant pharmacokinetic differences due to gender or race have been identified for sofosbuvir and GS-331007.

Elderly

Population pharmacokinetic analysis in HCV infected subjects showed that within the age range (19 to 75 years) analysed, age did not have a clinically relevant effect on the exposure to sofosbuvir and GS-331007. Clinical studies of sofosbuvir included 65 subjects aged 65 and over. The response rates observed for subjects over 65 years of age were similar to that of younger subjects across treatment groups.

Renal impairment

The pharmacokinetics of sofosbuvir were studied in HCV negative subjects with mild (eGFR ≥ 50 and < 80 mL/min/1.73 m²), moderate (eGFR ≥ 30 and < 50 mL/min/1.73 m²), severe renal impairment (eGFR < 30 mL/min/1.73 m²) and subjects with ESRD requiring haemodialysis following a single 400 mg dose of sofosbuvir. Relative to subjects with normal renal function (eGFR > 80 mL/min/1.73 m²), the sofosbuvir AUC_{0-inf} was 61%, 107% and 171% higher in mild, moderate and severe renal impairment, while the GS-331007 AUC_{0-inf} was 55%, 88% and 451% higher, respectively. In subjects with ESRD, relative to subjects with normal renal function, sofosbuvir AUC_{0-inf} was 28% higher when sofosbuvir was dosed 1 hour before haemodialysis compared with 60% higher when sofosbuvir was dosed 1 hour after haemodialysis. The AUC_{0-inf} of GS-331007 in subjects with ESRD could not be reliably determined. However, data indicate at least 10-fold and 20-fold higher exposure to GS-331007 in ESRD compared to normal subjects when Sofosbuvir was administered 1 hour before or 1 hour after haemodialysis, respectively.

Haemodialysis can efficiently remove (53% extraction ratio) the predominant circulating metabolite GS-331007. A 4-hour haemodialysis session removed approximately 18% of administered dose. No dose adjustment is required for patients with mild or moderate renal impairment. The safety of Sofosbuvir has not been assessed in patients with severe renal impairment or ESRD.

Hepatic impairment

Population pharmacokinetics analysis in HCV infected subjects indicated that cirrhosis had no clinically relevant effect on the exposure to sofosbuvir and GS-331007. No dose adjustment of sofosbuvir is recommended for patients with mild, moderate and severe hepatic impairment.

Paediatric population

The pharmacokinetics of sofosbuvir and GS-331007 in paediatric subjects have not been established.
Pharmacokinetic/pharmacodynamic relationship(s)

Efficacy, in terms of rapid virologic response, has been shown to correlate with exposure to sofosbuvir as well as GS-331007. However, neither of these entities has been evidenced to be a general surrogate marker for efficacy (SVR12) at the therapeutic 400 mg dose.

INDICATION

Sofosvir is indicated in combination with other medicinal products for the treatment of chronic hepatitis C (CHC) in adults (see Pharmacology: Pharmacodynamics, Warning and Precaution, Dosage & Administration).

For hepatitis C virus (HCV) genotype specific activity, see Pharmacology: Pharmacodynamics, Warning and Precaution.

RECOMMENDED DOSE

Sofosvir treatment should be initiated and monitored by a physician experienced in the management of patients with CHC.

The recommended dose is one 400 mg tablet, taken orally, once daily with food (see Pharmacology: Pharmacokinetics).

Sofosvir should be used in combination with other medicinal products. Monotherapy of Sofosvir is not recommended (see Pharmacology: Pharmacodynamics). Refer also to the Summary of Product Characteristics of the medicinal products that are used in combination with Sofosvir. The recommended co-administered medicinal product(s) and treatment duration for Sofosvir combination therapy are provided in the table below.

Recommended co-administered medicinal product(s) and treatment duration for Sofosvir combination therapy

Patient Population*	Treatment	Duration
Patients with genotype 1, 4 CHC	Sofosvir + ribavirin + peginterferon alfa	12 weeks ^{a, b}
Patients with genotype 2 CHC	Sofosvir + ribavirin	12 weeks ^b
Patients with genotype 3 CHC	Sofosvir + ribavirin + peginterferon alfa	12 weeks ^b
	Sofosvir + ribavirin	24 weeks
Patients with genotype 1, 2, 3 or 4 CHC awaiting liver transplantation	Sofosvir + ribavirin	Until liver transplantation ^c

* includes patients co-infected with human immunodeficiency virus (HIV)

^a For previously treated patients with HCV genotype 1 infection, no data exist with the combination of Sofosvir, ribavirin and peginterferon alfa

^b Consideration should be given to potentially extending the duration of therapy beyond 12 weeks and up to 24 weeks; especially for those subgroups who have one or more factors historically associated with lower response rates to interferon-based therapies (eg, advanced fibrosis/cirrhosis, high baseline viral concentrations, black race, IL28B non CC genotype prior null response to peginterferon alfa and ribavirin therapy)

^c See special patient populations – Patient hepatocellular awaiting liver transplantation

Sofosvir in combination with ribavirin for 24 weeks can be considered as a therapeutic option for CHC patients with genotype 1 infection who are ineligible to receive an interferon-based regimen (see Pharmacology: Pharmacodynamics and Precautions). Treatment decision should be guided by an assessment of the potential benefits and risks for the individual patient.

The dose of ribavirin, when used in combination with Sofovir is weight-based (<75 kg = 1,000 mg and ≥75 kg = 1,200 mg) and administered orally in two divided doses with food.

Concerning co-administration with other direct-acting antivirals against HCV, see Precautions.

Dose modification: Dose reduction of Sofovir is not recommended.

If sofosbuvir is used in combination with peginterferon alfa, and a patient has a serious adverse reaction potentially related to this drug, the peginterferon alfa dose should be reduced or discontinued.

Refer to the peginterferon alfa Summary of Product Characteristics for additional information about how to reduce and/or discontinue the peginterferon alfa dose.

If a patient has a serious adverse reaction potentially related to ribavirin, the ribavirin dose should be modified or discontinued, if appropriate, until the adverse reaction abates or decreases in severity.

The table below provides guidelines for dose modifications and discontinuation based on the patient's haemoglobin concentration and cardiac status.

Ribavirin dose modification guideline for co-administration with Sofovir

Laboratory values	Reduce ribavirin dose to 600 mg/day if:	Discontinue ribavirin if:
Haemoglobin in subjects with no cardiac disease	< 10 g/dL	< 8.5 g/dL
Haemoglobin in subjects with history of stable cardiac disease	≥ 2 g/dL decrease in haemoglobin during any 4 weeks treatment period	< 12 g/dL despite 4 weeks at reduced rate

Once ribavirin has been withheld due to either a laboratory abnormality or clinical manifestation, an attempt may be made to restart ribavirin at 600 mg daily and further increase the dose to 800 mg daily.

However, it is not recommended that ribavirin be increased to the original assigned dose (1,000 mg to 1,200 mg daily).

Discontinuation of dosing: If the other medicinal products used in combination with Sofovir are permanently discontinued, Sofovir should also be discontinued (see Precautions).

Special patient populations: Elderly: No dose adjustment is warranted for elderly patients (see Pharmacology: Pharmacokinetics).

Renal impairment: No dose adjustment of Sofovir is required for patients with mild or moderate renal impairment. The safety and appropriate dose of Sofovir have not been established in patients with severe renal impairment (estimated glomerular filtration rate [eGFR] <30 mL/min/1.73 m²) or end stage renal disease (ESRD) requiring haemodialysis (see Pharmacology: Pharmacokinetics).

Hepatic impairment: No dose adjustment of Sofovir is required for patients with mild, moderate or severe hepatic impairment (Child-Pugh-Turcotte [CPT] class A, B or C) (see Pharmacology: Pharmacokinetics). The safety and efficacy of Sofovir have not been established in patients with decompensated cirrhosis.

Patients awaiting liver transplantation: The duration of administration of Sofovir in patients awaiting liver transplantation should be guided by an assessment of the potential benefits and risks for the individual patient (see Pharmacology: Pharmacodynamics).

Liver transplant recipients: Sofovir in combination with ribavirin is recommended for 24 weeks in liver transplant recipients. A starting ribavirin dose of 400 mg administered orally in two divided doses with food is recommended. If the starting dose of ribavirin is well-tolerated, the dose can be titrated up to a maximum of 1,000-1,200 mg daily (1,000 mg for patients weighing <75 kg and 1,200 mg for patients weighing ≥75 kg). If the starting dose of ribavirin is not well-tolerated, the dose

should be reduced as clinically indicated based on haemoglobin levels (see Pharmacology: Pharmacodynamics).

Paediatric population: The safety and efficacy of Sofosvir in children and adolescents aged <18 years have not yet been established. No data are available.

Administration: The film-coated tablet is for oral use. Patients should be instructed to swallow the tablet whole. The film-coated tablet should not be chewed or crushed, due to the bitter taste of the active substance. The tablet should be taken with food (see Pharmacology: Pharmacokinetics). Patients should be instructed that if vomiting occurs within 2 hours of dosing an additional tablet should be taken. If vomiting occurs more than 2 hours after dosing, no further dose is needed. These recommendations are based on the absorption kinetics of sofosbuvir and GS-331007 suggesting that the majority of the dose is absorbed within 2 hours after dosing.

If a dose is missed and it is within 18 hours of the normal time, patients should be instructed to take the tablet as soon as possible and then patients should take the next dose at the usual time. If it is after 18 hours then patients should be instructed to wait and take the next dose at the usual time. Patients should be instructed not to take a double dose.

CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients.

WARNINGS AND PRECAUTIONS

General

Sofosbuvir is not recommended for administration as monotherapy and should be prescribed in combination with other medicinal products for the treatment of hepatitis C infection. If the other medicinal products used in combination with Sofosbuvir are permanently discontinued, Sofosbuvir should also be discontinued. Consult the Summary of Product Characteristics for co-prescribed medicinal products before starting therapy with Sofosbuvir.

Severe bradycardia and heart block

Cases of severe bradycardia and heart block have been observed when Sofosvir is used in combination with daclatasvir and concomitant amiodarone with or without other drugs that lower heart rate. The mechanism is not established.

The concomitant use of amiodarone was limited through the clinical development of sofosbuvir plus direct-acting antivirals (DAAs). Cases are potentially life threatening, therefore amiodarone should only be used in patients on Sofosvir + daclatasvir when other alternative anti-arrhythmic treatments are not tolerated or are contraindicated.

Should concomitant use of amiodarone be considered necessary it is recommended that patients are closely monitored when initiating Sofosvir + daclatasvir. Patients who are identified as being high risk of bradyarrhythmia should be continuously monitored for 48 hours in an appropriate clinical setting.

Due to the long half-life of amiodarone, appropriate monitoring should also be carried out for patients who have discontinued amiodarone within the past few months and are to be initiated on Sofosvir in combination with daclatasvir.

All patients receiving Sofosvir + daclatasvir in combination with amiodarone with or without other drugs that lower heart rate should also be warned of the symptoms of bradycardia and heart block and should be advised to seek medical advice urgently should they experience them.

Treatment-experienced patients with genotype 1, 4, 5 and 6 HCV infection

Sofosbuvir has not been studied in a Phase 3 study in treatment-experienced patients with genotype 1, 4, 5 and 6 HCV infection. Thus, the optimal treatment duration in this population has not been established.

Consideration should be given to treating these patients, and potentially extending the duration of therapy with sofosbuvir, peginterferon alfa and ribavirin beyond 12 weeks and up to 24 weeks; especially for those subgroups who have one or more factors historically associated with lower response rates to interferon-based therapies (advanced fibrosis/cirrhosis, high baseline viral concentrations, black race, IL28B non CC genotype).

Treatment of patients with genotype 5 or 6 HCV infection

The clinical data to support the use of Sofosbuvir in patients with genotype 5 and 6 HCV infection is very limited.

Interferon-free therapy for genotype 1, 4, 5 and 6 HCV infection

Interferon-free regimens for patients with genotype 1, 4, 5 and 6 HCV infection with Sofosbuvir have not been investigated in Phase 3 studies. The optimal regimen and treatment duration have not been established. Such regimens should only be used for patients that are intolerant to or ineligible for interferon therapy, and are in urgent need of treatment.

Co-administration with other direct-acting antivirals against HCV

Sofosbuvir should only be co-administered with other direct-acting antiviral medicinal products if the benefit is considered to outweigh the risks based upon available data. There are no data to support the co-administration of Sofosbuvir and telaprevir or boceprevir. Such co-administration is not recommended.

Pregnancy and concomitant use with ribavirin

When Sofosbuvir is used in combination with ribavirin or peginterferon alfa/ribavirin, women of childbearing potential or their male partners must use an effective form of contraception during the treatment and for a period of time after the treatment as recommended in the Summary of Product Characteristics for ribavirin. Refer to the Summary of Product Characteristics for ribavirin for additional information.

Use with potent P-gp inducers

Medicinal products that are potent P-glycoprotein (P-gp) inducers in the intestine (e.g. rifampicin, St. John's wort [*Hypericum perforatum*], carbamazepine and phenytoin) may significantly decrease sofosbuvir plasma concentration leading to reduced therapeutic effect of Sofosbuvir. Such medicinal products should not be used with Sofosbuvir.

Use in diabetic patients

Diabetics may experience improved glucose control, potentially resulting in symptomatic hypoglycaemia, after initiating HCV direct-acting antiviral treatment. Glucose levels of diabetic patients initiating direct-acting antiviral therapy should be closely monitored, particularly within the first 3 months, and their diabetic medication modified when necessary. The physician in charge of the diabetic care of the patient should be informed when direct-acting antiviral therapy is initiated.

Renal impairment

The safety of Sofosbuvir has not been assessed in subjects with severe renal impairment (eGFR <30 mL/min/1.73 m²) or ESRD requiring haemodialysis. Furthermore, the appropriate dose has not been established. When Sofosbuvir is used in combination with ribavirin or peginterferon alfa/ribavirin, refer also to the Summary of Product Characteristics for ribavirin for patients with creatinine clearance (CrCl) <50 mL/min

HCV/HBV (hepatitis B virus) co-infection

There are no data on the use of Sofosbuvir in patients with HCV/HBV co-infection.

Pediatric population

Sofosbuvir is not recommended for use in children and adolescents under 18 years of age because the safety and efficacy have not been established in this population.

Elderly patients

Clinical studies of sofosbuvir included 65 subjects aged 65 and over. The response rates observed for subjects over 65 years of age were similar to that of younger subjects across treatment groups. In

general, caution should be exercised when administering sofosbuvir in elderly patients reflecting the greater frequency of anaemia, decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Post-liver transplant patients

The safety and efficacy of Sofosbuvir have not been established in post-liver transplant patients.

INTERACTIONS WITH OTHER MEDICAMENTS

Sofosbuvir is a nucleotide prodrug. After oral administration of Sofosbuvir, sofosbuvir is rapidly absorbed and subject to extensive first-pass hepatic and intestinal metabolism. Intracellular hydrolytic prodrug cleavage catalysed by enzymes including carboxylesterase 1 and sequential phosphorylation steps catalysed by nucleotide kinases result in formation of the pharmacologically active uridine nucleoside analogue triphosphate. The predominant inactive circulating metabolite GS-331007 that accounts for greater than 90% of drug-related material systemic exposure is formed through pathways sequential and parallel to formation of active metabolite. The parent sofosbuvir accounts for approximately 4% of drug-related material systemic exposure. In clinical pharmacology studies, both sofosbuvir and GS-331007 were monitored for purposes of pharmacokinetic analyses.

Sofosbuvir is a substrate of drug transporter P-gp and breast cancer resistance protein (BCRP) while GS-331007 is not. Medicinal products that are potent P-gp inducers in the intestine (e.g. rifampicin, St. John's wort, carbamazepine and phenytoin) may decrease sofosbuvir plasma concentration leading to reduced therapeutic effect of Sofosbuvir and thus should not be used with Sofosbuvir. Co-administration of Sofosbuvir with medicinal products that inhibit P-gp and/or BCRP may increase sofosbuvir plasma concentration without increasing GS-331007 plasma concentration, thus Sofosbuvir may be co-administered with P-gp and/or BCRP inhibitors. Sofosbuvir and GS-331007 are not inhibitors of P-gp and BCRP and thus are not expected to increase exposures of medicinal products that are substrates of these transporters.

The intracellular metabolic activation pathway of sofosbuvir is mediated by generally low affinity and high capacity hydrolase and nucleotide phosphorylation pathways that are unlikely to be affected by concomitant medicinal products.

Other interactions

Drug interaction information for Sofosbuvir with potential concomitant medicinal products is summarised in Table 3 below (where 90% confidence interval (CI) of the geometric least-squares mean (GLSM) ratio were within “↔”, extended above “↑”, or extended below “↓” the predetermined equivalence boundaries). The table is not all-inclusive.

Medicinal product by therapeutic areas	Effects on drug levels. Mean ratio (90% confidence interval) for AUC, C _{max} , C _{min}	Recommendation concerning co administration with Sofosbuvir
ANALEPTICS		

Modafinil	Interaction not studied. <i>Expected:</i> ↓ Sofosbuvir ↓ GS-331007	Co-administration of Sofosbuvir with modafinil is expected to decrease the concentration of sofosbuvir, leading to reduced therapeutic effect of Sofosbuvir. Such co-administration is not recommended.
ANTIARRHYTHMICS		
Amiodarone	Interaction not studied	Use only if no other alternative is available. Close monitoring is recommended if this medicinal product is administered with Sofosvir + Daclatasvir (See Precautions and Adverse Reactions)
ANTICONVULSANTS		
Carbamazepine Phenytoin Phenobarbital Oxcarbazepine	Interaction not studied. <i>Expected:</i> ↓ Sofosbuvir ↔ GS-331007	Co-administration of Sofosbuvir with carbamazepine, phenytoin, phenobarbital or oxcarbazepine is expected to decrease the concentration of sofosbuvir, leading to reduced therapeutic effect of Sofosbuvir. Such co-administration is not recommended. Sofosbuvir should not be used with carbamazepine, phenytoin, phenobarbital or oxcarbazepine, potent intestinal P-gp inducers.
ANTIMYCOBACTERIALS		
Rifampicin (600mg single dose)	Interaction not studied. <i>Expected:</i> ↓ Sofosbuvir ↔ GS-331007	Sofosbuvir contraindicated with rifampicin, a potent intestinal P-gp inducer
Rifabutin Rifapentine	Interaction not studied. <i>Expected:</i> ↓ Sofosbuvir ↔ GS-331007	Sofosbuvir is contraindicated with rifabutin, a potent intestinal P-gp inducer Co-administration of Sofosbuvir with rifapentine is expected to decrease the concentration of sofosbuvir, leading to reduced therapeutic effect of Sofosbuvir. Such co-administration is not recommended.

HERBAL SUPPLEMENTS		
St. John's wort (<i>Hypericum perforatum</i>)	Interaction not studied. <i>Expected:</i> ↓ Sofosbuvir ↓ GS-331007	Sofosbuvir should not be used with St. John's wort, a potent intestinal P-gp inducer.
HCV ANITIVIRAL AGENTS: HCV PROTEASE INHIBITORS		
Boceprevir (BOC) Telaprevir (TPV)	Interaction not studied. <i>Expected:</i> ↑ Sofosbuvir (TPV) ↔ Sofosbuvir (BOC) ↔ GS-331007 (TPV or BOC)	No drug-drug interaction data exists regarding the co-administration of Sofosbuvir with boceprevir or telaprevir.
NARCOTIC ANALGESICS		
Methadone (Methadone maintenance therapy [30 to 130 mg/daily])	↔ R-methadone ↔ S-methadone <i>Sofosbuvir</i> ↓ C _{max} ↑ AUC <i>GS-331007</i> ↓ C _{max} ↔ AUC	No dose adjustment of sofosbuvir or methadone is required when sofosbuvir and methadone are used concomitantly.
IMMUNOSUPPRESSANTS		
Ciclosporin (600 mg single dose)	↔ <i>Ciclosporin</i> ↑ <i>Sofosbuvir</i> <i>GS-331007</i> ↓ C _{max} ↔ AUC	No dose adjustment of sofosbuvir or ciclosporin is required when sofosbuvir and ciclosporin are used concomitantly

Tacrolimus (5 mg single dose)	<i>Tacrolimus</i> ↓ C_{max} ↔ AUC <i>Sofosbuvir</i> ↓ C_{max} ↑ AUC ↔ GS-331007	No dose adjustment of sofosbuvir or tacrolimus is required when sofosbuvir and tacrolimus are used concomitantly.
HIV ANTIVIRAL AGENTS: REVERSE TRANSCRIPTASE INHIBITORS		
Efavirenz (600 mg once daily) Administered as a single Tenofovir Disoproxil Fumarate 300mg + Emtricitabine 200mg + Efavirenz 600mg Tablet	↔ <i>Efavirenz</i> <i>Sofosbuvir</i> ↓ C_{max} ↔ AUC GS-331007 ↓ C_{max} ↔ AUC	No dose adjustment of sofosbuvir or efavirenz is required when sofosbuvir and efavirenz are used concomitantly.
Emtricitabine (200 mg once daily) Administered as a single Tenofovir Disoproxil Fumarate 300mg + Emtricitabine 200mg + Efavirenz 600mg Tablet	↔ <i>Emtricitabine</i> <i>Sofosbuvir</i> ↓ C_{max} ↔ AUC GS-331007 ↓ C_{max} ↔ AUC	No dose adjustment of sofosbuvir or emtricitabine is required when sofosbuvir and emtricitabine are used concomitantly.
Tenofovir disoproxil fumarate (300 mg once daily) Administered as a single Tenofovir Disoproxil Fumarate 300mg + Emtricitabine 200mg + Efavirenz 600mg Tablet	<i>Tenofovir</i> ↑ C_{max} ↔ AUC <i>Sofosbuvir</i> ↓ C_{max} ↔ AUC GS-331007 ↓ C_{max} ↔ AUC	No dose adjustment of sofosbuvir or tenofovir disoproxil fumarate is required when sofosbuvir and tenofovir disoproxil fumarate are used concomitantly.

Rilpivirine (25 mg once daily)	↔ <i>Rilpivirine</i> <i>Sofosbuvir</i> ↑ C _{max} ↔ AUC ↔ <i>GS-331007</i>	No dose adjustment of sofosbuvir or rilpivirine is required when sofosbuvir and rilpivirine are used concomitantly.
HIV ANTIVIRAL AGENTS: HIV PROTEASE INHIBITORS		
Darunavir boosted with ritonavir (800/100 mg once daily)	↔ <i>Darunavir</i> ↑ <i>Sofosbuvir</i> ↔ <i>GS-331007</i>	No dose adjustment of sofosbuvir or darunavir (ritonavir boosted) is required when sofosbuvir and darunavir are used concomitantly.
HIV ANTIVIRAL AGENTS: INTEGRASE INHIBITORS		
Raltegravir (400 mg twice daily)	↓ <i>Raltegravir</i> ↔ <i>Sofosbuvir</i> ↔ <i>GS-331007</i>	No dose adjustment of sofosbuvir or raltegravir is required when sofosbuvir and raltegravir are used concomitantly.
ORAL CONTRACEPTIVES		
Norgestimate/ethinyl estradiol	↔ <i>Norgestromin</i> ↔ <i>Norgestrel</i> ↔ <i>Ethinyl estradiol</i>	No dose adjustment of norgestimate/ethinyl estradiol is required when sofosbuvir and norgestimate/ethinyl estradiol are used concomitantly.

Medicinal products that are potent P-gp inducers in the intestine (rifampicin, St. John's wort, carbamazepine and phenytoin) may significantly decrease sofosbuvir plasma concentration leading to reduced therapeutic effect. For this reason, sofosbuvir should not be co-administered with known inducers of P-gp.

PREGNANCY AND LACTATION

Women of childbearing potential / contraception in males and females

When Sofosbuvir is used in combination with ribavirin or peginterferon alfa/ribavirin, extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients. Significant teratogenic and/or embryocidal effects have been demonstrated in all animal species exposed to ribavirin. Women of childbearing potential or their male partners must use an effective

form of contraception during treatment and for a period of time after the treatment has concluded as recommended in the Summary of Product Characteristics for ribavirin. Refer to the Summary of Product Characteristics for ribavirin for additional information.

Pregnancy

There are no or limited amount of data (less than 300 pregnancy outcomes) from the use of sofosbuvir in pregnant women.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. No effects on foetal development have been observed in rats and rabbits at the highest doses tested. However, it has not been possible to fully estimate exposure margins achieved for sofosbuvir in the rat relative to the exposure in humans at the recommended clinical dose.

As a precautionary measure, it is preferable to avoid the use of Sofosbuvir during pregnancy. However, if ribavirin is co-administered with sofosbuvir, the contraindications regarding use of ribavirin during pregnancy apply (see also the Summary of Product Characteristics for ribavirin).

Breast-feeding

It is unknown whether sofosbuvir and its metabolites are excreted in human milk. Available pharmacokinetic data in animals has shown excretion of metabolites in milk. A risk to newborns/infants cannot be excluded. Therefore, Sofosbuvir should not be used during breast-feeding.

Fertility

No human data on the effect of Sofosbuvir on fertility are available. Animal studies do not indicate harmful effects on fertility.

SIDE EFFECTS

Summary of the safety profile

During treatment with sofosbuvir in combination with ribavirin or with peginterferon alfa and ribavirin, the most frequently reported adverse drug reactions were consistent with the expected safety profile of ribavirin and peginterferon alfa treatment, without increasing the frequency or severity of the expected adverse drug reactions.

Tabulated summary of adverse reactions

Sofosbuvir has been studied in combination with ribavirin, with or without peginterferon alfa. In this context, no adverse drug reactions specific to sofosbuvir have been identified. The most common adverse drug reactions occurring in patients receiving sofosbuvir and ribavirin or sofosbuvir, ribavirin and peginterferon alfa were fatigue, headache, nausea and insomnia.

The following adverse drug reactions have been identified with sofosbuvir in combination with ribavirin or in combination with peginterferon alfa and ribavirin (the table below). The adverse reactions are listed below by body system organ class and frequency. Frequencies are defined as follows: very common, common, uncommon, rare or very rare

Adverse drug reactions identified with sofosbuvir in combination with ribavirin or peginterferon alfa and ribavirin

Frequency	SOF^a + RBV^b	SOF + PEG^c + RBV
<i>Infections and infestations:</i>		
Common	nasopharyngitis	
<i>Blood and lymphatic system disorders:</i>		
Very common	haemoglobin decreased	anaemia, neutropenia, lymphocyte count decreased, platelet count decreased
Common	anaemia	
<i>Metabolism and nutrition disorders:</i>		
Very common		decreased appetite
Common		weight decreased
<i>Psychiatric disorders:</i>		
Very common	insomnia	insomnia
Common	depression	depression, anxiety, agitation
<i>Nervous system disorders:</i>		
Very common	headache	dizziness, headache
Common	disturbance in attention	migraine, memory impairment, disturbance in attention
<i>Eye disorders:</i>		
Common		vision blurred
<i>Respiratory, thoracic and mediastinal disorders:</i>		
Very common		dyspnoea, cough
Common	dyspnoea, dyspnoea exertional, cough	dyspnoea exertional
<i>Gastrointestinal disorders:</i>		
Very common	nausea	diarrhoea, nausea, vomiting
Common	Abdominal discomfort, constipation, dyspepsia	constipation, dry mouth, gastroesophageal reflux
<i>Hepatobiliary disorders:</i>		
Very common	blood bilirubin increased	blood bilirubin increased
<i>Skin and subcutaneous tissue disorders:</i>		
Very common		rash, pruritus
Common	alopecia, dry skin, pruritus	alopecia, dry skin
<i>Musculoskeletal and connective tissue disorders:</i>		
Very common		arthralgia, myalgia
Common	arthralgia, back pain, muscle spasms, myalgia	back pain, muscle spasms
<i>General disorders and administration site conditions:</i>		
Very common	fatigue, irritability	chills, fatigue, influenza-like illness, irritability, pain, pyrexia
Common	pyrexia, asthenia	chest pain, asthenia

a. SOF = sofosbuvir; b. RBV = ribavirin; c. PEG = peginterferon alfa.

Other special population(s)

HIV/HCV co-infection

The safety profile of sofosbuvir and ribavirin in HCV/HIV co-infected subjects was similar to that observed in mono-infected HCV subjects treated with sofosbuvir and ribavirin

Patients awaiting liver transplantation

The safety profile of sofosbuvir and ribavirin in HCV infected subjects prior to liver transplantation was similar to that observed in subjects treated with sofosbuvir and ribavirin

Description of selected adverse reactions:

Cardiac arrhythmias: Cases of severe bradycardia and heart block have been observed when Sofosbuvir is used in combination with daclatasvir and concomitant amiodarone and/or other drugs that lower heart rate (see Precautions and Interactions).

Less Common Adverse Reactions Reported in Clinical Trials (<1%):

The following adverse drug reactions occurred in <1% of subjects receiving sofosbuvir in combination with ribavirin or peginterferon alfa and ribavirin. These events have been included because of their seriousness or assessment of potential causal relationship.

Hematologic Effects: Pancytopenia (particularly in subjects receiving concomitant pegylated interferon).

Psychiatric Disorders: Severe depression (particularly in subjects with pre-existing history of psychiatric illness), including suicidal ideation and suicide.

SYMPTOMS AND TREATMENT OF OVERDOSE

The highest documented dose of sofosbuvir was a single supra therapeutic dose of sofosbuvir 1,200 mg administered to 59 healthy subjects. In that study, there were no untoward effects observed at this dose level, and adverse reactions were similar in frequency and severity to those reported in the placebo and sofosbuvir 400 mg treatment groups. The effects of higher doses are unknown.

No specific antidote is available for overdose with Sofosbuvir. If overdose occurs the patient must be monitored for evidence of toxicity. Treatment of overdose with Sofosbuvir consists of general supportive measures including monitoring of vital signs as well as observation of the clinical status of the patient. Haemodialysis can efficiently remove (53% extraction ratio) the predominant circulating metabolite GS-331007. A 4-hour haemodialysis session removed 18% of the administered dose.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Sofosbuvir has moderate influence on the ability to drive and use machines. Patients should be informed that fatigue and disturbance in attention, dizziness and blurred vision have been reported during treatment with sofosbuvir in combination with peginterferon alfa and ribavirin (see Adverse Reactions).

STORAGE CONDITION

Store below 30°C and protect from moisture

SHELF LIFE

2 Years

CONTAINER PACK

28's tablets in HDPE Bottle

MANUFACTURER

HETERO LABS LIMITED

UNIT-V, Sy. No. 439, 440, 441 & 458, TSIIC Formulation SEZ,

Polepally Village, Jadcherla Mandal, Mahaboobnagar Dist., Telangana, India.

PRODUCT REGISTRATION HOLDER

SYNERRV SDN BHD, S0-29-2, Menara 1, KL Eco City, Jalan Bangsar, Kg Haji Abdullah Hukum, 59200

Kuala Lumpur

DATE OF REVISION: August 2019