

Unwinding Direction

Each mL contains Oxaliplatin 5 mg
Never use sodium chloride or chloride containing solutions for dilution.
Keep medicine out of reach of children
Jauhkan Ubat daripada capaian Kanak-kanak
Ubat Terkawal / Controlled Medicine
Unopened Vial: Store below 30°C. Keep the vial in the outer carton in order to protect from light. Do not freeze. Dilution before infusion: After dilution with glucose 5% (50mg/ml) solution under aseptic conditions, chemical and physical in-use stability has been demonstrated for 48 hours at 2°C to 8°C and for 24 hours at 25°C. Refer to package insert for full details.

RECOXALI

Rx
Oxaliplatin 5 mg/mL Concentrate
for solution for Infusion **[50mg/10mL]**

Single Dose Vial
For Intravenous Use Only
Must be Diluted Before use



Caution:
Cytotoxic Agent

Manufactured by: **Eugia Pharma Specialities Limited**,
Sy No. 550, 551 552, Kolthur Village, Shamirpet Mandal,
Medchal-Malkajgiri District, Telangana 500101, INDIA.
Product Registration holder in Malaysia:
Synergiv Sdn Bhd, SO-29-2, Menara 1, KL ECO City,
Jalan Bangsar, KG Haji Abdullah Hukum,
59200 Kuala Lumpur Malaysia.
M.L. No.: 24/RR/TS/2015/F/G
MAL No:

P1429962

Batch No.:
Exp Date:
Mfg Date:

Coding Zone will be
over coded during
production(35 x 9.5 mm)

Size : 90 x 25 mm

* Batch:, Expiry: will be over coded during production

■ Pantone 639 C ■ Pantone 3275 C ■ Pantone 072 C ■ Black ■ Cool Gray 5 C

eugia Packaging Development		Product Name	Component	Item Code	Date & Time
		RECOXALI Inj 50 mg-10ml	Label	P1429962	10.01.2024 & 3:00 PM
		Customer / Country	Version No.	Reason Of Issue	Reviewed / Approved by
		Malaysia_Eugia	07	Submission	
Team Leader	Narender	Dimensions	No. of Colours : 05		
Initiator	Sagar	90 x 25 mm			
Artist:	Vision Graphic Designers	2D Code			
Additional Information :		NA			
					Sign / Date