

Pharmacode position may change as per Supplier’s m/c requirement & additional small pharma code may appear on the front / back panel



SUMMARY OF PRODUCT CHARACTERISTICS
AZASQUARE

Azacitidine Powder for Suspension for Injection 25 mg/mL (100 mg/Vial)

Name of the medicinal product

Azacitidine Powder for Suspension for Injection 25 mg/mL (100mg/ Vial)

Qualitative and quantitative composition

Each vial contains 100 mg Azacitidine. (After reconstitution, each mL of suspension contains 25 mg Azacitidine). For the full list of excipients, see section *List of excipients*.

Pharmaceutical form

Powder for suspension for injection.

Before reconstitution: White to off-white lyophilized cake or powder in clear glass vial stoppered with gray igloo lyo rubber stopper and sealed with aluminium seal having white colour with PP disc.

After reconstitution (with 4.0 mL water for injection): White to off-white uniform cloudy suspension.

Clinical particulars Therapeutic indications

Azacitidine is indicated for the treatment of adult patients who are not eligible for haematopoietic stem cell transplantation (HSCT) with:

- intermediate-2 and high-risk myelodysplastic syndromes (MDS) according to the International Prognostic Scoring System (IPSS),
- chronic myelomonocytic leukaemia (CMML) with 10-29 % marrow blasts without myeloproliferative disorder,
- acute myeloid leukaemia (AML) with 20-30 % blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) classification,

Posology and method of administration

Azacitidine treatment should be initiated and monitored under the supervision of a physician experienced in the use of chemotherapeutic agents. Patients should be premedicated with antiemetics for nausea and vomiting.

Posology

The recommended starting dose for the first treatment cycle, for all patients regardless of baseline haematology laboratory values, is 75 mg/m² of body surface area, injected subcutaneously, daily for 7 days, followed by a rest period of 21 days (28-day treatment cycle).

It is recommended that patients be treated for a minimum of 6 cycles. Treatment should be continued as long as the patient continues to benefit or until disease progression. Patients should be monitored for haematologic response/toxicity and renal toxicities (see section *Special warnings and precautions for use*); a delay in starting the next cycle or a dose reduction as described below may be necessary.

AZASQUARE should not be used interchangeably with oral Azacitidine. Due to differences in the exposure, the dose and schedule recommendations for oral Azacitidine are different from those for injectable Azacitidine. Healthcare professionals are recommended to verify the name of the medicinal product, dose and administration route.

Laboratory tests

Liver function tests, serum creatinine and serum bicarbonate should be determined prior to initiation of therapy and prior to each treatment cycle. Complete blood counts should be performed prior to initiation of therapy and as needed to monitor response and toxicity, but at a minimum, prior to each treatment cycle.

Dose adjustment due to haematological toxicity

Haematological toxicity is defined as the lowest count reached in a given cycle (nadir) if platelets ≤ 50.0 x 10⁹ /l and/ or absolute neutrophil count (ANC) ≤ 1 x 10⁹ /l.

Recovery is defined as an increase of cell line(s) where haematological toxicity was observed of at least half of the difference of nadir and the baseline count plus the nadir count (i.e., blood count at recovery ≥ nadir count + (0.5 x [baseline count – nadir count])).

Patients without reduced baseline blood counts (i.e., *White Blood Cells (WBC) ≥ 3.0 x 10⁹ /l and ANC ≥ 1.5 x 10⁹ /l, and platelets ≥ 75.0 x 10⁹ /l*) prior to the first treatment.

If haematological toxicity is observed following Azacitidine treatment, the next cycle of the therapy should be delayed until the platelet count and the ANC have recovered. If recovery is achieved within 14 days, no dose adjustment is necessary.

However, if recovery has not been achieved within 14 days, the dose should be reduced according to the following table. Following dose modifications, the cycle duration should return to 28 days.

Cycle Nadir Counts		% Dose in the next cycle, if recovery* is not achieved within 14 days
ANC (x 10 ⁹ /L)	Platelets (x 10 ⁹ /L)	
≤ 1.0	≤ 50.0	50 %
> 1.0	> 50.0	100 %

* Recovery – counts ≥ nadir count + (0.5 x [base line count – nadir count])

Patients with reduced baseline blood counts (i.e., *WBC < 3.0 x 10⁹ /L or ANC < 1.5 x 10⁹ /L or platelets < 75.0 x 10⁹ /l*) prior to the first treatment.

Following Azacitidine treatment, if the decrease in WBC or ANC or platelets from that prior to treatment is ≤ 50 %, or greater than 50 % but with an improvement in any cell line differentiation, the next cycle should not be delayed and no dose adjustment made.

If the decrease in WBC or ANC or platelets is greater than 50 % from that prior to treatment, with no improvement in cell line differentiation, the next cycle of Azacitidine therapy should be delayed until the platelet count and the ANC have recovered. If recovery is achieved within 14 days, no dose adjustment is necessary. However, if recovery has not been achieved within 14 days, bone marrow cellularity should be determined. If the bone marrow cellularity is > 50 %, no dose adjustments should be made. If bone marrow cellularity is ≤ 50 %, treatment should be delayed and the dose reduced according to the following table:

Bone marrow cellularity	Dose in the next cycle if recovery is not achieved within 14 days	
	%	
	Recovery * ≤ 21 days	Recovery * ≥ 21 days
15 – 50 %	100 %	50 %
< 15 %	100 %	33 %

* Recovery = counts ≥ nadir count + (0.5 x [baseline count – nadir count])

Following dose modifications, the cycle duration should return to 28 days.

Special populations

Elderly patients

No specific dose adjustments are recommended for the elderly. Because elderly patients are more likely to have decreased renal function, it may be useful to monitor renal function.

Patients with renal impairment

Azacitidine can be administered to patients with renal impairment without initial dose adjustment (see section *Pharmacokinetic properties*). If unexplained reductions in serum bicarbonate levels to less than 20 mmol/L occur, the dose should be reduced by 50 % on the next cycle. If unexplained elevations in serum creatinine or blood urea nitrogen (BUN) to ≥ 2-fold above baseline values and above upper limit of normal (ULN) occur, the next cycle should be delayed until values return to normal or baseline and the dose should be reduced by 50 % on the next treatment cycle (see section *Special warnings and precautions for use*).

Patients with hepatic impairment

No formal studies have been conducted in patients with hepatic impairment (see section *Special warnings and precautions for use*). Patients with severe hepatic organ impairment should be carefully monitored for adverse events. No specific modification to the starting dose is recommended for patients with hepatic impairment prior to starting treatment; subsequent dose modifications should be based on haematology laboratory values. Azacitidine is contraindicated in patients with advanced malignant hepatic tumours (see sections *Contraindications* and *Special warnings and precautions for use*).

Paediatric population

The safety and efficacy of Azacitidine in children aged 0-17 years have not yet been established. No data are available.

Method of administration

Reconstituted Azacitidine should be injected subcutaneously into the upper arm, thigh or abdomen.

Injection sites should be rotated. New injections should be given at least 2.5 cm from the previous site and never into areas where the site is tender, bruised, red, or hardened.

After reconstitution, the suspension should not be filtered. For instructions on reconstitution of the medicinal product before administration, see section *Special precautions for disposal and other handling*.

Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section *List of excipients*. Advanced malignant hepatic tumours (see section *Special warnings and precautions for use*). Breast-feeding (see section *Fertility, pregnancy and breastfeeding*).

Special warnings and precautions for use

Haematological toxicity

Treatment with Azacitidine is associated with anaemia, neutropenia and thrombocytopenia, particularly during the first 2 cycles (see section *Undesirable effects*). Complete blood counts should be performed as needed to monitor response and toxicity, but at least prior to each treatment cycle. After administration of the recommended dose for the first cycle, the dose for subsequent cycles should be reduced or its administration delayed based on nadir counts and haematological response (see section *Posology and method of administration*). Patients should be advised to promptly report febrile episodes. Patients and physicians are also advised to be observant for signs and symptoms of bleeding.

Hepatic impairment

No formal studies have been conducted in patients with hepatic impairment. Patients with extensive tumour burden due to metastatic disease have been reported to experience progressive hepatic coma and death during Azacitidine treatment, especially in such patients with baseline serum albumin < 30 g/L. Azacitidine is contraindicated in patients with advanced malignant hepatic tumours (see section *Contraindications*).

Renal impairment

Renal abnormalities ranging from elevated serum creatinine to renal failure and death were reported in patients treated with intravenous Azacitidine in combination with other chemotherapeutic agents. In addition, renal tubular acidosis, defined as a fall in serum bicarbonate to < 20 mmol/L in association with an alkaline urine and hypokalaemia (serum potassium < 3 mmol/L) developed in 5 subjects with chronic myelogenous leukaemia (CML) treated with Azacitidine and etoposide. If unexplained reductions in serum bicarbonate (< 20 mmol/L) or elevations of serum creatinine or BUN occur, the dose should be reduced or administration delayed (see section *Posology and method of administration*). Patients should be advised to report oliguria and anuria to the health care provider immediately.

Although no clinically relevant differences in the frequency of adverse reactions were noted between subjects with normal renal function compared to those with renal impairment, patients with renal impairment should be closely monitored for toxicity since Azacitidine and/ or its metabolites are primarily excreted by the kidney (see section *Posology and method of administration*).

Laboratory tests

Liver function tests, serum creatinine and serum bicarbonate should be determined prior to initiation of therapy and prior to each treatment cycle. Complete blood counts should be performed prior to initiation of therapy and as needed to monitor response and toxicity, but at a minimum, prior to each treatment cycle, see also section *Undesirable effects*.

Cardiac and pulmonary disease

Patients with a history of severe congestive heart failure, clinically unstable cardiac disease or pulmonary disease were excluded from the pivotal registration studies (AZA PH GL 2003 CL 001 and AZA-AML-001) and therefore the safety and efficacy of Azacitidine in these patients has not been established. Recent data from a clinical trial in patients with a known history of cardiovascular or pulmonary disease showed a significantly increased incidence of cardiac events with Azacitidine (see section *Undesirable effects*). It is therefore advised to exercise caution when prescribing Azacitidine to these patients. Cardiopulmonary assessment before and during the treatment should be considered.

Necrotising fasciitis

Necrotising fasciitis, including fatal cases, have been reported in patients treated with Azacitidine. Azacitidine therapy should be discontinued in patients who develop necrotising fasciitis and appropriate treatment should be promptly initiated.

Tumour lysis syndrome

The patients at risk of tumour lysis syndrome are those with high tumour burden prior to treatment. These patients should be monitored closely and appropriate precautions taken.

Differentiation syndrome

Cases of differentiation syndrome (also known as retinoic acid syndrome) have been reported in patients receiving injectable Azacitidine. Differentiation syndrome may be fatal and symptoms and clinical findings include respiratory distress, pulmonary infiltrates, fever, rash, pulmonary oedema, peripheral oedema, rapid weight gain, pleural effusions, pericardial effusions, hypotension and renal dysfunction (see section *Undesirable effects*). Treatment with high-dose IV corticosteroids and haemodynamic monitoring should be considered at first onset of symptoms or signs suggestive of differentiation syndrome. Temporary discontinuation of injectable Azacitidine should be considered until resolution of symptoms and if resumed, caution is advised.

Interaction with other medicinal products and other forms of interactions

Based on *in vitro* data, Azacitidine metabolism does not appear to be mediated by cytochrome P450 isoenzymes (CYPs), UDP-glucuronosyltransferases (UGTs), sulfotransferases (SULTs), and glutathione transferases (GSTs); interactions related to these metabolizing enzymes *in vivo* are therefore considered unlikely.

Clinically significant inhibitory or inductive effects of Azacitidine on cytochrome P450 enzymes are unlikely (see section *Pharmacokinetic properties*).

No formal clinical drug interaction studies with Azacitidine have been conducted.

Fertility, pregnancy and lactation

Women of childbearing potential / Contraception in males and females

Women of childbearing potential have to use effective contraception during and for at least 6 months after treatment. Men should be advised not to father a child while receiving treatment and must use effective contraception during and for at least 3 months after treatment.

Pregnancy

There are no adequate data on the use of Azacitidine in pregnant women. Studies in mice have shown reproductive toxicity. The potential risk for humans is unknown. Based on results from animal studies and its mechanism of action, Azacitidine should not be used during pregnancy, especially during the first trimester, unless clearly necessary. The advantages of treatment should be weighed against the possible risk for the foetus in every individual case.

Breast-feeding

It is not known whether Azacitidine or its metabolites are excreted in human milk. Due to the potential serious adverse reactions in the nursing child, breast-feeding is contraindicated during Azacitidine therapy.

Fertility

There are no human data on the effect of Azacitidine on fertility. In animals, adverse reactions with Azacitidine use on male fertility have been documented. Men should be advised not to father a child while receiving treatment and must use effective contraception during and up to 3 months after treatment. Before starting treatment, male patients should be advised to seek counselling on sperm storage.

Effects on ability to drive and use machines

Azaciitidine has minor or moderate influence on the ability to drive and use machines. Fatigue has been reported with the use of Azacitidine. Therefore, caution is recommended when driving or operating machines.

Undesirable effects

Summary of the safety profile

Adverse reactions considered to be possibly or probably related to the administration of Azacitidine have occurred in 97 % of patients.

The most common serious adverse reactions noted from the pivotal study (AZA PH GL 2003 CL 001) included febrile neutropenia (8.0 %) and anaemia (2.3 %), which were also reported in the supporting studies (CALGB 9221 and CALGB 8921). Other serious adverse reactions from these 3 studies included infections such as neutropenic sepsis (0.8 %) and pneumonia (2.5 %) (some with fatal outcome), thrombocytopenia (3.5 %), hypersensitivity reactions (0.25 %) and haemorrhagic events (e.g., cerebral haemorrhage [0.5 %], gastrointestinal haemorrhage [0.8 %] and intracranial haemorrhage [0.5%]).

The most commonly reported adverse reactions with Azacitidine treatment were haematological reactions (71.4 %) including thrombocytopenia, neutropenia and leukopenia (usually Grade 3-4), gastrointestinal events (60.6 %) including nausea, vomiting (usually Grade 1-2) or injection site reactions (77.1 %; usually Grade 1-2).

Tabulated list of adverse reactions

Table 1 below contains adverse reactions associated with Azacitidine treatment obtained from the main clinical studies in MDS and AML and post marketing surveillance.

Frequencies are defined as: very common (≥ 1/10), common (≥ 1/100 to < 1/10); uncommon (≥ 1/1,000 to < 1/100); rare (≥ 1/10,000 to < 1/1,000); very rare (< 1/10,000); not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. Adverse reactions are presented in the table below according to the highest frequency observed in any of the main clinical studies.

Table 1: Adverse reactions reported in patients with MDS or AML treated with Azacitidine (clinical studies and post-marketing)

System organ class	Very common	Common	Uncommon	Rare	Not known
Infections and infestations	pneumonia* (including bacterial, viral and fungal), nasopharyngitis	sepsis* (including bacterial, viral and fungal), neutropenic sepsis*, respiratory tract infection (includes upper and bronchitis), urinary tract infection, cellulitis, diverticulitis, oral fungal infection, sinusitis, pharyngitis, rhinitis, herpes simplex, skin infection			necrotising fasciitis*
Neoplasms benign, malignant and unspecified (including cysts and polyps)					differentiation syndrome* ^a
Blood and lymphatic system disorders	febrile neutropenia*, neutropenia, leukopenia, thrombocytopenia, anaemia	Pancytopenia*, bone marrow failure			
Immune system disorders			hypersensitivity reactions		
Metabolism and nutrition disorders	anorexia, decreased appetite, hypokalemia	dehydration		tumour lysis syndrome	
Psychiatric disorders	Insomnia	confusional state, anxiety			
Nervous system disorders	Dizziness, headache	intracranial haemorrhage*, syncope, somnolence, lethargy			
Eye disorders		eye haemorrhage, conjunctival haemorrhage			
Cardiac diorders		pericardial effusion	pericarditis		
Vascular disorders		hypotension*, hypertension, orthostatic hypotension, haematoma			
Respiratory, thoracic and mediastinal disorders	dyspnoea, epistaxis	pleural effusion, dyspnoea exertional, pharyngolaryngeal pain		Interstitial lung disease	
Gastrointestinal disorders	diarrhoea, vomiting, constipation, nausea, abdominal pain (includes upper and abdominal discomfort)	gastrointestinal haemorrhage* (includes mouth haemorrhage), haemorrhoidal haemorrhage, stomatitis, gingival bleeding, dyspepsia			
Hepatobiliary disorders			hepatic failure*, progressive hepatic coma		

Black A/s: 350 x 720 mm

eugia Packaging Development		Product Name	Component	Item Code	Date & Time
		AZASQUARE	Leaflet	P1530284	29.11.2024 & 3:00 PM
		Customer / Country	Version No.	Reason Of Issue	Reviewed / Approved by
		Eugia_Pharma_Malaysia	08	NEW	
Team Leader	Narender	Dimensions	No. of Colours : 01		
Initiator	Sagar	350 x 720 mm			
Artist:	Vision Graphic Designers	Pharmacode	 30284		
		30284			
Additional Information :		Font Type: Times New Roman			Sign / Date
		Font Size: 10 pt			

Skin and subcutaneous tissue disorders	petechiae, pruritus (includes generalized), rash, ecchymosis	purpura, alopecia, urticaria, erythema, rash macular	acute febrile neutrophilic dermatosis, pyoderma gangrenosum		
Musculoskeletal and connective tissue disorders	arthralgia, musculoskeletal pain (includes back, bone and pain in extremity)	muscle spasms, myalgia			
Renal and urinary disorders		renal failure*, haematuria, elevated serum creatinine	renal tubular acidosis		
General disorders and administration site conditions	pyrexia*, fatigue, asthenia, chest pain, injection site erythema, injection site pain, injection site reaction (unspecified)	bruising, haematoma, induration, rash, pruritus, inflammation, discoloration, nodule and haemorrhage (at injection site), Malaise, chills, catheter site hemorrhage		injection site necrosis (at injection site)	
Investigations	Weight decreased				

* = rarely fatal cases have been reported
^ = see section *Special warnings and precautions for use*

Description of selected adverse reactions

Haematologic adverse reactions

The most commonly reported (≥ 10 %) haematological adverse reactions associated with azacytidine treatment include anaemia, thrombocytopenia, neutropenia, febrile neutropenia and leukopenia, and were usually Grade 3 or 4. There is a greater risk of these events occurring during the first 2 cycles, after which they occur with less frequency in patients with restoration of haematological function.

Most haematological adverse reactions were managed by routine monitoring of complete blood counts and delaying Azacitidine administration in the next cycle, prophylactic antibiotics and/or growth factor support (e.g., G-CSF) for neutropenia and transfusions for anaemia or thrombocytopenia as required.

Infections

Myelosuppression may lead to neutropenia and an increased risk of infection. Serious adverse reactions such as sepsis, including neutropenic sepsis and pneumonia were reported in patients receiving Azacitidine, some with a fatal outcome. Infections may be managed with the use of anti- infectives plus growth factor support (e.g., G-CSF) for neutropenia.

Bleeding

Bleeding may occur with patients receiving Azacitidine. Serious adverse reactions such as gastrointestinal haemorrhage and intracranial haemorrhage have been reported. Patients should be monitored for signs and symptoms of bleeding, particularly those with pre-existing or treatment related thrombocytopenia.

Hypersensitivity

Serious hypersensitivity reactions have been reported in patients receiving Azacitidine. In case of an anaphylactic-like reaction, treatment with Azacitidine should be immediately discontinued and appropriate symptomatic treatment initiated.

Skin and subcutaneous tissue adverse reactions

The majority of skin and subcutaneous adverse reactions were associated with the injection site. None of these adverse reactions led to discontinuation of Azacitidine, or reduction of Azacitidine dose in the pivotal study. The majority of adverse reactions occurred during the first 2 cycles of treatment and tended to decrease with subsequent cycles. Subcutaneous adverse reactions such as injection site rash/ inflammation/ pruritus, rash, erythema and skin lesion may require management with concomitant medicinal products, such as antihistamines, corticosteroids and non-steroidal anti- inflammatory medicinal products (NSAIDs). These cutaneous reactions have to be distinguished from soft tissue infections, sometimes occurring at injection site. Soft tissue infections, including cellulitis and necrotising fasciitis in rare cases leading to death, have been reported with Azacitidine in the post marketing setting. For clinical management of infectious adverse reactions, see section *Undesirable effects: Infections*.

Gastrointestinal adverse reactions

The most commonly reported gastrointestinal adverse reactions associated with Azacitidine treatment included constipation, diarrhoea, nausea and vomiting. These adverse reactions were managed symptomatically with anti-emetics for nausea and vomiting, anti-diarrhoeals for diarrhoea, and laxatives and/or stool softeners for constipation.

Renal adverse reactions

Renal abnormalities, ranging from elevated serum creatinine and haematuria to renal tubular acidosis, renal failure and death were reported in patients treated with Azacitidine (see section *Special warnings and precautions for use*).

Hepatic adverse reactions

Patients with extensive tumour burden due to metastatic disease have been reported to experience hepatic failure, progressive hepatic coma and death during Azacitidine treatment (see section *Special warnings and precautions for use*).

Cardiac events

Data from a clinical study allowing enrolment of patients with known history of cardiovascular or pulmonary disease showed an increase in cardiac events in patients with newly diagnosed AML treated with Azacitidine (see section *Special warnings and precautions for use*).

Overdosage

One case of overdose with Azacitidine was reported during clinical trials. A patient experienced diarrhoea, nausea, and vomiting after receiving a single intravenous dose of approximately 290 mg/m², almost 4 times the recommended starting dose.

In the event of overdose, the patient should be monitored with appropriate blood counts and should receive supportive treatment, as necessary. There is no known specific antidote for Azacitidine overdose.

Pharmacological properties

Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, Pyrimidine analogues; ATC code: L01BC07

Azacitidine is believed to exert its antineoplastic effects by multiple mechanisms including cytotoxicity on abnormal haematopoietic cells in the bone marrow and hypomethylation of DNA. The cytotoxic effects of Azacitidine may result from multiple mechanisms, including inhibition of DNA, RNA and protein synthesis, incorporation into RNA and DNA, and activation of DNA damage pathways. Non-proliferating cells are relatively insensitive to Azacitidine. Incorporation of Azacitidine into DNA results in the inactivation of DNA methyltransferases, leading to hypomethylation of DNA. DNA hypomethylation of aberrantly methylated genes involved in normal cell cycle regulation, differentiation and death pathways may result in gene re-expression and restoration of cancer- suppressing functions to cancer cells.

The relative importance of DNA hypomethylation versus cytotoxicity or other activities of Azacitidine to clinical outcomes has not been established.

Pharmacokinetic properties

Absorption

Following subcutaneous administration of a single 75 mg/m² dose, Azacitidine was rapidly absorbed with peak plasma concentrations of 750 ± 403 ng/mL occurring at 0.5 h after dosing (the first sampling point). The absolute bioavailability of Azacitidine after subcutaneous relative to intravenous administration (single 75 mg/m² doses) was approximately 89 % based on area under the curve (AUC). Area under the curve and maximum plasma concentration (Cmax) of subcutaneous administration of Azacitidine were approximately proportional within the 25 to 100 mg/m² dose range.

Distribution

Following intravenous administration, the mean volume of distribution was 76 ± 26 L, and systemic clearance was 147 ± 47 L/h.

Biotransformation

Based on in vitro data, Azacitidine metabolism does not appear to be mediated by cytochrome P450 isoenzymes (CYPs), UDP-glucuronosyltransferases (UGTs), sulfotransferases (SULTs), and glutathione transferases (GSTs).

Azacitidine undergoes spontaneous hydrolysis and deamination mediated by cytidine deaminase. In human liver S9 fractions, formation of metabolites was independent of NADPH implying that Azacitidine metabolism was not mediated by cytochrome P450 isoenzymes. An in vitro study of Azacitidine with cultured human hepatocytes indicates that at concentrations of 1.0 µM to 100 µM (i.e., up to approximately 30-fold higher than clinically achievable concentrations), Azacitidine does not induce CYP 1A2, 2C19, or 3A4 or 3A5. In studies to assess inhibition of a series of P450 isoenzymes (CYP 1A2, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1 and 3A4) Azacitidine up to 100 µM did not produce inhibition. Therefore, CYP enzyme induction or inhibition by Azacitidine at clinically achievable plasma concentrations is unlikely.

Elimination

Azacitidine is cleared rapidly from plasma with a mean elimination half-life (t½) after subcutaneous administration of 41 ± 8 minutes. No accumulation occurs after subcutaneous administration of 75 mg/m² Azacitidine once daily for 7 days. Urinary excretion is the primary route of elimination of Azacitidine and/or its metabolites. Following intravenous and subcutaneous administration of 14C- Azacitidine, 85 and 50 % of the administered radioactivity was recovered in urine respectively, while < 1 % was recovered in faeces.

Special populations

The effects of hepatic impairment (see section *Posology and method of administration*), gender, age, or race on the pharmacokinetics of Azacitidine have not been formally studied.

Renal impairment

Renal impairment has no major effect on the pharmacokinetic exposure of Azacitidine after single and multiple subcutaneous administrations. Following subcutaneous administration of a single 75 mg/m² dose, mean exposure values (AUC and Cmax) from subjects with mild, moderate and severe renal impairment were increased by 11-21 %, 15-27 %, and 41-66 %, respectively, compared to normal renal function subjects. However, exposure was within the same general range of exposures observed for subjects with normal renal function.

Azacitidine can be administered to patients with renal impairment without initial dose adjustment provided these patients are monitored for toxicity since Azacitidine and/or its metabolites are primarily excreted by the kidney.

Pharmacogenomics

The effect of known cytidine deaminase polymorphisms on Azacitidine metabolism has not been formally investigated.

Pharmaceutical particulars List of excipients

Mannitol

Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section *Special precautions for storage* and *Special precautions for disposal and other handling*.

Special precautions for storage

Unopened vial:

Store the medicinal product in its original package at below 30°C.

After reconstitution:

When AZASQUARE is reconstituted using water for injections that has not been refrigerated, chemical and physical in-use stability of the reconstituted medicinal product has been demonstrated at 25°C for 45 minutes and at 2°C to 8°C for 8 hours.

The shelf life of the reconstituted medicinal product can be extended by reconstituting with refrigerated (2°C to 8°C) water for injections. When AZASQUARE is reconstituted using refrigerated (2°C to 8°C) water for injections, the chemical and physical in-use stability of the reconstituted medicinal product has been demonstrated at 2°C to 8°C for 22 hours.

From a microbiological point of view, the reconstituted product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and must not be longer than 8 hours at 2°C to 8°C when reconstituted using water for injections that has not been refrigerated or not longer than 22 hours when reconstituted using refrigerated (2°C to 8°C) water for injections.

Shelf life

Unopened vial:

24 months

Nature and contents of container

30mL Type-I, clear transparent moulded glass vial stoppered with 20 mm (Lyo igloo), grey bromo butyl rubber stopper and sealed with 20 mm aluminium seal having white color polypropylene disc, containing 100mg of Azacitidine. Each vial is packed with protective plastic cover.

Pack size: 1 vial with package insert.

Special precautions for disposal and other handling

Recommendations for safe handling

AZASQUARE is a cytotoxic medicinal product and, as with other potentially toxic compounds, caution should be exercised when handling and preparing Azacitidine suspensions. Procedures for proper handling and disposal of anticancer medicinal products should be applied. If reconstituted Azacitidine comes into contact with the skin, immediately and thoroughly wash with soap and water. If it comes into contact with mucous membranes, flush thoroughly with water.

Reconstitution procedure

AZASQUARE should be reconstituted with water for injections. The shelf life of the reconstituted medicinal product can be extended by reconstituting with refrigerated (2°C to 8°C) water for injections. Details on storage of the reconstituted product are provided below.

- The following supplies should be assembled:
Vial(s) of Azacitidine; vial(s) of water for injections; non-sterile surgical gloves; alcohol wipes; 5 mL injection syringe(s) with needle(s).
- 4 mL of water for injections should be drawn into the syringe, making sure to purge any air trapped within the syringe.
- The needle of the syringe containing the 4 mL of water for injections should be inserted through the rubber top of the Azacitidine vial followed by injection of the water for injections into the vial.
- Following removal of the syringe and needle, the vial should be vigorously shaken until a uniform cloudy suspension is achieved. After reconstitution each mL of suspension will contain 25 mg of Azacitidine (100 mg/4 mL). The reconstituted product is a homogeneous, cloudy suspension, free of agglomerates. The product should be discarded if it contains large particles or agglomerates. Do not filter the suspension after reconstitution since this could remove the active substance. It must be taken into account that filters are present in some adaptors, spikes and closed systems; therefore, such systems should not be used for administration of the drug after reconstitution.
- The rubber top should be cleaned and a new syringe with needle inserted into the vial. The vial should then be turned upside down, making sure the needle tip is below the level of the liquid. The plunger should then be pulled back to withdraw the amount of medicinal product required for the proper dose, making sure to purge any air trapped within the syringe. The syringe with needle should then be removed from the vial and the needle disposed of.
- A fresh subcutaneous needle (recommended 25-gauge) should then be firmly attached to the syringe. The needle should not be purged prior to injection, in order to reduce the incidence of local injection site reactions.
- When more than 1 vial is needed all the above steps for preparation of the suspension should be repeated. For doses requiring more than 1 vial, the dose should be equally divided e.g., dose 150 mg = 6 mL, 2 syringes with 3 mL in each syringe. Due to retention in the vial and needle, it may not be feasible to withdraw all of the suspension from the vial.
- The contents of the dosing syringe must be re-suspended immediately prior to administration. The syringe filled with reconstituted suspension should be allowed up to 30 minutes prior to administration to reach a temperature of approximately 20°C-25°C. If the elapsed time is longer than 30 minutes, the suspension should be discarded appropriately, and a new dose prepared. To re-suspend, vigorously roll the syringe between the palms until a uniform, cloudy suspension is achieved. The product should be discarded if it contains large particles or agglomerates.

Storage of the reconstituted product

For storage conditions after reconstitution of the medicinal product, see section *Special precautions for storage*.

Calculation of an individual dose

The total dose, according to the body surface area (BSA) can be calculated as follows: Total dose (mg) = Dose (mg/ m²) x BSA (m²)

The following table is provided only as an example of how to calculate individual Azacitidine doses based on an average BSA value of 1.8 m².

Dose mg/ m² (% of recommended starting dose)	Total dose based on BSA value of 1.8 m²	Number of vials required	Total volume of reconstituted suspension required
75 mg/ m² (100%)	135 mg	2 vials	5.4 mL
37.5 mg/m² (50%)	67.5 mg	1 vial	2.7 mL
25 mg/m² (33%)	45 mg	1 vial	1.8 mL

Method of administration

Reconstituted AZASQUARE should be injected subcutaneously (insert the needle at a 45-90° angle) using a 25-gauge needle into the upper arm, thigh or abdomen.

Doses greater than 4 mL should be injected into two separate sites.

Injection sites should be rotated. New injections should be given at least 2.5 cm from the previous site and never into areas where the site is tender, bruised, red, or hardened.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Manufacturer

Eugia Pharma Specialities Limited, Unit-1, Survey No. 550, 551 and 552, Kolthur Village, Shamirpet Mandal, Medchal Malkajgiri District, Shamirpet, Telangana, 500101, India.

Product Registration Holder in Malaysia

Synerrv Sdn. Bhd. SO-29-2, Menara 1, KL Eco City, Jalan Bangsar Kg Haji Abdullah Hukum, 59200 Bangsar Wilayah Persekutuan, Kuala Lumpur, Malaysia

Leaflet Revision Date

28 November 2024