

Fasdol Suspension 250mg/5ml (Strawberry)

Fasdol Suspension 250mg/5ml (Orange)

Name and Strength of Active Substance

Paracetamol 250mg/5ml

It also contains preservative Methylparaben 0.1% w/v

Product Description:

Fasdol Suspension Strawberry: A pink rose coloured and strawberry flavour suspension.

Fasdol Suspension Orange: Orange coloured and orange flavour suspension

Pharmacodynamics

Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

Analgesic

The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

Antipyretic

Paracetamol act centrally on the hypothalamic heat regulating center to produce peripheral vasodilatation resulting in increased blood flow through the skin, sweating and heat loss. Paracetamol reduces fever by inhibiting the formation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermoregulatory center.

Pharmacokinetics

Following oral administration paracetamol is rapidly absorbed. Paracetamol absorption takes place mainly in the small intestine and therefore the rate of absorption is depending on the rate of gastric emptying. It has been shown that drugs which delay gastric emptying also delay the absorption of paracetamol whereas metoclopramide (a drug which increases the rate of gastric emptying) accelerates absorption of the analgesic through the total amount absorbed doses not increase.

The presence of food in the stomach has also been reported to reduce the rate of absorption of paracetamol. Alterations in gastric pH have no appreciable effect on paracetamol absorption. During absorption, the amount of paracetamol which is inactivated is negligible and it has been shown that paracetamol does not affect gastric mucosal permeability and does not produce mucosal bleeding.

Peak plasma concentrations are reached 1 hour after absorption. The plasma half-life is 1 to 3 hours.

Paracetamol penetrates the brains and is present in breast milk of human.

Paracetamol is metabolized by the microsomal enzyme system of the liver. This metabolism is mainly to the glucuronide and sulphate conjugates, accounting for approximately 49% and 26% of the ingested dose respectively. About 4% is excreted as free paracetamol. Other minor pathways include the production of catechol derivatives and cysteine conjugates (via glutathione).

Paracetamol excretion is rapid and occurs via the urine.

Indication

Indicated for the relief of fever, headache and symptoms of cold and flu, toothache, discomfort of teething and fever after vaccination.

Recommend Doses

Children: 1-3 years: 3 ml – 5 ml

4-6 years: 5 ml – 7 ml

7-9 years: 7 ml – 10 ml

10-12 years: 10 ml – 15 ml

Shake well before use. Do not give more than 4 times daily.

Mode of Administration

Oral

Contraindications

Hypersensitivity to paracetamol or any of the other ingredients/components of the product.

Severe and active hepatic impairment.

Warning and Precautions

**This preparation contains PARACETAMOL.
Do not take any other paracetamol
containing medicines at the same time.**

- Keep out of reach of children.
- Do not take if allergic to paracetamol.
- Patients should contact their health care provider if symptom persists (if the pain lasts for more than 10 days, If there is redness or fever lasts more than 3 days).
- Paracetamol should be given with care to patients with impaired kidney or liver function.
- Large doses should be avoided in patients with hepatic impairment. Paracetamol overdose may harm the liver.
- Do not exceed recommended dose.
- Paracetamol provides symptomatic relief only, additional therapy to treat the cause of the pain or fever should be instituted when necessary.
- Allergy alert: Paracetamol may cause severe skin reactions, symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Effects on Ability to Drive and Use Machines

It is unlikely to impair a patient's ability to drive or use machinery.

Interaction with Other Medicaments

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Statement on Using During Pregnancy and Lactation

Use in pregnancy:

- Considered to be the analgesic of choice in pregnant patients.
- Although it crosses placenta, paracetamol is considered to be safe in normal therapeutic doses for short-term use as a minor analgesic / antipyretic in pregnancy.

Use in lactation:

- Excreted in breast milk.
- Maternal ingestion of paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

Adverse Effects / Undesirable Effects

Adverse effects of paracetamol are rare and usually mild, although haematological reactions have been reported. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Overdose and treatment

Symptoms:

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse of acute overdose of paracetamol is a dose-dependent, potentially fatal hepatic necrosis. Clinical and laboratory evidence of hepatotoxicity may be delayed for up to one week. Major manifestations of liver failure such as jaundice, hypoglycaemia and metabolic acidosis may take at least 3 days to develop.

Treatment:

On cases of overdose, methods of reducing the absorption of ingested drug are important. Gastric lavage is essential even if several hours have elapsed. Prompt administration of 50g activated charcoal and 500ml iced mannitol 20% by mouth, may reduce absorption. If the history suggests that 15g Paracetamol or more has been ingested, administer one of the following antidotes:

Acetylcysteine 20% i.v.: Administer intravenously, 20% acetylcysteine (Parvolex, Glaxo) immediately without waiting for positive urine test or plasma level results: initial dose of 150mg/kg over 15 minutes, followed by continuous infusion of mg/kg in 500ml 5% glucose/dextrose over hours and mg/kg in L 5% glucose/dextrose over 16 hours;

OR

Oral Methionine: 2.5g immediately followed by three further doses of 2.5g at four hourly intervals. For a 3 years old child, 1g methionine every four hours for four does has been used;

OR

Oral Acetylcysteine 5%: 140mg/kg as a loading dose, then 70mg/kg every 4 hours for a total of 17 maintenance doses. If more than ten hours have elapsed since the over dosage was taken, the antidote may be ineffective.

Storage Condition

Store below 30°C.

Shelf Life

3 years from the date of manufacture.

Dosage Forms and Packaging Available

Amber PET 90 ml.

Product registration holder:

Advance Pharma Sdn Bhd 539885-W
28, Jalan BP6/6, Bandar Bukit Puchong,
47120 Puchong, Selangor.

Manufacturer

AV MANUFACTURING SDN. BHD. 667760-V
Lot 10621 (PT16700). JLN Permata 2,
Arab Malaysian Industrial Park,
71800 Nilai, Negeri Sembilan.

Date of Revision

Aug 2019

Ver.01